AN EFFICIENT CONSTRUCTION OF 1,2-*TRANS*- β -GLYCOSIDIC LINKAGES VIA BENZYL-PROTECTED GLYCOPYRANOSYL *P,P*-DIPHENYL-*N*-(*p*-TOLUENESULFONYL)-PHOSPHINIMIDATES

Shun-ichi HASHIMOTO, Takeshi HONDA, and Shiro IKEGAMI*

Faculty of Pharmaceutical Sciences, Teikyo University, Sagamiko, Kanagawa 199-01, Japan

An efficient 1,2-*trans*-glycosidation reaction without neighbouring group participation has been developed using benzyl-protected *P,P*-diphenyl-*N*-(*p*-toluenesulfonyl)phosphinimides as glycosyl donors in the presence of trimethylsilyl triflate or boron trifluoride etherate.

KEYWORDS glycosidation; 1,2-*trans*-glycosidic linkage; phosphinimide; steroidal glycoside

Due to the increasingly recognized biological significance of saccharide residues of carbohydrate-containing biomolecules, the development of expeditious methods for the stereocontrolled construction of the glycosidic linkages is becoming of paramount importance in synthetic organic chemistry.^{1,2)} Our interest in this area has led to the design of the leaving groups of glycosyl donors allowing excellent shelf-lives, as well as their activation without resorting to precious, explosive, or toxic heavy-metal salts as promoters. In an effort to capitalize on the phosphorus-containing leaving groups, we recently developed rapid and efficient methods for the stereocontrolled construction of 1,2-*trans*- β -glycosidic linkages by the devise of shelf-stable benzyl- or benzoyl-protected glycopyranosyl diphenyl phosphates³⁾ or benzoyl-protected *P,P*-diphenyl-*N*-(*p*-toluenesulfonyl)phosphinimides⁴⁾ as glycosyl donors, in which the respective promoter of choice was trimethylsilyl triflate (TMSOTf) or BF₃·OEt₂. As an extension of the phosphinimide method, we now turned our attention to the glycosidation using benzyl-protected glycopyranosyl *P,P*-diphenyl-*N*-(*p*-toluenesulfonyl)phosphinimides as glycosyl donors, some features of which are described herein.

Benzyl-protected glycopyranosyl phosphinimides (**1** and **2**) were readily prepared by condensation of the 1-*O*-lithium salts of 2,3,4,6-tetra-*O*-benzyl- α -D-glucopyranose and - α -D-galactopyranose with *P,P*-diphenyl-*N*-(*p*-toluenesulfonyl)phosphinimide chloride⁵⁾ (THF, -78 °C, 0.5 h). These donors could be stored in the freezer (at -20 °C) for at least several weeks without detectable deterioration.

Initial attempts at the glycosidation reactions were performed with suitably protected glycoside alcohols as acceptors. After surveying a variety of conditions, coupling of the phosphinimides (**1** and **2**) (1.1 eq) with alcohols (**3-8**) (1.0 eq) in propionitrile in the presence of TMSOTf (1.2 eq) at -55 °C was found to proceed smoothly to give the 1,2-*trans*- β -linked disaccharides in high yields and with high levels of stereoselectivity (method A, entries 1-6 in Table I). The combined use of TMSOTf as promoter and propionitrile as solvent proved to be the best choice for allowing considerable levels of 1,2-*trans*-selectivity. It should be noted that the alcohols bearing the acid-labile epoxy or acetal groups are safely glycosylated (entries 3, 4, and 6). From the result that the glycosidation of a 33:67 mixture⁶⁾ of **1** and its β -anomer under the above conditions gave essentially the same yields and stereoselectivities as that of **1**, the glycosidation reaction apparently proceeds through the thermodynamically more stable α -D-glycopyranosyl triflate followed by the backside attack of alcohols on this intermediate.

Table I. 1,2-*trans*-Glycosidation of Benzyl-Protected Glycopyranosyl *P,P*-Diphenyl-*N*-(*p*-toluenesulfonyl)-phosphinimides (**1** and **2**)

Entry	Donor	Acceptor	Method a)	Time (h)	Yield (%) ^{b)}	$\alpha:\beta$ ^{c)}	$\delta^{13}\text{C}$ ^{d)}
1	1	3	A	0.3	88	4:96	103.8(97.3)
2	1	5	A	0.5	84	12:88	102.4(96.6)
3	1	6	A	0.5	82	8:92	103.7(95.5)
4	1	7	A	0.5	79	13:87	103.2(96.9)
5	2	4	A	0.3	84	9:91	104.3(98.0)
6	2	8	A	0.3	82	8:92	104.7(97.6)
7	1	9	B	6	74	10:90	102.0(94.7)
8	1	10	B	6	72	13:87	102.3(94.7) ^{e)}
9	1	11	B	6	72	7:93	101.0(94.8)
10	1	12	B	8	70	12:88	101.9(94.8)
11	2	11	B	6	73	11:89	101.1(95.4)

a) Method A: TMSOTf (1.0 M in CH_2Cl_2 , 0.12 mmol) was added to a solution of donor (0.11 mmol) and acceptor (0.1 mmol) in propionitrile (2 ml) at -70°C , and the reaction was performed at -55°C under argon atmosphere. Method B: $\text{BF}_3\cdot\text{OEt}_2$ (1.0 M in CH_2Cl_2 , 0.11 mmol) was added to a mixture of donor (0.11 mmol), acceptor (0.1 mmol), and pulverized 4Å molecular sieves (50 mg) in CH_2Cl_2 (2.5 ml) at -23°C , and the reaction was carried out at this temperature in a sealed tube. After the reaction was complete, the product was isolated by extractive workup and subsequent chromatography on silica gel. b) Isolated total yield based on the acceptor used. c) Determined by HPLC analysis. d) Chemical shifts (δ ppm) in ^{13}C -NMR spectrum (100 MHz, CDCl_3) for the anomeric centers newly formed. Values in parentheses correspond to those of 1,2-*cis*-linked glycosides and disaccharides. e) Ref. 9.

On these positive notes, we then directed our efforts to the extension of this protocol to the construction of steroidal glycosides, with which the phosphate method³⁾ without neighbouring group participation proved to be less than satisfactory.⁷⁾ While the foregoing procedure (method A) also met with less success,⁸⁾ it was found that condensation of the phosphinimides (**1** and **2**) (1.1 eq) with steroidal alcohols (**9-12**) (1.0 eq) in dichloromethane in the presence of BF₃·OEt₂ (1.1 eq) and pulverized 4Å molecular sieves at -23 °C led to the preponderant formation of 1,2-*trans*-β-linked steroidal glycosides in good yields (method B, entries 7-11 in Table I).¹⁰⁾ In view of the lack of versatile methods for 1,2-*trans*-glycosidation of steroidal glycosides without the anchimeric assistance by the *O*-2-acyl group,¹¹⁾ our procedure should be of great promise.

In summary, the effectiveness of the benzyl-protected glycopyranosyl phosphinimides as glycosyl donors has been demonstrated by proper choice of promoters and solvents. More complex application of the glycosidation method are currently under investigation.

ACKNOWLEDGEMENT We are grateful to Misses Y. Ikenoya and H. Ishido for spectroscopic measurements. Partial financial support for this research from the Japan Research Foundation for Optically Active Compounds, the Japan Science Society, and the Ministry of Education, Science and Culture of Japan is gratefully acknowledged.

REFERENCES AND NOTES

- 1) For recent reviews, see: a) H. Paulsen, *Angew. Chem. Int. Ed. Engl.*, **21**, 155 (1982); b) R. R. Schmidt, *ibid.*, **25**, 212 (1986); c) H. Kunz, *ibid.*, **26**, 294 (1987).
- 2) For other more recent works, see: a) F. Dasgupta and P. J. Garegg, *Carbohydr. Res.*, **177**, C13 (1988); b) Y. Ito and T. Ogawa, *Tetrahedron Lett.*, **29**, 1061 (1988); c) M. Nishizawa, Y. Kan, and H. Yamada, *ibid.*, **29**, 4597 (1988); d) D. R. Mootoo, P. Konradsson, U. Udodong, and B. Fraser-Reid, *J. Am. Chem. Soc.*, **110**, 5583 (1988); e) G. V. Reddy, V. R. Kulkarni, and H. B. Mereyala, *Tetrahedron Lett.*, **30**, 4283 (1989); f) K. Suzuki, H. Maeta, and T. Matsumoto, *ibid.*, **30**, 4853 (1989); g) N. K. Kochetkov, E. M. Klimov, and N. N. Malysheva, *ibid.*, **30**, 5459 (1989); h) A. G. M. Barrett, B. C. B. Bezuidenhout, A. F. Gasielki, A. R. Howell, and M. A. Russell, *J. Am. Chem. Soc.*, **111**, 1392 (1989); i) D. Kahne, S. Walker, Y. Cheng, and D. V. Engen, *ibid.*, **111**, 6881 (1989); j) G. H. Veeneman and J. H. van Boom, *Tetrahedron Lett.*, **31**, 275 (1990); k) P. Konradsson, D. R. Mootoo, R. E. McDevitt, and B. Fraser-Reid, *J. Chem. Soc., Chem. Commun.*, **1990**, 270.
- 3) S. Hashimoto, T. Honda, and S. Ikegami, *J. Chem. Soc., Chem. Commun.*, **1989**, 685.
- 4) S. Hashimoto, T. Honda, and S. Ikegami, *Heterocycles*, **30**, 775 (1990).
- 5) H. Bock and W. Wiegand, *Chem. Ber.*, **99**, 1068 (1966).
- 6) An anomeric mixture of the phosphinimide was prepared by phosphinimidation of the 1-*O*-thallium(I) salt under Perlin's conditions. A. Granata and A. S. Perlin, *Carbohydr. Res.*, **94**, 165 (1981).
- 7) Glycosidation of 2,3,4,6-tetra-*O*-benzyl-α-D-glucopyranosyl diphenyl phosphate (1.1 eq) with **9** (1.0 eq) gave the glycoside with the α:β ratio of 10:90 in 17% yield.
- 8) Glycosidation of **1** with **9** resulted in the α:β ratio of 12:88 and 20% yield.
- 9) G. Wulff, U. Schröder, and J. Wichelhaus, *Carbohydr. Res.*, **72**, 280 (1979).
- 10) Glycosidation of a 33:67 mixture of **1** and its β-anomer with **9** under the similar conditions furnished the glycoside with the α:β ratio of 45:55 in 75% yield.
- 11) a) S. Shoda and T. Mukaiyama, *Chem. Lett.*, **1979**, 847; b) G. Wulff and J. Wichelhaus, *Chem. Ber.*, **112**, 2847 (1979); d) S. Koto, T. Sato, N. Morishima, and S. Zen, *Bull. Chem. Soc. Jpn.*, **53**, 1761 (1980); e) S. Hashimoto, M. Hayashi, and R. Noyori, *Tetrahedron Lett.*, **25**, 1379 (1984); f) R. R. Schmidt and J. Michel, *J. Carbohydr. Chem.*, **4**, 141 (1985).

(Received May 30, 1990)