TODDACOUMAQUINONE, A UNIQUE COUMARIN-NAPHTHOQUINONE DIMER, FROM *TODDALIA ASIATICA* (L.) LAM. (*T. ACULEATA* PERS.)

Hisashi ISHII*,^a Jun-Ichi KOBAYASHI,^a Hiroko SEKI^b and Tsutomu ISHIKAWA*,^a

Faculty of Pharmaceutical Sciences^a and Chemical Analysis Center,^b Chiba University, 1-33 Yayoi-cho, Inage, Chiba 263, Japan

The spectroscopic structural elucidation of toddacoumaquinone (2), a unique coumarin-naphthoquinone dimer, isolated from *Toddalia asiatica* (L.) Lam. (*T. aculeata* Pers.) is described.

KEYWORDS coumarin-naphthoquinone dimer; structural establishment; 2D-NMR; *Toddalia asiatica* (*T. aculeata*); Rutaceae

Recently we¹⁾ described the structural determination of toddacoumalone (1), a novel mixed dimer of coumarin and quinolone which had been reported as unknown I,²⁾ from *Toddalia asiatica* (L.) Lam. (*T. aculeata* Pers.). In this paper we present the structural elucidation of unknown II,²⁾ a unique coumarin-naphthoquinone dimer designated as toddacoumaquinone (2), by spectroscopic means.

Toddacoumaquinone (2) was isolated as orange prisms, mp 278–281 °C (AcOEt), in 0.015 % yield in a racemic form.³⁾ The molecular formula of C₂₃H₁₈O₇ (M⁺ 406.1052. Calcd 406.1052) was confirmed by high resolution mass spectrometry. The spectral data [$\nu_{\max}$ (KBr): 1730 and 1620 cm⁻¹; $\lambda_{\max}$ (MeOH): 201 (log ϵ 4.76), 224sh (4.39), 249sh (4.40), 256 (4.43), 283 (4.36) and 327 (4.20) nm; NMR (see Table I)] indicated the presence of a 8-substituted 5, 7-dimethoxycoumarin moiety in 2 like 1.¹⁾

On the other hand, a *p*-naphthoquinone⁴⁾ unit in the molecule was deduced by additional carbonyl bands ($\nu_{\max}$: 1683 and 1650 cm⁻¹) in the IR spectrum, the lowest frequency π – π^* transition of the UV absorption at 327 nm and the two lowest field shifted signals due to carbonyl functions at around 180 ppm in the ¹³C-NMR spectrum. The following facts allowed us to extend the unit to an 8-substituted 2-methoxy-6-methylnaphthoquinone: 1. Two 3H singlets attributable to methyl and methoxy groups appeared at δ 2.50 and 3.80 in the ¹H-NMR spectrum. 2. The NOE enhancements were observed between the methoxy group and a quinoid proton (δ 6.10) and between the methyl group and a pair of *meta* coupled aromatic protons (δ 7.29 and 8.00). 3. Two carbons at ring junction, 4'a-C and 8'a-C, showed three independent bond correlations to the quinoid and to the aromatic protons in the COLOC experiments (see Table I).

Thus, toddacoumaquinone should be a biaryl compound consisting of a coumarin and a naphthoquinone, the structure of which could be depicted as 2. It could be reasonable that 2 would be synthesized in a plant body through [4+2] cycloaddition reaction between the coumarin diene (3) and a *p*-benzoquinone derivative because of the coexistence^{1, 2)} of formal Diels-Alder dimeric products derived from 3. The synthetic work of 2 based on this biogenetic consideration is now in progress.

Table I. NMR Data^a of Toddacoumaquinone (2)

C# ^b	δ_H	NOE	δ_C	COLOC	
				J=8Hz	J=4Hz
2	—		161.34	4-H(3)	4-H(3), 3-H(2)
3	6.10(d, J=9.6Hz)	4-H	110.95		
4	8.03(d, J=9.6Hz)		138.82		
4a	—		103.89	3-H(3), 6-H(3)	
5	—		156.72	5-OMe(3), 6-H(2)	5-OMe(3), 6-H(2)
6	6.41 (s)	5-OMe, 7-OMe	90.45		
7	—		160.33	7-OMe(3), 6-H(2)	7-OMe(3), 6-H(2)
8	—		110.30	6-H(3), 7'-H(3)	6-H(3), 7'-H(3)
8a	—		152.78	4-H(3)	4-H(3)
5-OMe	3.99(s)	4-H, 6-H	56.03		
7-OMe	3.77(s)	6-H	56.13		
1'	—		185.21		5'-H(4)
2'	—		160.64	2'-OMe(3), 3'-H(2)	2'-OMe(3), 3'-H(2)
3'	6.10(s)	2'-OMe	108.60		2'-OMe(4)
4'	—		179.73	3'-H(2)	
4'a	—		133.50	3'-H(3)	3'-H(3)
5'	8.00(d, J=1.1Hz)	6'-Me	127.40	6'-Me(3)	6'-Me(3)
6'	—		144.77	6'-Me(2)	6'-Me(2)
7'	7.29(d, J=1.1Hz)	6'-Me	138.63	6'-Me(3)	6'-Me(3)
8'	—		133.14		6-H(4)
8'a	—		127.11	5'-H(3), 7'-H(3)	
2'-OMe	3.80(s)	3'-H	56.29		
6'-Me	2.50(s)	5'-H, 7'-H	21.86	5'-H(3), 7'-H(3)	5'-H(3), 7'-H(3)

^a ¹H-NMR (500 MHz in CDCl₃) are reported downfield from internal TMS at 0.00 ppm. ¹³C-NMR assignments are related to internal CDCl₃ at 77.00 ppm. ¹H- and ¹³C-NMR assignments are based on H-C COSY, DEPT and COLOC experiments. For the COLOC experiments the number in parentheses denotes the number of bonds involved in the correlation. ^b The number is arbitrarily shown in each structure.

ACKNOWLEDGEMENTS We thank Prof. M. Suzuki and Dr. K.-I. Harada, Meijo University, for measurement of mass spectrometry.

REFERENCES AND NOTES

- 1) H. Ishii, J.-I. Kobayashi and T. Ishikawa, *Tetrahedron Lett.*, **32**, 6907(1991).
- 2) H. Ishii, J.-I. Kobayashi, M. Ishikawa, J. Haginiwa and T. Ishikawa, *Yakugaku Zasshi*, **111**, 365(1991).
- 3) No optical rotation or Cotton effect was observed at range from 400 to 240 nm in the ORD and CD spectra (MeOH).
- 4) S. Berger, P. Hertl and A. Rieker, "The Chemistry of the Quinonoid Compounds," Vol. 2, ed. by S. Pathai and Z. Rappoport, John Wiley and Sons, Inc., New York, 1988, p. 29.

(Received April 3, 1992)