TOTAL SYNTHESIS OF NATURAL (+)-DUOCARMYCIN SA

Hideaki MURATAKE, Naoshige MATSUMURA, and Mitsutaka NATSUME*

Research Foundation Itsuu Laboratory, 2-28-10 Tamagawa, Setagaya-ku, Tokyo 158, Japan

A total synthesis of natural (+)-duocarmycin SA (**1**) was achieved as shown in Chart 1, starting from L-malic acid (**5**) by using a Lewis acid-mediated indole formation reaction of a pyrrole precursor **14** to form the key compound **15**.

KEY WORDS duocarmycin SA; antitumor antibiotic; total synthesis; indole formation reaction

Last year, we reported a 12-step total synthesis of duocarmycin SA (**1**),¹⁾ a potent antitumor antibiotic isolated from a culture broth of the *Streptomyces* species,²⁾ in the racemic form by employing a new strategy (**2** → **3**) for construction of the *N*-acylcyclopropanoindolinone unit (dashed line in **3**), which plays a pivotal role in exerting extremely potent biological activities.³⁾ In contrast to the previous syntheses of **1**,⁴⁾ duocarmycin A⁵⁾ and CC-1065⁶⁾ using **4** as a synthetic intermediate, our adoption of **2** made it easy to design an enantioselective access to **2**, and here we report a total synthesis of natural (+)-duocarmycin SA (**1**).

Our synthetic plan was to select L-malic acid (**5**) for the starting material, and this was converted into an indole formation precursor **14**. Our procedure⁷⁾ was applied to **14** for the synthesis of a hydroxyindole to finally identify appropriate reaction conditions affording **15**. Succeeding operations, *i.e.*, transformation of the carboxylic acid into a nitrogen function (**17** → **18**) and subsequent construction of the piperidine ring (**19** → **20**), would provide **2** having the hydroxyl group of the requisite absolute configuration.

The synthesis was started with diol **9**, prepared from **5** according to the literature.⁸⁾ Its dibenzyl ether **10** was reduced with diisobutylaluminum hydride (DIBAL),⁹⁾ and the resulting aldehyde **11** was condensed with an anion derived from benzyl acetate to form **12**,¹⁰⁾ which was oxidized to the β-keto ester **13**. Coupling of **13** with the pyrrole part **8**, prepared from **6**¹¹⁾ by way of **7**, afforded a cyclization precursor **14**. Indole formation was accomplished by treatment of **14** with 2-ethyl-2-methyl-1,3-dioxane (50 mol eq) in CH₂Cl₂ in the presence of BF₃·OEt₂ (6 mol eq) at room temperature for 44 h, and **15** was produced in 54% yield, along with a furan

* To whom correspondence should be addressed.

derivative **27** (10% yield) as an inevitable by-product. The hydroxypropyl function in **15** was conveniently used as a protecting group of the phenol function. The acetate **16**, prepared from **15**, was partially hydrogenated with Pd-C in EtOAc in particular to give mostly a mono-debenzylated acid **17** for the Curtius rearrangement. This reaction with diphenylphosphoryl azide (DPPA)¹² had to be carried out with some caution, and a modified procedure¹³ for the reaction of **17** with DPPA and iso-Pr₂NEt in benzene, followed by treatment with MeOH, afforded **18** in a good yield.

The benzyl group at the primary alcohol in **18** was predominantly eliminated with BBr₃, and the resulting **19** was treated under Mitsunobu conditions.¹⁴ Clean cyclization to the piperidine ring was accomplished in a very good yield to afford **20**. The phenol protecting group was removed by i) methanolysis of the acetate, ii) Dess-Martin oxidation and iii) β -elimination with Et₃N to produce **22**, whose benzyl group was taken off by hydrogenation over Pearlman's catalyst in MeOH. Formation of the cyclopropanoindolinone **24** from **23** was

a: Cl₃CC(=NH)OBn, CF₃SO₃H, cyclohexane-CH₂Cl₂, 0-26°C, 52%. *b*: DIBAL, PhMe, -76 – -63°C, 87%. *c*: LDA + CH₃COOBn in THF, -71 – -66°C, 78%. *d*: Dess-Martin periodinane, CH₂Cl₂, reflux, 86%. *e*: *tert*-BuOK, THF, 0 – 23°C, 86%. *f*: 2-ethyl-2-methyl-1,3-dioxane, BF₃·OEt₂, CH₂Cl₂, 20°C, 54%. *g*: Ac₂O, pyridine, CH₂Cl₂, 16°C, 97%. *h*: H₂, 10% Pd-C, EtOAc, 13°C, 75%. *i*: i) DPPA, iso-Pr₂NEt, benzene, reflux; ii) MeOH, reflux, 69%. *j*: BBr₃, CH₂Cl₂, -80 – -55°C, 61%. *k*: DEAD, Ph₃P, THF, 21°C, 90%. *l*: K₂CO₃, MeOH, 21°C, 98%. *m*: i) Dess-Martin periodinane, CH₂Cl₂, reflux; ii) Et₃N, CH₂Cl₂, reflux, 86%. *n*: H₂ (5 atm), 20% Pd(OH)₂-C, MeOH, 25°C, 96%. *o*: *N*-piperidyl-CO-N=N-CO-*N*-piperidyl, *n*-Bu₃P, THF, 20°C, 88%. *p*: K₂CO₃, MeOH, 18°C, 96%. *q*: i) NaH, DMF-THF; ii) **26**, 0°C, 74%.

Chart 1

effected with 1,1'-(azodicarbonyl)dipiperidine¹⁵⁾ as in the case of the racemic series.¹⁾ Methanolysis of **24**, followed by condensation of **25** with 1-(5,6,7-trimethoxyindole-2-carbonyl)imidazole (**26**), completed a total synthesis of natural (+)-duocarmycin SA (**1**), {[α]_D²¹ +192° (*c* = 0.352, MeOH); lit.^{2a)}: [α]_D²⁴ +180° (*c* = 0.1, MeOH)}.

In summary, a novel indole formation reaction was applied to **14** for construction of the important intermediate **15** carrying in its side chain the requisite chiral center derived from L-malic acid. Mitsunobu reaction was successfully applied not only to the cyclopropanoindolinone formation (**23** → **24**) but also to dehydrative bond connection between the methyl carbamate group and the primary alcohol to give the *N*-methoxycarbonylpiperidine ring system (**19** → **20**). Using these reactions in key steps, a total synthesis of natural (+)-duocarmycin SA (**1**) was achieved starting from L-malic acid (**5**) and a pyrrole derivative **6**.

ACKNOWLEDGMENT Thanks are due to the Ministry of Education, Science, and Culture of Japan for financial support.

REFERENCES AND NOTES

- 1) Muratake H., Abe I., Natsume M., *Tetrahedron Lett.*, **35**, 2573 — 2576 (1994).
- 2) a) Ichimura M., Ogawa T., Takahashi K., Kobayashi E., Kawamoto I., Yasuzawa T., Takahashi I., Nakano H., *J. Antibiot.*, **43**, 1037 — 1038 (1990). b) Yasuzawa T., Saitoh Y., Ichimura M., Takahashi I., Sano H., *J. Antibiot.*, **44**, 445 — 447 (1991). c) Ichimura M., Ogawa T., Katsumata S., Takahashi K., Takahashi I., Nakano H., *J. Antibiot.*, **44**, 1045—1053 (1991).
- 3) a) Hurley L. H., Needham-VanDevanter D. R., *Acc. Chem. Res.*, **19**, 230 — 237 (1986); and literatures cited therein. b) Boger D. L., *Acc. Chem. Res.*, **28**, 20 — 29 (1995).
- 4) a) Boger D. L., Machiya K., *J. Am. Chem. Soc.*, **114**, 10056 — 10058 (1992). b) Boger D. L., Machiya K., Hertzog D. L., Kitos P. A., Holmes D., *J. Am. Chem. Soc.*, **115**, 9025 — 9036 (1993).
- 5) a) Fukuda Y., Nakatani K., Ito Y., Terashima S., *Tetrahedron Lett.*, **31**, 6699 — 6702 (1990). b) Fukuda Y., Nakatani K., Terashima S., *Bioorg. Med. Chem. Lett.*, **2**, 755 — 758 (1992).
- 6) Kelly R. C., Gebhard I., Wicnienski N., Aristoff P. A., Johnson P. D., Martin D. G., *J. Am. Chem. Soc.*, **109**, 6837 — 6838 (1987).
- 7) Fuji M., Muratake H., Natsume M., *Chem. Pharm. Bull.*, **40**, 2344 — 2352 (1992).
- 8) Saito S., Hasegawa T., Inaba M., Nishida R., Fujii T., Nomizu S., Moriwake T., *Chem. Lett.*, **1984**, 1389 — 1392.
- 9) An over-reduction product was obtained in 7% yield as a by-product; this was oxidized back to **11** with Dess-Martin reagent in 75% yield.
- 10) Benzyl (5*ξ*, 7*S*)-7,8-dibenzyloxy-5-hydroxy-3-oxooctanoate was obtained in 10% yield as a by-product.
- 11) Anderson H. J., Huang C. W., *Can. J. Chem.*, **45**, 897 — 902 (1967).
- 12) Ninomiya K., Shioiri T., Yamada S., *Tetrahedron*, **30**, 2151 — 2157 (1974).
- 13) Actual experimental details will be described in the forthcoming full paper.
- 14) Mitsunobu O., *Synthesis*, **1981**, 1 — 28.
- 15) Tsunoda T., Yamamiya Y., Itô S., *Tetrahedron Lett.*, **34**, 1639 — 1642 (1993).

(Received April 12, 1995; accepted May 12, 1995)