ASYMMETRIC SYNTHESIS OF (*S*)-(+)- AND (*R*)-(-)-NZ-105 THROUGH THE MODIFIED MICHAELIS-ARBUZOV REARRANGEMENT AS A KEY STEP

Tatsuhisa KATO, Mamiko TEJIMA, Hirosato EBIKE and Kazuo ACHIWA*

School of Pharmaceutical Sciences, University of Shizuoka 52-1 Yada, Shizuoka 422, Japan

The asymmetric synthesis of the (*S*)-(+)- and (*R*)-(-)-NZ-105 from the prochiral compound (**1**) was realized by using the modified Hunsdiecker reaction followed by the modified Michaelis-Arbuzov reaction with zerovalent palladium.

KEY WORDS 1,4-dihydropyridine; calcium antagonist; NZ-105; Hunsdiecker reaction; Michaelis-Arbuzov reaction

4-Aryl-1,4-dihydro-2,6-dimethylpyridine-3,5-dicarboxylate derivatives show calcium antagonistic activity and are therapeutically useful vasodilators.¹⁾ These calcium antagonists have a high affinity with the voltage-dependent calcium channel. Recently, a novel and structurally unique calcium antagonist (NZ-105 : 2-[Benzyl(phenyl)amino]ethyl 5-(5,5-Dimethyl-2-oxo-1,3,2-dioxaphosphorinane-2-yl)-1,4-dihydro-2,6-dimethyl-4-(3-nitrophenyl)-3-pyridinecarboxylate) was prepared.^{2,3)} The very slow onset and recovery from NZ-105-induced vasodilation may be attributable to the slow and long-lasting inhibition of transmembrane calcium uptake which accompanies its very slow binding to and dissociation from DHP receptors.⁴⁾ Some of these 1,4-dihydropyridines have a chiral center at 4-position, and these enantiomers have been reported to show different biological activities; for example, (*S*)-NZ-105 exhibited no less than 850-fold calcium antagonistic activity compared with that of *R* isomer.²⁾ But these racemic 1,4-dihydropyridines have been used as vasodilators. NZ-105 so far has been synthesized as racemate³⁾ or has been enantiomerically synthesized with separation via diastereomeric ester²⁾ or chemoenzymatic separation.⁵⁾ We describe here the first asymmetric synthesis of (*S*)- and (*R*)-NZ-105, which includes a brominating process through the modified Hunsdiecker reaction and a coupling of the phosphorous to an sp^2 carbon atom.

The Michaelis-Arbuzov rearrangement,⁶⁾ also known as the Arbuzov reaction, is one of the most extensively investigated in organophosphorous chemistry and is widely used to prepare phosphonates, phosphinates and phosphine oxides.⁷⁾ To our knowledge, there is no report in the literature on a Michaelis-Arbuzov rearrangement applying to 1,4-dihydropyridines.

The difficulty of nucleophilic substitution at an sp^2 carbon atom by conventional organic techniques is overcome by using transition metals.⁸⁾ It has been reported that zerovalent palladium is an effective catalyst for cross-coupling of alkenyl halides with nucleophiles.⁹⁾ Adaptations of this palladium-catalyzed reaction to the Arbuzov reaction have also been reported.¹⁰⁾ The phosphorous-carbon bond formation by cross-coupling of alkenyl halides with the phosphite is one of the attractive and important pathways for the first asymmetric synthesis of NZ-105. We accomplished this type of the Arbuzov reaction with trialkyl phosphite, though the Arbuzov reactions in previous studies were carried out with dialkyl phosphite.

* To whom correspondence should be addressed.

Chart 1

The prochiral compound (**1**) was hydrolyzed by lipase AH (*Pseudomonas sp.*), according to our previous report,¹¹⁾ in diisopropyl ether saturated with water at room temperature (Chart 1) to give the optically active (*S*)-monocarboxylic acid (**2a**) in 95 % yield. (>> 99 % e.e. : Optical yield was determined by HPLC analysis after conversion into methyl ester.) Bromination of the monocarboxylic acid (**2a**) was carried out with pyridinium bromide perbromide in dichloromethane¹²⁾ at -20 °C to give the bromide (**3a**) in 63 % yield. Phosphination of the bromide (**3a**), as a key step, was carried out with the tri-alkyl phosphite (**4**) in the presence of tetrakis-triphenylphosphine palladium and triethylamine in toluene for 5 hours at 110 °C to give the product (**5a**) in 87 % yield. It was found that this reaction proceeded in good yield without racemization. Deprotection of the methoxymethyl group of **5a** with hydrochloric acid and the pivaloyloxymethyl group with sodium hydroxide gave the monocarboxylic acid (**6a**) in 86 % yield (>> 99 % e.e.) from **5a**. And the (*R*)- NZ-105 was obtained by esterification with the 2-

[benzyl(phenyl)amino]-ethanol in the presence of WSC according to our reported method. For the synthesis of the (*S*)-NZ-105, the prochiral compound (**1**) was hydrolyzed by lipase PS (*Pseudomonas* sp.) similarly to give the (*R*)-monocarboxylic acid (**2b**) in 44 % yield (82 % e.e.), which was then converted to (*S*)-NZ-105 by the same process as the synthesis of *R* isomer.

The commercially available monocarboxylic acid¹³⁾ (**2c**: (*S*)-monopivaloyloxymethyl ester: (*S*)-MPE) was also converted to the (*R*)-(-)-NZ-105 (Chart 1). Consequently, it was found that phosphination of the bromide (**3c**) with the trialkyl phosphite (**4**) catalyzed by tetrakis-triphenyl phosphine palladium proceeded to give the desired product (**5c**) in 52 % yield and the recovered bromide (**3c**) in 17 % yield without racemization. And hydrolysis of the (*R*)-**5c** gave the (*R*)-**6a** in 43 % yield.

By using the modified Hunsdiecker reaction with pyridinium ion and the Michaelis-Arbuzov rearrangement catalyzed by zerovalent palladium, the first asymmetric synthesis of both (*S*)-(+)- and (*R*)-(-)-NZ-105 was established.

REFERENCES

- 1) Bossert F., Vater W., *Res. Rev.*, **1989**, 9, 291.
- 2) Sakoda R., Matsumoto H., Seto K., *Chem. Pharm. Bull.*, **1992**, 40, 2377.
- 3) a) Sakoda R., Kamikawaji Y., Seto K., *Chem. Pharm. Bull.*, **1992**, 40, 2362. b) Idem, *ibid.*, **1992**, 40, 2370.
- 4) Yamashita T., Masuda Y., Sakai T., Tanaka S., Kasuya Y., *Jpn. J. Pharmacol.*, **1991**, 57, 337.
- 5) Ebiike H., Yamazaki Y., Achiwa K., *Chem. Pharm. Bull.*, **1995**, 43, 1251.
- 6) a) Michaelis A., Kaehene R., *Chem. Ber.*, **1898**, 31, 1408. b) Arbuzov A. E., *J. Russ. Phys. Chem. Soc.*, **1906**, 38, 687.
- 7) a) Arbuzov B. A., *Pure Appl. Chem.*, **1964**, 9, 307. b) Henning H. G., Hilgetag G., *Z. Chem.*, **1967**, 7, 169. c) Arbuzov B. A., *Z. Chem.*, **1974**, 14, 41. d) Bhattacharya A. K., Thyagarajan G., *Chem. Rev.*, **1981**, 415.
- 8) a) Kozikowski A. D., Wetter H. F., *Synthesis*, **1976**, 561. b) Trost B. M., *Tetrahedron*, **1977**, 33, 2615.
- 9) a) Yamamura N., Moritani I., Murahashi S.-I., *J. Organomet. Chem.*, **1975**, 91, C39. b) Dang H. P., Linstumelle G., *Tetrahedron Lett.*, **1978**, 191. c) Mallan J. M., Bebb R. L., *Chem. Rev.*, **1969**, 69, 693. d) Baba S., Negishi E., *J. Am. Chem. Soc.*, **1976**, 98, 6729. e) Negishi E., Van Horn D. E., *J. Am. Chem. Soc.*, **1977**, 99, 3168. f) Dieck H. A., Heck F. R., *J. Organomet. Chem.*, **1975**, 93, 259. g) Cassar L., *ibid.*, **1975**, 93, 253. h) Sonogashira K., Tohda Y., Hagihara N., *Tetrahedron Lett.*, **1975**, 4469. i) King A. O., Okukado N., Negishi E., *J. Chem. Soc., Chem. Commun.*, **1977**, 683. j) Yamamura K., Murahashi S.-I., *Tetrahedron Lett.*, **1977**, 4429. k) Murahashi S.-I., Yamamura M., Yanagisawa K.-I., Mita N., Kondo K., *J. Org. Chem.*, **1979**, 44, 2408.
- 10) a) Hirao T., Masunaga T., Yamada N., Ohshiro Y., Agawa T., *Bull. Chem. Soc. Jpn.*, **1982**, 55, 909. b) Blumb J., Obrycki R., Griffin C. E., *J. Org. Chem.*, **1966**, 31, 2455. c) Obruchki R., Griffin C. E., *ibid.*, **1968**, 33, 632.
- 11) a) Ebiike H., Terao Y., Achiwa K., *Tetrahedron Lett.*, **1991**, 32, 5805. b) Hirose Y., Kariya K., Sasaki I., Kurono Y., Ebiike H., Achiwa K., *ibid.*, **1992**, 33, 7157. c) Hirose Y., Kariya K., Nakanishi Y., Kurono Y., Achiwa K., *ibid.*, **1995**, 36, 1063.
- 12) a) Johnson R. G., *Chem. Rev.*, **1956**, 56, 219. b) Hunsdiecker H., Hunsdiecker C., *Ber.*, **1942**, 75, 291.
- 13) This compound **2c** was established the synthetic method in our laboratory and is produced by Amano Pharmaceutical Co., Ltd. using this method.; See reference 11).

(Received February 15, 1996; accepted March 21, 1996)