Astertarone A: A Triterpenoid Ketone Isolated from the Roots of *Aster tataricus* L.

Toshihiro AKIHISA,^{*,a} Yumiko KIMURA,^b Kazuo KOIKE,^c Takaaki TAI,^d Ken YASUKAWA,^b Koichi ARAI,^e Yasuhiro SUZUKI,^a and Tamotsu NIKAI^d

College of Science and Technology, Nihon University,^a 1–8 Kanda Surugadai, Chiyoda-ku, Tokyo 101–8308, Japan, College of Pharmacy, Nihon University,^b 7–7–1 Narashinodai, Funabashi-shi, Chiba 274–8555, Japan, School of Pharmaceutical Sciences, Toho University,^c 2–2–1 Miyama, Funabashi-shi, Chiba 274–8510, Japan, Central Research Laboratory, Kotaro Pharmaceutical Co. Ltd.,^d 47–3 Suga-cho, Takatsuki-shi, Osaka 569–0022, Japan, and Meikai University School of Dentistry,^e 1–1 Keyakidai, Sakado-shi, Saitama 350–0283, Japan.

Received July 13, 1998; accepted August 22, 1998

The structure of astertarone A isolated from the root extract of *Aster tataricus* L. (Compositae) was established to be D:A-friedoeuph-21-en-3-one (1) based on spectroscopic methods. This is the first example of a naturally occurring triterpenoid with a D:A-friedoeuphane skeleton. Isolation and identification of epishionol (2), shionone (3), friedelin, epifriedelinol, and ψ -taraxasterol are also described.

Key words *Aster tataricus*; Compositae; roots; triterpenoid ketone; astertarone A

The roots of *Aster tataricus* L. (Compositae) are used as the Chinese crude drug Asteris Radix (Japanese name: Shion) and are combined in various traditional Chinese prescriptions for use as cough medicines, expectorants and diuretics.¹⁾ Constituents of the root extract have been investigated and the presence of a triterpenoid ketone, shionone (shion-21-en-3-one; 3),^{2–4)} seven triterpenoid glycosides, aster saponins A–G,^{5–7)} and three monoterpenoid glycosides, shionosides A–C,^{7,8)} have been reported. In this paper, we report the isolation and structure elucidation of a new triterpenoid ketone designated astertarone A (1) from the methanol extract of *A. tataricus* roots. Isolation and characterization of epishionol (shion-21-en-3 β -ol; 2), a known synthetic triterpene alcohol,^{2–4,9–11)} which has not been isolated before as a natural product, is also described.

Column chromatography of the hexane soluble portion of the methanol extract of *A. tataricus* roots on silica gel afforded the triterpenoid ketone and triterpene alcohol fractions. Crystallization of the ketone fraction from acetone–methanol followed by preparative HPLC of the filtrate portion eventually yielded compound 1 in addition to 3^{2–4)} and friedelin (D: A-friedoeuph-21-en-3-one).¹²⁾

Compound 1, molecular formula C₃₀H₅₀O determined from its high-resolution mass spectrum (HR-MS) ([M]⁺, *m/z* 426.3848), gave IR absorptions at 1712 (ketone) and 824 cm^{–1} (trisubstituted double bond). Compound 1 displayed four methyl singlets at δ 0.72, 0.79, 0.86, and 0.87, two methyl doublets at δ 0.86 (*J*=6.5 Hz) and 0.88 (*J*=6.6 Hz), two olefinic methyl singlets at δ 1.61 and 1.69, with an olefinic methine signal at δ 5.09 (tt, *J*=1.4, 6.9 Hz) in the ¹H-NMR spectrum. These data, in combination with MS fragment ions having *m/z* 313 (base peak) [loss of side-chain (C₈H₁₅) with 2H transfer], 273 [loss of side-chain and ring D (C₃H₆)], and 69 [CH₂CH=C(Me)₂]⁺ (C-20–C-22, and C-29 and C-30),¹³⁾ suggested that compound 1 has a C₈ side-chain containing an isopropylidene group, like euphol (eupha-8,24-dien-3 β -ol) and tirucallol (tirucalla-8,24-dien-3 β -ol),¹⁴⁾ and a tetracyclic triterpenoid skeleton with a keto group most probably located at C-3. Moreover, the presence as a secondary methyl group of one of the five skeletal methyl groups sug-

gested it to possess a D:A-friedoeuphane or tirucallane skeleton. From the foregoing, compound 1 was assigned a D:A-friedoeuph/tirucall-21-en-3-one structure with as-yet-to-be determined stereochemistry. Analysis of the ¹³C distortionless enhancement by polarization transfer (DEPT), ¹H–¹H correlation spectroscopy (COSY), ¹H detected multiple quantum coherence (HMQC), and heteronuclear multiple-bond correlation (HMBC) spectra, and the ¹³C- and ¹H-NMR spectral comparison of 1 (Table 1) with 3 (see the Experimental section), friedelin,¹²⁾ and euphol and tirucallol acetates¹⁴⁾ confirmed the above assumption.

The stereochemistry of compound 1 was established by phase-sensitive nuclear Overhauser and exchange spectroscopy (NOESY). Compound 1 showed significant NOE correlations between [H-4 α –H-10 α] and [H-23 (4 β -Me)–H-24 (5 β -Me)–H-25 (9 β -Me)–H-7 β –H-26 (14 β -Me)] (Fig. 1), which were observed also for compound 3 (see the Experimental section), demonstrating that 1 possessed the same stereochemistry as 3 as far as rings A, B and C and the C/D ring junction (13 α , 14 β) were concerned. In addition, compound 1 exhibited NOE correlations between [H-26 (14 β -Me)–H-17–H-28], [H-28–H-16 α , β], and [H-16 α –H-18–H-27 (13 α -Me)] which were consistent with those observed for euphol acetate.¹³⁾ From these data, it was concluded that 1 was D:A-friedoeuph-21-en-3-one with a 18*R*-chirality which

Chart 1. Structures of Triterpenoids from *Aster tataricus* Roots

* To whom correspondence should be addressed.

Table 1. ^{13}C -NMR and ^1H -NMR Spectral Data (δ Values; CDCl_3) and ^1H - ^{13}C Long-Range Correlations of Astertarone A (**1**) by ^1H - ^{13}C COSY, HMBC and HMQC

C No.	δ_{C}		δ_{H}^a	Cross peaks (δ_{C}) in HMBC spectrum
1	22.8	CH_2	1.96 (α), 1.74 (β)	
2	41.6	CH_2	2.30 (α), 2.39 (β)	
3	213.2	C		
4	58.3	CH	2.28	6.8 (C-23), 14.7 (C-24), 42.4 (C-5), 213.2 (C-3)
5	42.4	C		
6	40.8	CH_2	1.32 (α), 1.72 (β)	
7	20.3	CH_2	1.29 (α), 1.57 (β)	
8	50.0	CH	1.58	16.0 (C-26), 18.5 (C-25), 20.3 (C-7), 37.9 (C-9), 59.1 (C-10)
9	37.9	C		
10	59.1	CH	1.60	18.5 (C-25), 22.8 (C-1), 36.8 (C-11), 41.6 (C-2), 42.4 (C-5)
11	36.8	CH_2	1.43 (2H)	
12	34.0	CH_2	1.22 (α), 1.14 (β)	
13	46.2	C		
14	48.2	C		
15	30.1	CH_2	1.76 (α), 1.56 (β)	
16	28.1	CH_2	1.30 (α), 1.89 (β)	
17	49.8	CH	1.46	19.3 (C-27), 35.3 (C-18), 46.2 (C-13)
18	35.3	CH	1.52	19.0 (C-28), 35.4 (C-19)
19	35.4	CH_2	1.11, 1.61	
20	24.7	CH_2	1.87, 2.04	
21	125.2	CH	5.09 (tt, 1.4, 6.9)	
22	131.0	C		
23	6.8	Me	0.88 (d, 6.6)	14.7 (C-24), 58.3 (C-4), 213.2 (C-3)
24	14.7	Me	0.72 (s)	40.8 (C-6), 42.4 (C-5), 58.3 (C-4), 59.1 (C-10)
25	18.5	Me	0.86 (s)	36.8 (C-11), 37.9 (C-9), 50.0 (C-8), 59.1 (C-10)
26	16.0	Me	0.87 (s)	30.1 (C-15), 42.4 (C-5), 46.2 (C-13), 48.2 (C-14)
27	19.3	Me	0.79 (s)	34.0 (C-12), 46.2 (C-13), 48.2 (C-14), 49.8 (C-17)
28	19.0	Me	0.86 (d, 6.5)	35.3 (C-18), 35.4 (C-19), 49.8 (C-17), 50.0 (C-8)
29	25.8	Me	1.69 (s)	17.7 (C-30), 125.2 (C-21), 131.0 (C-22)
30	17.7	Me	1.61 (s)	25.8 (C-29), 125.2 (C-21), 131.0 (C-22)

a) Figures in parentheses in the ^1H chemical shift column denote J values (Hz).

1

Fig. 1. Energy-Minimized Conformations and Some Representative NOE Correlations ($\leftrightarrow$) for Astertarone A (**1**)

we named astertarone A.

The most stable conformation of **1** was simulated using MacroModel. The results of the calculations^{15,16} are shown in Fig. 1 together with the significant NOE's ($\leftrightarrow$). The conformer of simulated **1** shows C-19 of the side chain at C-17 oriented into a "left-handed" conformation (C-19 *trans*-oriented to C-13) similar to that of butyrospermol (eupha-7,24-dien-3 β -ol) acetate¹⁷ and the crystal structure of euphol acetate.¹⁸ This was consistent with the NOE experiment done in solution, thus confirming the proposed structure of **1**.

Preparative HPLC of the triterpene alcohol fraction yielded epishionol (**2**),^{2-4, 9-11} along with epifriedelinol

(D:A-friedoolean-3 β -ol)^{2,10,11} and ψ -taraxasterol (taraxast-20-en-3 β -ol).¹⁹

Astertarone A (**1**) is the first naturally occurring triterpenoid possessing a D:A-friedoeuphane skeleton. A triterpenoid hydrocarbon possessing a closely related structure with **1**, named aonena-3,21-diene with a (17 ξ , 18S)-D:A-friedotirucallane-type skeleton, has been isolated from the rhizomes of *Polypodiodes niponica*.²⁰ Although epishionol (**2**) has previously been synthesized from shionone (**3**) by reduction,^{2-4, 9-11} this is the first report for its detection as a natural product.

Experimental

Crystallizations were performed from acetone–MeOH. Melting points measured were uncorrected. Reverse-phase HPLC was carried out on an octadecyl silica column (Supiorex ODS S-5 μ m column, 10 mm i.d. $\times$ 25 cm; Shiseido Co., Ltd., Tokyo) at 25 °C with MeOH (4 ml/min) as mobile phase. GLC was performed using a DB-17 fused-silica capillary column (30 m $\times$ 0.3 mm i.d., column temp., 275 °C). In both HPLC and GLC, cholesterol (cholest-5-en-3 β -ol) was the standard for the determination of R_f of triterpenoids. Electron-impact MS and HR-MS were recorded at 70 eV. NMR spectra were recorded at 400 MHz (1 H-NMR) and 100.6 MHz (13 C-NMR) in $CDCl_3$ with tetramethylsilane (TMS) (1 H-NMR) and $CDCl_3$ at δ 77.0 (13 C-NMR) as internal standard, and chemical shifts were recorded in δ values. IR spectra were recorded in KBr. Instrumental details were the same as described previously.²¹⁾ The dried roots of *A. tataricus* were purchased from Tochimoto Co., Ltd. (Osaka). Identification of the plant material was performed by Prof. Norio Sahashi (Department of Biology, School of Pharmaceutical Sciences, Toho University). Friedelin¹²⁾ and ψ -taraxasterol¹⁹⁾ were used as reference compounds.

Isolation Procedures Dried and ground *A. tataricus* roots (2.5 kg) were extracted 3 $\times$ for 3 d with MeOH (6 l each) at room temperature to give an extract (600 g) which was partitioned between EtOAc–H₂O to afford an EtOAc fraction (34.5 g). This extract was partitioned between hexane–MeOH–H₂O (19 : 19 : 2). The hexane fraction (19.9 g) was subjected to column chromatography (CC) on silica gel (600 g) using hexane and a hexane–EtOAc gradient of 50 : 1–1 : 1, which yielded triterpenoid ketone (A) and triterpene alcohol (B) fractions. Repeated CC of the fractions using the same solvent system eventually yielded purified fractions A (1.8 g) and B (0.5 g). Recrystallization of the fraction A from acetone–MeOH gave crystallized (1.2 g), constituted with shionone (3), and filtrate portions (360 mg). Preparative HPLC of the filtrate portion yielded astertarone A [1; 5 mg; amorphous solid; R_f : 1.09 (HPLC), 2.28 (GLC)], 3 [227 mg; mp 162–164 °C (colorless needles) (lit.,²⁾ mp 158.5–159.5 °C; lit.,⁸⁾ mp 155–156 °C]; R_f : 1.19 (HPLC), 2.54 (GLC)], and friedelin [32 mg; mp 266–269 °C (colorless plates) (lit.,¹²⁾ mp 268–269 °C]; R_f : 1.48 (HPLC), 3.05 (GLC)]. On the other hand, preparative HPLC of the fraction B afforded epishionol [2; 10 mg; mp 116–119 °C (colorless needles) (lit.,²⁾ mp 117.5–118.5 °C]; R_f : 1.15 (HPLC), 2.32 (GLC)], epifriedelinol [35 mg; mp 275–279 °C (colorless plates) (lit.,²⁾ mp 279–283 °C; lit.,⁸⁾ mp 299 °C]; R_f : 1.54 (HPLC), 2.76 (GLC)], and ψ -taraxasterol [67 mg; mp 290–292 °C (colorless plates); R_f : 1.23 (HPLC), 2.72 (GLC)]. Identification of 2,^{10,11)} 3,^{10,11)} and epifriedelinol^{10,11)} was based on spectroscopic (MS and 1 H-NMR) comparison with data in the literature, whereas that of friedelin and ψ -taraxasterol was done by chromatographic (HPLC, GLC) and spectroscopic (MS, 1 H-NMR) comparison with reference compounds. Since the fully assigned 1 H- and 13 C-NMR spectral data for 2 and 3 were unavailable in the literature, these also are shown below accompanied by some representative NOE correlations. The NMR assignments were aided by 13 C DEPT, 1 H– 1 H COSY, HMQC, HMBC, and phase-sensitive NOESY spectroscopy.

Astertarone A (1): IR ν_{max} cm^{-1} : 1712, 824. MS m/z (%): 426 (M^+ , 13), 411 (21), 341 (19), 313 (100), 273 (3), 245 (3), 218 (14), 203 (7), 191 (10), 189 (6), 69 (97). HR-MS: m/z 426.3848 [Calcd for $C_{30}H_{50}O$ (M^+): 426.3859]; 313.2506 [Calcd for $C_{22}H_{33}O$: 313.2528]; 273.2243 [Calcd for $C_{19}H_{29}O$: 273.2217]; 69.0697 [Calcd for C_5H_9 : 69.0703]. See Table 1 for the 13 C- and 1 H-NMR data.

Epishionol (2): 13 C- and 1 H-NMR: C-1 [δ_C 15.8; δ_H 1.46 (α), 1.56 (β)], C-2 [35.2; 1.57 (α), 1.90 (β)], C-3 [72.8; 3.73, dt, $J=2.5, 3.1$ Hz], C-4 [49.2; 1.25], C-5 [37.9], C-6 [41.7; 0.95 (α), 1.72 (β)], C-7 [17.3; 1.26 (α), 1.37 (β)], C-8 [50.0; 1.22], C-9 [38.2], C-10 [61.1; 0.96], C-11 [35.3; 1.39 (2H)], C-12 [32.5; 0.89 (α), 1.53 (β)], C-13 [36.9], C-14 [38.7], C-15 [29.2; 1.30 (α), 1.24 (β)], C-16 [34.8; 1.33 (α), 1.60 (β)], C-17 [31.8], C-18 [44.7; 1.13 (α), 1.19 (β)], C-19 [43.6; 1.16, 1.67], C-20 [23.6; 1.85, 1.99], C-21 [125.4; 5.10, tt, $J=1.5, 7.3$ Hz], C-22 [130.7], C-23 [11.6; 0.93, d, $J=7.0$ Hz], C-24 [16.4; 0.95, s], C-25 [20.0; 0.92, s], C-26 [15.1; 0.87, s], C-27 [20.7; 1.09, s], C-28 [33.0; 0.89, s], C-29 [25.7; 1.68, s], C-30 [17.6; 1.60, s]. Significant NOE correlations observed were between [H-3 α –H-4 α –H-10 α –H-8 α –H-27 (13 α -Me)], [H-23 (4 β -Me)–H-6 β –H-24 (5 β -Me)], [H-6 β –H-7 β –H-25

(9 β -Me)], and [H-7 β –H-26 (14 β -Me)–H-18 β –H-28 (17 β -Me)].

Shionone (3): 13 C- and 1 H-NMR: C-1 [δ_C 22.3; δ_H 1.97 (α), 1.70 (β)], C-2 [41.5; 2.40 (α), 2.30 (β)], C-3 [213.1], C-4 [58.2; 2.25, q, $J=6.2$ Hz], C-5 [42.2], C-6 [41.2; 1.24 (α), 1.73 (β)], C-7 [17.9; 1.47 (α), 1.35 (β)], C-8 [50.0; 1.32], C-9 [38.5], C-10 [59.7; 1.58], C-11 [35.3; 1.40, 1.52], C-12 [32.3; 0.92 (α), 1.58 (β)], C-13 [36.9], C-14 [38.6], C-15 [29.3; 1.34 (α), 1.27 (β)], C-16 [34.7; 1.36 (α), 1.59 (β)], C-17 [31.7], C-18 [44.6; 1.12 (α), 1.22 (β)], C-19 [43.6; 1.17, 1.67], C-20 [23.2; 1.83, 2.00], C-21 [125.3; 5.10, tt, $J=1.3, 1.5, 7.2$ Hz], C-22 [130.8], C-23 [6.8; 0.88, d, $J=6.6$ Hz], C-24 [14.6; 0.72, s], C-25 [19.6; 0.93, s], C-26 [15.2; 0.90, s], C-27 [20.6; 1.14, s], C-28 [33.0; 0.91, s], C-29 [25.7; 1.69, s], C-30 [17.6; 1.61, s]. The following diagnostic NOE correlations were observed between [H-4 α –H-10 α –H-8 α , H-11 α –H-27 (13 α -Me)] and [H-23 (4 β -Me)–H-24 (5 β -Me)–H-25 (9 β -Me)–H-7 β –H-26 (14 β -Me)–H-18 β –H-28 (17 β -Me)].

Acknowledgements The authors are indebted to Prof. Norio Sahashi (Toho University) for the identification of the plant material.

References and Notes

- 1) Namba T., "The Encyclopedia of Wakan-Yaku (Traditional Sino-Japanese Medicines) with Color Pictures," Revised edn., Vol. I, Hoikusha, Osaka, 1994, p. 161.
- 2) Takahashi M., Kamisako W., Ishimasa S., Miyamura K., *Yakugaku Zasshi*, **79**, 1281–1283 (1959).
- 3) Tanahashi Y., Takahashi T., Patil F., Ourisson G., *Bull. Soc. Chim. Fr.*, **1964**, 584–590.
- 4) Tanahashi Y., Moriyama Y., Takahashi T., *Bull. Soc. Chim. Fr.*, **1966**, 1670–1677.
- 5) Nagao T., Hachiyama S., Okabe H., Yamauchi T., *Chem. Pharm. Bull.*, **37**, 1977–1983 (1989).
- 6) Nagao T., Okabe H., Yamauchi T., *Chem. Pharm. Bull.*, **38**, 783–785 (1990).
- 7) Cheng D., Shao Y., *Phytochemistry*, **35**, 173–176 (1994).
- 8) Nagao T., Okabe H., Yamauchi T., *Chem. Pharm. Bull.*, **36**, 571–577 (1988).
- 9) Tanahashi Y., Moriyama Y., Takahashi T., Patil F., Ourisson G., *Bull. Soc. Chim. Fr.*, **1966**, 2374–2377.
- 10) Hirota H., Moriyama Y., Tsuyuki T., Tanahashi Y., Takahashi T., Katoh Y., Satoh H., *Bull. Chem. Soc. Jpn.*, **48**, 1884–1888 (1975).
- 11) Kikuchi T., Yokoi T., Niwa M., Shingu T., *Chem. Pharm. Bull.*, **28**, 2014–2023 (1980).
- 12) Akihisa T., Yamamoto K., Tamura T., Kimura Y., Iida T., Nambara T., Chang F. C., *Chem. Pharm. Bull.*, **40**, 789–791 (1992).
- 13) Goad L. J., Akihisa T., "Analysis of Sterols," Blackie Academic and Professional, London, 1997.
- 14) Akihisa T., Kimura Y., Koike K., Shibata T., Yoshida Z., Nikaido T., Tamura T., *J. Nat. Prod.*, **61**, 409–412 (1998).
- 15) Calculations were performed using MacroModel Ver. 6.0 with extended MM3 parameters. Conformation with minimum steric energy was obtained through a Metropolis Monte Carlo procedure. Final structures were depicted using Chem3D program (Cambridge Scientific Computing Inc., Cambridge, MA).
- 16) Mohamadi F., Richards N. G. L., Guida W. C., Liskamp R., Lipton M., Caufield C., Chang G., Hendrickson T., Still W. C., *J. Comput. Chem.*, **11**, 440–467 (1990).
- 17) Akihisa T., Kimura Y., Kokke W. C. M. C., Takase S., Yasukawa K., Tamura T., *J. Chem. Soc., Perkin Trans. 1*, **1996**, 2379–2384.
- 18) Nes W. D., Wong R. Y., Benson J. R., Landrey J. R., Nes W. R., *Proc. Natl. Acad. Sci. U.S.A.*, **81**, 5896–5900 (1984).
- 19) Akihisa T., Yasukawa K., Oinuma H., Kasahara Y., Yamanouchi S., Takido M., Kumaki K., Tamura T., *Phytochemistry*, **43**, 1255–1260 (1996).
- 20) Arai Y., Hirohata M., Ageta H., *Tetrahedron Lett.*, **30**, 7209–7212 (1989).
- 21) Akihisa T., Kimura Y., Kokke W. C. M. C., Itoh T., Tamura T., *Chem. Pharm. Bull.*, **44**, 1202–1207 (1996).