New Syntheses of Murrayaquinone A and Furostifoline

Hitomi HAGIWARA, Tominari CHOSHI, Hiroyuki FUJIMOTO, Eiichi SUGINO, and Satoshi HIBINO*

Faculty of Pharmacy and Pharmaceutical Sciences, Fukuyama University, Fukuyama, Hiroshima 729-0292, Japan.

Received September 29, 1998; accepted November 13, 1998

Starting from 2-chloro-3-formylindole, new syntheses of murrayaquinone A and furostifoline were achieved by an allene-mediated electrocyclic reaction involving the indole 2,3-bond.

Key words murrayaquinone A; carbazole-1,4-quinone; furostifoline; furo[3,2-*a*]carbazole; synthesis; electrocyclic reaction

The carbazole-1,4-quinone alkaloid, murrayaquinone A (**1**), was isolated from the root bark of *Murraya eucrestifolia* HAYATA (Rutaceae) together with three related alkaloids (murrayaquinone B—D) by Furukawa and co-workers from 1983 to 1985.¹⁾ The furo[3,2-*a*]carbazole alkaloid, furostifoline (**3**), was also isolated in 1990 from the same origin by the same group (Chart 1).²⁾ Extracts of the leaves and bark of this tree have been used as a folk medicine in Taiwan. Among these alkaloids, murrayaquinone A (**1**) has cardiotonic activity on guinea-pig papillary muscle.³⁾ Since they were isolated, synthetic studies of these alkaloids have been reported by many groups.^{4,5)}

We were also interested in the syntheses of the carbazole-1,4-quinone alkaloid, murrayaquinone A (**1**), and the novel furo[3,2-*a*]carbazole alkaloid, furostifoline (**3**), in the course of our synthetic studies. Recently, we reported the total syntheses of the 3-oxygenated and 3,4-dioxygenated carbazole alkaloids, carazostatins and carbazoquinocins (B—F), based on the thermal electrocyclic reaction of the 3-alkenyl-2-alkenylindole intermediate derived from 3-alkenyl-2-propargylindole.⁶⁾

In this paper, we describe the synthesis of 4-oxygenated carbazole (**2**) which consists of a formal total synthesis of murrayaquinone A (**1**) and a new total synthesis of the furo[3,2-*a*]carbazole, furostifoline (**3**), based on the thermal electrocyclic reaction of a new type of hexatriene system involving the 2-alkenyl-3-allenylindole intermediate derived from 2-alkenyl-3-propargylindole.

We chose 2-chloro-3-formylindole (**4**) as the common starting material for the syntheses of the two target molecules. The 2-chloroindole (**4**) was treated with benzyl-oxymethyl chloride (BOMCl) and K₂CO₃ to give the *N*-BOM-indole (**5**) (96%). The cross-coupling reaction of **5** with tributylvinyltin in the presence of PdCl₂(PPh₃)₂ and tetraethylammonium chloride in DMF at 80 °C was carried out to yield the 2-ethenylindole (**6**) (78%). The subsequent Grignard reaction of 3-formylindole (**6**) with ethynylmagnesium bromide followed by treatment with BOMCl afforded

the 2-ethenylindole (**6**) (78%). The subsequent Grignard reaction of 3-formylindole (**6**) with ethynylmagnesium bromide followed by treatment with BOMCl afforded

Chart 1

Chart 2

* To whom correspondence should be addressed.

Chart 3

the 2-ethenyl-3-propargylindole (**8**) (56%) as a new precursor of the hexatriene system. The thermal electrocyclic reaction of **8** was carried out in the presence of potassium *t*-butoxide at 90 °C⁶) to produce 3-methyl-4-oxygenated carbazole (**9**) (81%), which was deprotected with Birch conditions to give a separable mixture of the *N*-hydroxymethyl-4-hydroxy-3-methylcarbazole (**10**) (75%) and the 4-hydroxy-3-methylcarbazole (**2**) (22%). Treatment of **10** with Triton B⁷) gave **2** [71%; mp 156–158 °C (Lit.,^{3g}) mp 161–163 °C] (Chart 2). This compound was identical in all respects to the precursor (**2**) of murrayaquinone A (**1**), as reported previously.^{3g}) This constitutes the formal total synthesis of murrayaquinone A (**1**).

The cross-coupling reaction of 2-chloroindole (**4**) with furan-3-boronic acid⁸) was carried out in the presence of Pd(OAc)₂, PPh₃, and triethylamine in DMF at 100 °C to obtain 2-(3-furyl)indole (**11**) (84%). After protection of the indole nitrogen atom with BOMCl, the subsequent Grignard reaction of **12** with ethynylmagnesium bromide gave the alcohol (**13**) (97%). The alcohol (**13**) was treated with BOMCl and NaH to yield the 2-(3-furyl)-3-propargylindole (**14**), which was subjected to the thermal electrocyclic reaction in the presence of potassium *t*-butoxide at 90 °C⁶) to produce the 4-oxygenated tetracyclic furocarbazole (**15**) (65% yields from **13**). Birch reduction of **15** for the deprotection of *N,O*-bisBOM groups also gave a separable mixture of the *N*-(hydroxymethyl)furocarbazole (**16**) (41%) and the furocarbazole (**17**) (51%). Treatment of **16** with Triton B⁷) afforded **17** (93%), which was converted into the triflate (**18**) (92%) with trifluoromethanesulfonic anhydride and pyridine. Finally, reductive elimination⁹) of the 4-trifluoromethanesulfonyloxy group of **18** with HCOOH, triethylamine, Pd(OAc)₂, and PPh₃ in DMF at 60 °C afforded furostifoline (**3**) (98%) (Chart 3).¹⁰) The synthetic furostifoline (**3**) was identical in

all respects to natural²⁾ and synthetic⁵⁾ furostifoline.

Thus the new synthetic route to the carbazole-1,4-quinone, murrayaquinone A (**1**), has been established by a new type of allene-mediated electrocyclic reaction. Based on this methodology, the total synthesis of furostifoline (**3**) has also been completed.

References

- For the isolation of murrayaquinone A: a) Wu T.-S., Ohta T., Furukawa H., *Heterocycles*, **20**, 1267 (1983); b) Furukawa H., Wu T.-S., Ohta T., Kouh C.-S., *Chem. Pharm. Bull.*, **33**, 4132 (1985).
- For the isolation of furostifoline: Ito C., Furukawa H., *Chem. Pharm. Bull.*, **38**, 1548 (1990).
- Takeya K., Itoigawa, M. Furukawa H., *Eur. J. Pharmacol.*, **169**, 137 (1989).
- For the synthesis of murrayaquinone A: a) Ramesh K., Kapil R. S., *Chem. Ind.*, **1986**, 186; b) Ramesh K., Kapil R. S., *J. Nat. Prod.*, **50**, 932 (1987); c) Martin T., Moody C. J., *J. Chem. Soc., Perkin Trans. 1*, **1988**, 235; d) Yogo M., Ito C., Furukawa H., *Chem. Pharm. Bull.*, **39**, 328 (1991); e) Miki Y., Hachiken H., *Synlett*, **1993**, 333; f) Wada A., Hirai S., Hanaoka M., *Chem. Pharm. Bull.*, **42**, 416 (1994); g) Matsuo K., Ishida S., *ibid.*, **42**, 1325 (1994).
- For the synthesis of furostifoline: a) Knölker H.-J., Fröhner W., *Tetrahedron Lett.*, **37**, 9183 (1996); b) Beccalli E. M., Clerici F., Marchesini A., *Tetrahedron*, **54**, 11675 (1998).
- a) Choshi T., Sada T., Fujimoto H., Nagayama C., Sugino E., Hibino S., *Tetrahedron Lett.*, **37**, 2593 (1996); b) Choshi T., Sada T., Fujimoto H., Nagayama C., Sugino E., Hibino S., *J. Org. Chem.*, **62**, 2535 (1997).
- Anderson H. J., Groves J. K., *Tetrahedron Lett.*, **1971**, 3165.
- Roques B. P., Florentin D., Callanquin M., *J. Heterocycl. Chem.*, **12**, 195 (1975).
- Cacchi S., Ciattini P. G., Morera E., Ortar G., *Tetrahedron Lett.*, **27**, 5541 (1986).
- Furostifoline (**3**): mp 171–173 °C (Lit., mp 174–175 °C^{5a}) and 175 °C^{5b}); ¹H-NMR (300 MHz, CDCl₃) δ 2.68 (3H, d, *J*=1.1 Hz), 7.00 (1H, d, *J*=2.2 Hz), 7.26 (1H, dt, *J*=1.1, 8.1 Hz), 7.38 (1H, dt, *J*=1.1, 8.1 Hz), 7.49 (1H, br d, *J*=8.1 Hz), 7.73 (1H, d, *J*=2.2 Hz), 7.78 (1H, br s), 8.06 (1H, br d, *J*=8.1 Hz), 8.26 (1H, br s). Ms *m/z*: 221 (M⁺).