Studies on Disease-Modifying Antirheumatic Drugs. III.¹⁾ Bone Resorption Inhibitory Effects of Ethyl 4-(3,4-Dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (TAK-603) and Related Compounds

Atsuo BABA,^{*,a} Tsuneo ODA,^a Shigehisa TAKETOMI,^a Kohei NOTOYA,^a Atsushi NISHIMURA,^b Haruhiko MAKINO,^a and Takashi SOHDA^a

Pharmaceutical Research Division^a and Discovery Research Division,^b Takeda Chemical Industries, Ltd., 17–85, Jusohonmachi 2-chome, Yodogawa-ku, Osaka 532–8686, Japan. Received October 7, 1998; accepted December 1, 1998

In the course of our studies aimed at obtaining new drugs for treatment of bone and joint diseases, chemical modification of the potent bone resorption inhibitors justicidins, was performed and various naphthalene lactones, quinoline lactones and quinoline derivatives bearing an azole moiety at the side chain were prepared. Their inhibitory effects on bone resorption were evaluated by Raisz's method, and several compounds, including ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (6c, TAK-603), were found to have activities comparable with or superior to the justicidins. The 4-(3-isopropoxy-4-methoxy)-phenyl derivative (6d), in particular, displayed a marked increase in potency. TAK-603 and compound 6d were very effective in preventing osteoclast formation and bone resorption by mature osteoclasts. Further, TAK-603 was shown to be effective in preventing bone loss in ovariectomized mice.

Key word TAK-603; justicidin; bone and joint disease; antiresorptive activity; bone resorption

Rheumatoid arthritis (RA) is a serious, chronic, and systemic disease characterized by inflammation and progressive joint destruction. Since RA is an autoimmune disease, disease-modifying antirheumatic drugs (DMARDs), which have selective and direct effects on the abnormal immune system, have attracted a great deal of attention as potentially effective treatments for RA.²⁾ Development of osteopenia in the region of inflamed joints is a common clinical feature of RA. Previous histological studies of bone specimens obtained from patients with chronic RA have demonstrated an increase in bone resorption, suggesting increased bone turnover.³⁾ Therefore, DMARDs possessing antiresorptive activity as well as immune system modulating effects are expected to be useful in the treatment of RA.

Our search for a new type of DMARD with antiresorptive activity began with chemical modification of the potent bone resorption inhibitors justicidins (1, Fig. 1),⁴⁾ which were isolated from *Justicia Procumbens*, since factors such as interleukin (IL)-1 β and prostaglandin (PG) E₂ play important roles in both RA and bone metabolism.⁵⁾ Although justicidins possess only weak anti-inflammatory activity,⁶⁾ these compounds with antiresorptive activity are very attractive lead compounds for the development a new type of DMARD. Thus, the justicidin structure was modified to obtain compounds with better pharmacological properties. Naphthalene lactones (2), quinoline lactones (3), and quinoline derivatives (4–6), bearing a heteroaryl moiety at the side chain of the 2-position, were prepared to evaluate their antiresorptive and anti-inflammatory activities. We previously reported structure–activity relationships (SAR) with respect to the anti-inflammatory effect of these derivatives, leading to the finding of a new type of DMARD, TAK-603 (6c).⁷⁾ This report details some SAR information concerning the bone resorption inhibitory effects of this series of compounds.⁸⁾

Chemistry

The naphthalene lactone derivatives (2, Table 1) were pre-

pared using the method of Stevenson and co-workers,^{9,10)} as shown in Chart 1. Condensation of carboxylic acids (7) with alcohols (8) gave the esters (9), which were converted to the dihydronaphthalene lactones (10) by heating in acetic anhydride. Aromatization of 10 yielded 2.

Friedländer reaction¹¹⁾ of 2-aminobenzophenones (11) with tetronic acid afforded quinoline lactones (3) (Chart 2).¹²⁾

The syntheses of 2-(heteroarylthiomethyl)-(4), 2-(1-methylimidazol-2-ylethyl)-(5) and 2-(azolylmethyl)quinolines (6), the structures of which are shown in Table 2, were detailed in our previous reports.^{1,7)} Compounds 4c, g, i and m were newly prepared for this study by the same methods, and their physical and spectral data are shown in Table 3.

Results and Discussion

The structures and physical data for the naphthalene and quinoline derivatives prepared are shown in Tables 1–3. Since we selected the potent bone resorption inhibitors, the justicidins, as lead compounds aiming at a new type of DMARD, these compounds were first evaluated for their inhibition of ⁴⁵Ca release from a fetal long-bone culture system (Raisz's assay).¹³⁾ The results are summarized in Table 4.

Modification was begun by changing the alkoxy substituents on the naphthalene skeleton and the pendent phenyl moiety. In general, these justicidin analogues (2a–c) were fully active.

Replacement of the naphthalene ring by a quinoline ring

Fig. 1

* To whom correspondence should be addressed.

Table 1. Physical Data for Naphthalene Lactones **2** and Quinoline Lactones **3**

Compd.	R ¹	R ²	R ³	R ⁴	X	Yield (%) ^{a)}	Formula	mp (°C)	Recrystn. solvent
2a		OCH ₂ O		OCH ₂ O	CH	31	C ₂₀ H ₁₂ O ₆	203—205	AcOEt–acetone
2b		OCH ₂ O	OMe	OMe	CH	36	C ₂₁ H ₁₆ O ₆	230—231	AcOEt
2c		OCH ₂ O	H	OMe	CH	37	C ₂₀ H ₁₄ O ₅	209—210	AcOEt
3a		OCH ₂ O	OMe	OMe	N	84	C ₂₀ H ₁₅ NO ₆	252—253	CHCl ₃ –MeOH
3b	OMe	OMe	OMe	OMe	N	90	C ₂₁ H ₁₉ NO ₆	241—242	CHCl ₃ –MeOH
3c	OMe	OMe	H	H	N	65	C ₁₉ H ₁₅ NO ₄	210—211	EtOH

a) Yields of **2a–c** were based on **10**. Yields of **3a–c** were based on **11**.

Chart 1

Chart 2

was tried next in an attempt to improve the pharmacological properties, and it was found that activity was retained in the quinoline lactone derivatives (**3a–c** vs. **2a–c**). This finding suggested that the quinoline ring was an effective core structure, encouraging us to introduce various heteroaryl moieties to the 2-methyl position, generated by ring opening of the lactone. The 2-(1-methylimidazol-2-ylthiomethyl)quinoline derivative (**4a**) was the first compound found with the desired biological activity in this series of compounds. Thus, further modification of **4a** was performed.

Firstly, the effect of changing a heteroaryl moiety on the side chain at the 2-position of the quinoline skeleton was evaluated for compounds with a 1-methylimidazolyl (**4a**), a 4-methyltriazolyl (**4b**) or a thiazolyl (**4c**) moiety. These 2-azolythio derivatives exhibited potent activity comparable with that of **4a**. Compounds **4d–g** represent variations of the 6,7-dimethoxy moiety on the quinoline ring possessing the 2-(1-methylimidazol-2-ylthio)methyl substituent. Among these derivatives, compounds bearing a 6,7-dialkoxy moiety (**4a**, **4d**, **4e**) had potent activity. This result is similar to that for the lactone derivatives. Concerning the substituents on the pendent phenyl ring at the 4-position, potent activity was

Table 2. Structures of 2-(Heteroarylthiomethyl)quinoline Derivatives **4**, 2-(1-Methylimidazol-2-ylethyl)quinoline Derivative **5** and 2-(Azolylmethyl)quinoline Derivatives **6**

Compd.	R ¹	R ²	G	X	Y	Z	R ³
4a	6,7-(MeO) ₂	3,4-(MeO) ₂	S	N-Me	CH		
4b	6,7-(MeO) ₂	3,4-(MeO) ₂	S	N-Me	N		
4c	6,7-(MeO) ₂	3,4-(MeO) ₂	S	S	CH		
4d	6,7-(EtO) ₂	3,4-(MeO) ₂	S	N-Me	CH		
4e	6,7-O(CH ₂) ₂ O-	3,4-(MeO) ₂	S	N-Me	CH		
4f	H	3,4-(MeO) ₂	S	N-Me	CH		
4g	6-Cl	3,4-(MeO) ₂	S	N-Me	CH		
4h	6,7-(MeO) ₂	4-MeO	S	N-Me	CH		
4i	6,7-(MeO) ₂	4-EtO	S	N-Me	CH		
4j	6,7-(MeO) ₂	4-Me	S	N-Me	CH		
4k	6,7-(MeO) ₂	4-Cl	S	N-Me	CH		
4l	6,7-(MeO) ₂	H	S	N-Me	CH		
4m	6-Cl	2-Cl	S	N-Me	CH		
5	6,7-(MeO) ₂	3,4-(MeO) ₂	CH ₂	N-Me	CH		
6a	6,7-(MeO) ₂	3,4-(MeO) ₂					CH H
6b	6,7-(MeO) ₂	3,4-(MeO) ₂					CH Et
6c	6,7-(MeO) ₂	3,4-(MeO) ₂					N H
6d	6,7-(MeO) ₂	3-iso-PrO, 4-MeO					N H

observed for compounds with 4-methoxy (**4h**) and 4-methyl (**4j**) substituents, despite the reduction in activity with the unsubstituted compound (**4l**).

Although more data need to be accumulated for further discussion, we proceeded to explore the SAR around **4a**. The activities of compounds **5** and **6a–c** illustrate the influence of the linker between the quinoline and the heteroaryl rings. The sulfide moiety of **4a** can be substituted with a methylene (**5**) or can be removed (**6b**, **c**). An abrupt increase in potency was achieved through replacement of the 3,4-dimethoxyphenyl group of **6c** by a 3-isopropoxy-4-methoxyphenyl group (**6d**).

Compounds **2–6**, which were derived from justicidins as lead compounds and found to possess antiresorptive activity, were evaluated for anti-inflammatory activity using an adjuvant arthritis model in the rat. Compounds **4**, **5** and **6** had

Table 3. Physical and Spectral Data for Compounds **4c**, **g**, **i** and **m**

Compd.	mp (°C)	Solvent ^{a)}	¹ H-NMR (ppm, in CDCl ₃ , <i>J</i> in Hz)	IR (KBr, cm ⁻¹)	Formula	Anal. Calcd (Found)			Yield ^{b)} (%)
						C	H	N	
4c	145—146	EA-H	0.98 (3H, t, <i>J</i> =7.2), 3.80 (3H, s), 3.88 (3H, s), 3.97 (3H, s), 4.05 (2H, q, <i>J</i> =7.2), 4.06 (3H, s), 4.89 (2H, s), 6.91—7.02 (4H, m), 7.22 (1H, d, <i>J</i> =3.4), 7.45 (1H, s), 7.69 (1H, d, <i>J</i> =3.4)	1710	C ₂₆ H ₂₇ N ₃ O ₆ S	59.30 (59.10)	4.98 (5.03)	5.32 (5.45)	81
4g	119—120	EA-H	0.99 (3H, t, <i>J</i> =7.2), 3.51 (3H, s), 3.88 (3H, s), 3.98 (3H, s), 4.08 (2H, q, <i>J</i> =7.2), 4.65 (2H, s), 6.85—7.03 (4H, m), 7.10 (1H, s), 7.64—7.69 (2H, m), 7.96 (1H, d, <i>J</i> =9.6)	1716	C ₂₅ H ₂₄ ClN ₃ O ₄ S	60.30 (60.39)	4.86 (5.00)	8.44 (8.34)	66
4i	117—118	EA-H	0.96 (3H, t, <i>J</i> =7.2), 1.47 (3H, t, <i>J</i> =7.0), 3.43 (3H, s), 3.78 (3H, s), 4.02 (2H, q, <i>J</i> =7.2), 4.04 (3H, s), 4.11 (2H, q, <i>J</i> =7.2), 4.61 (2H, s), 6.86 (1H, s), 6.88 (1H, d, <i>J</i> =1.2), 7.00 (2H, d, <i>J</i> =8.8), 7.09 (1H, d, <i>J</i> =1.2), 7.26 (2H, d, <i>J</i> =8.8), 7.37 (1H, s)	1714	C ₂₇ H ₂₉ N ₃ O ₅ S	63.89 (64.01)	5.76 (5.88)	8.28 (8.36)	83
4m	137—138	A-E	0.93 (3H, t, <i>J</i> =7.2), 3.32 (3H, s), 4.04 (2H, q, <i>J</i> =7.2), 4.59 (1H, d, <i>J</i> =13.6), 4.70 (1H, d, <i>J</i> =13.6), 6.85 (1H, d, <i>J</i> =1.2), 7.10 (1H, d, <i>J</i> =1.2), 7.20—7.30 (2H, m), 7.36—7.60 (3H, m), 7.66 (1H, dd, <i>J</i> =9.0, 2.0), 7.97 (1H, d, <i>J</i> =9.0)	1739	C ₂₃ H ₁₉ C ₁₂ N ₃ O ₂ S	58.48 (58.20)	4.05 (4.12)	8.90 (8.73)	48

a) Recrystallization solvent, EA=ethyl acetate, H=hexane, A=acetone, E=Et₂O. b) Yield from the corresponding 2-chloromethyl quinoline derivative (see ref.7).

Table 4. Bone Resorption Inhibitory Effect of the Compounds Prepared^{a)}

Compd.	Conct. (μM)	⁴⁵ Ca release (% vs. control)	Compd.	Conct. (μM)	⁴⁵ Ca release (% vs. control)
2a	10	50***	4h	10	63**
2b	10	57**	4i	10	>80
2c	10	48***	4j	10	67*
3a	27 ^{b)}	74*	4k	10	>80
3b	26 ^{b)}	>80	4l	10	>80
3c	31 ^{b)}	59**	4m	10	>80
4a	30	57***	5	10	79**
4b	10	76*	6a	10	>80
4c	10	59***	6b	10	76**
4d	10	72*	6c	30	59**
4e	10	66***	6d	3	44**
4f	10	>80	1a	25	48**
4g	10	70***	(Justicidin A)		
			1b	25	47**
			(Justicidin B)		

a) Bone resorption inhibitory effects were evaluated by Raisz's method (see Biological Procedures). b) Actual assay was performed at a concentration of 10 μg/ml, which is equivalent to this μM value. Statistically significant at **p*<0.05, ***p*<0.01 and ****p*<0.001 by Student's *t*-test.

Table 5. Inhibitory Effect of **6c** and **6d** on FBS, hPTH and IL-1β Stimulated Release of ⁴⁵Ca from Fetal Rat Long Bones in Organ Culture

Stimulator	Compd.	Conct. (μM)	⁴⁵ Ca release (%) ^{a)}
FBS	6d	0	28.9±2.0 (100.0)
		0.3	23.1±2.3 (79.9)***
		1	14.8±1.3 (51.2)***
		3	10.8±0.3 (37.4)***
		30	17.4±1.6 (60.2)**
hPTH	6d	Basal	20.0±2.7 (58.3)**
		0	34.3±2.1 (100.0)
		0.5	23.6±1.3 (68.8)**
		1	17.8±1.3 (51.9)***
		30	21.8±3.2 (63.6)**
IL-1β	6d	Basal	16.9±1.7 (48.6)**
		0	34.8±3.3 (100.0)
		0.3	26.9±1.4 (77.3)
		1	20.0±1.3 (57.5)**
		30	19.1±0.4 (54.9)**

a) These effects were evaluated by Raisz's method (see Biological Procedures). Data are the mean±S.E. (*n*=5—6). Statistically significant at **p*<0.05, ***p*<0.01 and ****p*<0.001 by Student's *t*-test.

potent anti-inflammatory activity.⁷⁾ Among compounds possessing both antiresorptive and anti-inflammatory activities, **6c** (TAK-603) was selected for clinical studies based on its pharmacological profile.¹⁴⁾ Compound **6d**, which was synthesized during study of the metabolites of **6c**,¹⁾ exhibited reduced anti-inflammatory activity,¹⁵⁾ suggesting that some factors other than bone-resorbing function participate in adjuvant-induced inflammation.

Considering the biological results described above, two compounds, **6c** (TAK-603) and **6d**, were selected for detailed evaluation of antiresorptive activity. Ca release from bone induced by bone-resorbing factors,¹⁶⁾ human IL-1β, human parathyroid hormone (hPTH) and fetal bovine serum (FBS), was reduced by **6c** and **6d** (Table 5). Both compounds inhibited pit formation caused by pre-existing tartrate-resistant acid phosphatase (TRAP)-positive multinucleated cells (MNCs) on dentin slices.¹⁷⁾ Furthermore, new TRAP-positive MNC formation in culture dishes, without dentin slices, was also inhibited (Table 6). These results indicate that **6c** and **6d** inhibited both the activation of mature osteoclasts and the for-

mation of new ones.

As shown in Table 7, **6c** was effective in an *in vivo* model, ovariectomized mice. In this model, **6c** (TAK-603) had a preventive effect at an oral dose of 10 mg/kg/d.¹⁸⁾ Since the final stage of RA is bone destruction, the antiresorptive activity of **6c** (TAK-603) may be useful in controlling cartilage destruction. Concerning **6d**, *in vivo* studies using this model are presently under way.

In conclusion, we have found that quinoline derivatives bearing an azole moiety at the side chain of the 2-position have effects on bone metabolism as well as anti-inflammatory activity. Ethyl 4-(3,4-dimethoxyphenyl)-6,7-dimethoxy-2-(1,2,4-triazol-1-ylmethyl)quinoline-3-carboxylate (TAK-603), which had the most promising profile¹⁹⁾ in this series of compounds, is expected to be useful as a new type of DMARD. Although several approaches to elucidate the detailed mechanism of TAK-603 are now in progress, we are convinced that the antiresorptive activity of this compound will be one of its most characteristic features as an antirheumatic drug.

Table 6. Effect of **6c** (TAK-603) and **6d** on Pit Formation and New Osteoclast Formation by Mature Osteoclasts

	Conct. (M)	Pit formation		New osteoclast formation
		No. of pits/dentin slice ^{a)}	No. of TRAP(+)-MNCs/dentin slice ^{b)}	No. of TRAP(+)-MNCs/culture dish ^{c)}
Control		70.5±4.7	57.5±2.4	66.3±3.0
6c (TAK-603)	10 ⁻⁷	71.3±6.7	55.3±7.3	62.3±7.0
	10 ⁻⁶	47.0±7.0*	38.0±3.9**	46.8±3.2**
	10 ⁻⁵	31.8±5.0**	26.0±2.9**	23.0±3.4**
6d	10 ⁻⁹	69.0±6.0	50.8±5.8	67.8±3.2
	10 ⁻⁷	34.0±6.7**	41.3±3.8**	63.8±3.4
	10 ⁻⁵	2.5±1.0**	21.0±2.4**	13.0±2.9**

a) Effect on the number of resorption pits in mouse unfractionated bone cell cultures. b) Effect on the number of pre-existing TRAP-positive MNCs in mouse unfractionated bone cell cultures. c) Effect on the formation of new TRAP-positive MNCs in culture dishes without dentin slices. Data are the mean±S.E. (n=4). Statistically significant at **p*<0.05, ***p*<0.01 by Student's *t*-test.

Table 7. Effect of **6c** (TAK-603) on Changes in Femoral Bone Dry Weight in Ovariectomized Mice

	Ovx	Dose (mg/kg)	Femur dis. 1/3 (mg)	(%)
Sham-operated	—		13.0±0.2**	100.0**
Ovx Control	+		11.7±0.2	0.0
6c (TAK-603)	+	10	12.8±0.3*	86.7*

Data are the mean±S.E. (n=6—7). Statistically significant at **p*<0.05, ***p*<0.01 by Student's *t*-test.

Experimental

Chemistry Melting points were determined on a Yanagimoto micro-melting point apparatus and are uncorrected. Elemental analyses (C, H, N) were carried out by the Analytical Department of Takeda Chemical Industries, Ltd. ¹H-NMR spectra of deuteriochloroform (CDCl₃) or dimethyl sulfoxide (DMSO-*d*₆) solutions (internal standard tetramethylsilane (TMS), δ 0) were recorded on a Varian EM-390 (CW-60 MHz) or a Gemini-200 (FT-200 MHz) spectrometer. Infrared (IR) spectra were recorded on a Hitachi IR-215 spectrometer. All compounds exhibited ¹H-NMR, IR, and analytical data consistent with the proposed structures. Column chromatography was performed with E. Merck Silica gel 60 (0.063—0.200 mm).

General Procedure for Quinoline Lactones 3. 4-(3,4-Dimethoxyphenyl)-2-hydroxymethyl-6,7-methylenedioxyquinoline-3-carboxylic Acid Lactone (**3a**) A mixture of 2-amino-3',4'-dimethoxy-4,5-methylenedioxybenzophenone (200 mg, 0.66 mol), tetronic acid (80 mg, 0.79 mmol), concentrated H₂SO₄ (1 drop) and AcOH (3 ml) was stirred at 100 °C for 1 h, and then concentrated *in vacuo*. The residue was neutralized with saturated aqueous NaHCO₃, and extracted with CH₂Cl₂. The extract was washed successively with H₂O and brine, dried over MgSO₄, and concentrated *in vacuo* to give crystals. Recrystallization from CHCl₃-MeOH afforded **3a** as colorless needles (175 mg, 84%). ¹H-NMR (CDCl₃) δ: 3.90 (3H, s), 3.99 (3H, s), 5.18 (2H, s), 5.37 (2H, s), 6.95 (1H, d, *J*=2.0 Hz), 6.98 (1H, dd, *J*=8.0, 2.0 Hz), 7.06 (1H, d, *J*=8.0 Hz), 7.22 (1H, s), 7.46 (1H, s). IR (KBr) ν: 1768 cm⁻¹. Anal. Calcd for C₂₀H₁₅NO₆: C, 65.75; H, 4.14; N, 3.83. Found: C, 65.45; H, 4.06; N, 3.64.

4-(3,4-Dimethoxyphenyl)-2-hydroxymethyl-6,7-dimethoxyquinoline-3-carboxylic Acid Lactone (**3b**): Colorless prisms. ¹H-NMR (CDCl₃) δ: 3.85 (3H, s), 3.91 (3H, s), 4.01 (3H, s), 4.11 (3H, s), 5.39 (2H, s), 6.98—7.14 (3H, m), 7.23 (1H, s), 7.50 (1H, s). IR (KBr) ν: 1769 cm⁻¹. Anal. Calcd for C₂₁H₁₉NO₆: C, 66.14; H, 5.02; N, 3.67. Found: C, 66.13; H, 5.06; N, 3.71.

2-Hydroxymethyl-6,7-dimethoxy-4-phenylquinoline-3-carboxylic Acid Lactone (**3c**): Colorless prisms. ¹H-NMR (CDCl₃) δ: 3.82 (3H, s), 4.10 (3H, s), 5.39 (2H, s), 7.10 (1H, s), 7.40—7.65 (6H, m). IR (KBr) ν: 1758 cm⁻¹. Anal. Calcd for C₁₉H₁₅NO₄: C, 71.02; H, 4.71; N, 4.36. Found: C, 71.17; H, 4.68; N, 4.34.

3,4-Methylenedioxypropionyl 3',4'-Methylenedioxyphenylpropionate (9a) *N,N*-dimethylformamide (DMF) (1 drop) was added to a stirred mixture of 3,4-methylenedioxyphenylpropionic acid (2.0 g, 11.0 mmol), (COCl)₂ (1.7 g, 13.2 mmol) in tetrahydrofuran (THF) (10 ml) at room temperature. The re-

sultant mixture was stirred at room temperature for 1.5 h, and then concentrated *in vacuo*. A mixture of the residue obtained, pyridine (1.3 g, 16.0 mmol), 3,4-methylenedioxypropionyl alcohol (1.9 g, 11.0 mmol) and benzene (20 ml) was refluxed for 40 min. After cooling, the mixture was washed successively with 2*N* aqueous HCl, H₂O, saturated aqueous NaHCO₃ and brine, dried over MgSO₄, and concentrated *in vacuo*. The residue was chromatographed on SiO₂ with AcOEt-hexane (2:3) to give **9a** as pale yellow powder (1.4 g, 36%), mp 100—101 °C. ¹H-NMR (CDCl₃) δ: 4.84 (2H, d, *J*=7 Hz), 5.97 (2H, s), 6.02 (2H, s), 6.16 (1H, dt, *J*=16, 7 Hz), 6.63 (1H, d, *J*=16 Hz), 6.76 (1H, d, *J*=8 Hz), 6.80 (1H, d, *J*=8 Hz), 6.85 (1H, dd, *J*=8, 2 Hz), 6.95 (1H, d, *J*=2 Hz), 7.01 (1H, d, *J*=2 Hz), 7.17 (1H, dd, *J*=8, 2 Hz). IR (KBr) ν: 1710 cm⁻¹.

3,4-Methylenedioxypropionyl 3',4'-Dimethoxyphenylpropionate (9b): The title compound was prepared by the same method for **9a**. Yellow powder (yield: 44%), mp 113—115 °C. ¹H-NMR (CDCl₃) δ: 3.86 (3H, s), 3.90 (3H, s), 4.84 (2H, d, *J*=6 Hz), 5.93 (2H, s), 6.15 (1H, dt, *J*=16, 6 Hz), 6.64 (1H, d, *J*=16 Hz), 6.77—6.93 (4H, m), 7.07 (1H, d, *J*=2 Hz), 7.23 (1H, dd, *J*=9, 2 Hz). IR (KBr) ν: 1700 cm⁻¹.

3,4-Methylenedioxypropionyl 4'-Methoxyphenylpropionate (9c) A solution of *N,N'*-dicyclohexylcarbodiimide (DCC) (2.6 g, 12.7 mmol) in pyridine (9 ml) was added dropwise to a stirred mixture of 3,4-methylenedioxypropionyl alcohol (2.0 g, 11.5 mmol), 4-methoxyphenylpropionic acid (2.0 g, 11.5 mmol), *p*-TsOH·H₂O (120 mg, 0.63 mmol) and pyridine (29 ml) at room temperature. After stirring at room temperature for 5 h, AcOH (12 ml) was added, the mixture cooled to 0 °C, and stirred for a further 2 h. The insoluble product was filtered off, the filtrate was diluted with H₂O (100 ml), acidified with concentrated HCl, and then extracted with AcOEt. The combined extracts were washed successively with H₂O, saturated aqueous NaHCO₃ and brine, dried over MgSO₄ and concentrated *in vacuo* to afford **9c** as pale brown powder (72%), mp 100—102 °C. ¹H-NMR (CDCl₃) δ: 3.82 (3H, s), 4.84 (2H, d, *J*=6 Hz), 5.94 (2H, s), 6.15 (1H, td, *J*=16, 6 Hz), 6.65 (1H, d, *J*=16 Hz), 6.79 (1H, d, *J*=2 Hz), 6.87 (4H, d, *J*=9 Hz), 7.54 (2H, d, *J*=9 Hz). IR (KBr) ν: 1700 cm⁻¹.

General Procedure for Dihydronaphthalene Lactones 10. **3,4-Dihydro-3-hydroxymethyl-6,7-methylenedioxy-1-(3,4-methylenedioxyphenyl)-naphthalene-2-carboxylic Acid Lactone (10a)** A solution of **9a** (1.4 g, 4.0 mmol) in Ac₂O (25 ml) was refluxed for 13 h. The mixture was concentrated *in vacuo*, and the residue was chromatographed on SiO₂ with AcOEt-hexane (1:1) to give crystals. Recrystallization from AcOEt-EtOH afforded **10a** as pale yellow needles (1.1 g, 78%), mp 140—141 °C. ¹H-NMR (CDCl₃) δ: 2.77—2.95 (2H, m), 3.33—3.41 (1H, m), 4.00 (1H, t, *J*=9 Hz), 4.69 (1H, t, *J*=9 Hz), 5.97 (2H, s), 6.02 (2H, s), 6.52 (1H, s), 6.74—6.90 (4H, m). IR (KBr) ν: 1745 cm⁻¹. Anal. Calcd for C₂₀H₁₄O₆: C, 68.57; H, 4.03. Found: C, 68.35; H, 4.28.

1-(3,4-Dimethoxyphenyl)-3,4-dihydro-3-hydroxymethyl-6,7-methylenedioxy-naphthalene-2-carboxylic Acid Lactone (10b): Yellow prisms (yield: 34%), mp 225—226 °C (EtOH). ¹H-NMR (CDCl₃) δ: 2.78—2.92 (2H, m), 3.34—3.42 (1H, m), 3.87 (3H, s), 3.94 (3H, s), 4.01 (1H, t, *J*=9 Hz), 4.70 (1H, t, *J*=9 Hz), 5.97 (2H, s), 6.75—6.92 (5H, m). IR (KBr) ν: 1745 cm⁻¹. Anal. Calcd for C₂₁H₁₈O₆: C, 68.85; H, 4.95. Found: C, 68.88; H, 4.88.

3,4-Dihydro-3-hydroxymethyl-1-(4-methoxyphenyl)-6,7-methylenedioxy-naphthalene-2-carboxylic Acid Lactone (10c): Pale yellow prisms (yield: 53%), mp 233—234 °C (AcOEt). ¹H-NMR (CDCl₃) δ: 2.78—2.91 (2H, m),

3.37–3.42 (1H, m), 3.86 (3H, s), 3.99 (1H, t, $J=9$ Hz), 4.69 (1H, t, $J=9$ Hz), 5.96 (2H, s), 6.49 (1H, s), 6.77 (1H, s), 6.96 (2H, d, $J=10$ Hz), 7.24 (2H, d, $J=10$ Hz). IR (KBr) ν : 1740 cm^{-1} . Anal. Calcd for $\text{C}_{20}\text{H}_{16}\text{O}_5$: C, 71.42; H, 4.79. Found: C, 71.60; H, 4.67.

General Procedure for Naphthalene Lactones 2. **3-Hydroxymethyl-6,7-methylenedioxy-1-(3,4-methylenedioxyphenyl)naphthalene-2-carboxylic Acid Lactone (2a)** A mixture of **10a** (840 mg, 2.4 mmol), *N*-bromosuccinimide (NBS) (510 mg, 2.9 mmol), benzoyl peroxide (70 mg, 0.29 mmol) and CCl_4 (120 ml) was refluxed for 3 h, and the solvent was evaporated. The residue was chromatographed on SiO_2 with AcOEt–hexane (1:1) to give crystals. Recrystallization from AcOEt–acetone afforded **2a** as pale yellow prisms. $^1\text{H-NMR}$ (CDCl_3) δ : 5.38 (2H, s), 6.07 (2H, s), 6.09 (2H, s), 6.79 (1H, dd, $J=8, 2$ Hz), 6.81 (1H, d, $J=2$ Hz), 6.97 (1H, d, $J=8$ Hz), 7.12 (1H, s), 7.20 (1H, s), 7.70 (1H, s). IR (KBr) ν : 1760 cm^{-1} . Anal. Calcd for $\text{C}_{20}\text{H}_{12}\text{O}_6$: C, 68.97; H, 3.47. Found: C, 68.34; H, 3.51.

1-(3,4-Dimethoxyphenyl)-3-hydroxymethyl-6,7-methylenedioxy-naphthalene-2-carboxylic Acid Lactone (2b): Pale yellow needles. $^1\text{H-NMR}$ (CDCl_3) δ : 3.87 (3H, s), 3.98 (3H, s), 5.38 (2H, s), 6.08 (2H, s), 6.87 (1H, d, $J=2$ Hz), 6.91 (1H, dd, $J=8, 2$ Hz), 7.03 (1H, d, $J=8$ Hz), 7.13 (1H, s), 7.21 (1H, s), 7.70 (1H, s). IR (KBr) ν : 1750 cm^{-1} . Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{O}_6$: C, 69.23; H, 4.43. Found: C, 69.10; H, 4.40.

3-Hydroxymethyl-1-(4-methoxyphenyl)-6,7-methylenedioxy-naphthalene-2-carboxylic Acid Lactone (2c): Yellow prisms. $^1\text{H-NMR}$ (CDCl_3) δ : 3.91 (3H, s), 5.37 (2H, s), 6.08 (2H, s), 7.06 (2H, dd, $J=7, 2$ Hz), 7.11 (1H, s), 7.20 (1H, s), 7.28 (2H, dd, $J=7, 2$ Hz), 7.69 (1H, s). IR (KBr) ν : 1765 cm^{-1} . Anal. Calcd for $\text{C}_{20}\text{H}_{14}\text{O}_5$: C, 71.85; H, 4.22. Found: C, 71.52; H, 4.43.

Biological Procedures. Bone Resorption Inhibitory Effect The bone resorption inhibitory effect was determined by Raisz's method¹³⁾. ^{45}Ca (radioisotope of calcium in $^{45}\text{CaCl}_2$ solution) (50 μCi) was subcutaneously injected into a Sprague–Dawley rat on the 18th day of pregnancy. On the next day, the abdomen was opened, and a fetus was aseptically removed. The left and right humeri (radii and ulnae) were removed under a dissecting microscope, and as much connective tissue and cartilage as possible were also removed. Bone culture samples were then prepared. The bone was incubated in 0.6 ml BCJ_2 medium (Fitton-Jackson modification; GIBCO Laboratories) containing 2 mg/ml bovine serum albumin at 37°C for 24 h in an atmosphere of 5% CO_2 air. The bones were transferred to further fresh medium containing the appropriate final dose of test compound, with or without stimulator, 3% FBS or hPTH (1–34, 100 ng/ml) or IL-1 β (300 U/ml). Bone was cultivated for 2 d in the resulting medium. The ^{45}Ca radioactivity in the medium and the ^{45}Ca radioactivity in the bone were determined. The ratio (%) of ^{45}Ca released from the bone into the medium was calculated from the following equation:

$$\text{ratio of } ^{45}\text{Ca} \text{ released from bone into medium (\%)} = \frac{^{45}\text{Ca} \text{ released into medium}}{(^{45}\text{Ca} \text{ released into medium}) + (^{45}\text{Ca} \text{ incorporated in bone})} \times 100$$

Unfractionated Bone Cell Preparation Unfractionated bone cells were prepared according to the method of Takeda *et al.*¹⁷⁾ In brief, the femur and tibia of 11- to 13-day-old ICR mice (Japan Charles River, Tokyo, Japan) were aseptically isolated and minced with scalpel blades into 20 ml culture medium consisting of α -minimum essential medium (α -MEM) containing 5% FBS, 10 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (pH 7.0), 100 $\mu\text{g}/\text{ml}$ kanamycin, and 80 $\mu\text{g}/\text{ml}$ gentamicin. The mixture of cell suspension and bone fragments was gently pipetted for 5 min and allowed to stand for 5 min. The resulting supernatant was used for the experiments. To determine the number of osteoclasts in the supernatant, an aliquot was smeared on a slide glass and stained for TRAP, a marker enzyme for osteoclasts, according to the method of Burstone.²⁰⁾ TRAP-positive MNCs with three or more nuclei (roughly corresponding to 0.05% of the total cells) were counted as osteoclast-like cells.

Pit-Formation Assay Dentin slices of 0.1 mm-thickness and 6 mm-diameter were prepared by our previous method²¹⁾ and placed in a 96-well culture dish (Nunc; Nunc, Roskilde, Denmark). The unfractionated bone cell suspension (4×10^5 cells/dentin slice) with a TRAP-positive MNC density of 200 cells/dentin slice was seeded on to dentin slices and allowed to stand for 2 h at 37°C in a CO_2 incubator. After removal of nonadherent cells, the cells were further cultured for 4 d in medium containing several concentrations of test compound in the presence of 10^{-8} M $1\alpha,25$ -dihydroxyvitamin D_3 ($1\alpha,25(\text{OH})_2\text{D}_3$). Experiments were completed by removing the culture medium and adding 0.1 M cacodylate buffer solution (pH 7.4) containing 2% paraformaldehyde. The total number of TRAP-positive MNC on each dentin slice was counted after TRAP staining of the cells. Cells were

then removed from the dentin slices by ultrasonication for 30 s in distilled water, and air-dried. The slices were stained with hematoxylin and the number of densely stained pits was counted under light microscopy.

Osteoclast-Formation Assay²²⁾ The unfractionated bone cell suspension (2×10^5 cells/well) with a TRAP-positive MNC density of 100 cells/well was cultured in a 96-well plate without dentin slices in the absence of $1\alpha,25(\text{OH})_2\text{D}_3$ for 4 d. After depletion of TRAP-positive MNC was confirmed, the cells were further incubated for 6 d in culture medium containing test compound in the presence of 10^{-8} M $1\alpha,25(\text{OH})_2\text{D}_3$. The number of newly formed TRAP-positive MNC in the culture medium was then counted after TRAP staining of the cells. Our previous study demonstrated that pre-existing TRAP-positive MNC were depleted by culturing in the absence of $1\alpha,25(\text{OH})_2\text{D}_3$, but new TRAP-positive MNC formation is induced by the addition of $1\alpha,25(\text{OH})_2\text{D}_3$ in culture dishes even after complete depletion of TRAP-positive MNCs.

Effect on Changes in Femoral Bone Weight in Ovariectomized Mice Thirteen-week-old female C3H mice were ovariectomized by a dorsal approach under ether anesthesia. Three weeks later, the right femur of each mouse was removed and cleaned of soft tissue. The dry weight of the femur, which was cut one-third from the distal end, was determined after heating at 110°C for 3 h in an oven. **6c** was administered orally once a day for three weeks after the ovariectomy.

References and Notes

- 1) Part II: Baba A., Makino H., Ohta Y., Sohda T., *Chem. Pharm. Bull.*, **46**, 1130–1134 (1998).
- 2) a) McCulloch J., Lydyard P. M., Rook G. A., *Clin. Exp. Immunol.*, **92**, 1–6 (1993); b) Wick I., McColl G., Harrison L., *Immunol. Today*, **15**, 553–556 (1994).
- 3) Suzuki Y., Mizushima Y., *Osteoporosis Int.*, **7** (Suppl. 3), S217–S222 (1997).
- 4) a) Sohda T., Tsuda M., Naruse Y., Taketomi S., *Chem. Abstr.*, **118**, 17577j (1993); b) Sohda T., Tsuda M., Naruse Y., Taketomi S., *Jpn. Kokai Tokkyo Koho*, JP 04,211,609 (1992).
- 5) a) Lipsky P. E., Davis L. S., Cush D. J., Oppenheimer-Marks N., *Springer Semin. Immunopathol.*, **11**, 123–162 (1989); b) Gowen M., Mundy G. R., *J. Immunol.*, **136**, 2478–2482 (1986).
- 6) Anti-inflammatory activity of justicidin A (**1a**) in an adjuvant arthritic rat model: 10% inhibition of paw edema (50 mg/kg, *p.o.*, 14 d) (see ref. 7).
- 7) Baba A., Kawamura N., Makino H., Ohta Y., Taketomi S., Sohda T., *J. Med. Chem.*, **39**, 5176–5182 (1996).
- 8) A part of this work has been presented at AFMC International Medicinal Chemistry Symposium 95 (AIMECS 95), The Pharmaceutical Society of Japan, Tokyo, Japan, Sept. 3–8, 1995; Abstract p. 108.
- 9) a) Block E., Stevenson R., *J. Org. Chem.*, **36**, 3453–3455 (1971); b) Kohen F., Maclean J., Stevenson R., *J. Chem. Soc. (C)*, **1966**, 1775–1780.
- 10) Synthesis of this type of naphthalene lactone was reported by several groups: a) Arnold B. J., Mellows S. M., Sammes P. G., *J. Chem. Soc., Perkin Trans. I*, **1973**, 1266–1270; b) Horii Z., Tsujiuchi M., Kanai K., Momose T., *Chem. Pharm. Bull.*, **25**, 1803–1808 (1977); c) Momose T., Kanai K., Hayashi K., *ibid.*, **26**, 3195–3198 (1978); d) Silva S. O., Denis C. S., Rodrigo R., *J. Chem. Soc., Chem. Comm.*, **1980**, 995–997; e) Mann J., Piper S. E., Yeung L. K. P., *J. Chem. Soc., Perkin Trans. I*, **1984**, 2081–2088; f) Takano S., Otaki S., Ogasawara K., *Tetrahedron Lett.*, **26**, 1659–1660 (1985); g) Ishii Y., Ikariya T., Saburi M., Yoshikawa S., *ibid.*, **27**, 365–368 (1986); h) Seko S., Tanabe Y., Suzukamo G., *ibid.*, **31**, 6883–6886 (1990); i) Ogiku T., Seki M., Takahashi M., Ohmizu H., Iwasaki T., *ibid.*, **31**, 5487–5490 (1990); j) Kobayashi K., Kanno Y., Seko S., Sugimoto H., *J. Chem. Soc., Perkin Trans. I*, **1992**, 3111–3117; k) Hattori T., Tanaka H., Okashi Y., Miyano S., *ibid.*, **1995**, 235–241. l) Ogiku T., Yoshida S., Ohmizu H., Iwasaki T., *J. Org. Chem.*, **60**, 4585–4590 (1995); m) Tanabe Y., Seko S., Nishii Y., Yoshida T., Utsumi N., Suzukamo G., *J. Chem. Soc., Perkin Trans. I*, **1996**, 2157–2165. n) Padwa A., Cochran J. E., Kappe C. O., *J. Org. Chem.*, **61**, 3706–3714 (1996).
- 11) Cheng C.-C., Yan S.-J., *Org. React.*, **28**, 37–201 (1982).
- 12) a) Sohda T., Taketomi S., Baba A., *Eur. Pat. Appl.*, EP634169; b) Hitotsuyanagi Y., Kobayashi M., Fukuyo M., Takeya K., Itokawa H., *Tetrahedron Lett.*, **38**, 8295–8296 (1997).
- 13) a) Raisz L. G., *J. Clin. Invest.*, **44**, 103–116 (1965); b) Tsuda M., Kitazaki T., Ito T., Fujita T., *J. Bone Miner. Res.*, **1**, 207–211 (1986).
- 14) Makino H., Ohta Y., Baba A., Sohda T., *Rheumatoid Arthritis ID Re-*

- search Alert.*, **1**, 573—582 (1997).
- 15) Sohda T., co-workers, unpublished result. Anti-inflammatory activity of **6d** in an adjuvant arthritic rat model: <50% inhibition of paw edema (25 mg/kg, *p.o.*, 14 d).
 - 16) a) Suda T., Takahashi N., Martin T. J., *Endocr. Rev.*, **13**, 66—80 (1992); b) Akatsu T., Takahashi N., Udagawa N., Sato K., Nagata N., Moseley J. M., Martin T. J., Suda T., *Endocrinology*, **125**, 20—27 (1989); c) Akatsu T., Takahashi N., Udagawa N., Imamura K., Yamaguchi A., Sato K., Nagata N., Suda T., *J. Bone Miner. Res.*, **6**, 183—189 (1991).
 - 17) Takeda Y., Kusuda M., Hiura K., Sato T., Mochizuki H., Nagao Y., Tomura M., Yashiro M., Hakeda Y., Kawashima H., Kumegawa M., *Bone and Miner.*, **17**, 347—359 (1992).
 - 18) Mizoguchi J., Nishimura A., Notoya K., Baba A., Sohda T., The 13th Annual Meeting of the Japanese Society for Bone and Mineral Research (JSBMR), Fukuoka, Japan, Aug. 17—19, 1995; Abst. p. 355.
 - 19) Anti-inflammatory activity of **6c** in an adjuvant arthritic rat model: 65% inhibition of paw edema (12.5 mg/kg, *p.o.*, 14 d). ED₅₀: 2.6 mg/kg (see ref. 7).
 - 20) Burstone M. S., *J. Natl. Cancer Inst.*, **21**, 523—539 (1958).
 - 21) Notoya K., Yoshida K., Taketomi S., Yamazaki I., Kumegawa M., *Calcif. Tissue Int.*, **53**, 206—209 (1993).
 - 22) Notoya K., Yoshida K., Shirakawa Y., Taketomi S., Tsuda M., *Bone*, **16**, Suppl., 349S—353S (1995).