Ozonolysis of isodihydrohinesene (Va), obtained by formic acid dehydration of dihydrohinesol (II), afforded acetone, 2,6-dimethyl-bicyclo[5.3.0]decan-9-one (VII), $C_{12}H_{20}O$ (2,4-dinitrophenylhydrazones, m.p. $212\sim 214^\circ$), and 2,6-dimethyl-7-isobutyroylmethylcycloheptane-1-carboxylic acid (VIII), $C_{15}H_{26}O_3$ (semicarbazones, m.p. $203\sim 204^\circ$). When (II) was converted into the chloride (X) and treated with either aniline or ethanolic potassium hydroxide, isodihydrohinesene (Vb or Vc) was formed. Oxidation of (Vb) or (Vc) with ozone yielded, in addition to acetone, (VII), and (VIII), formaldehyde and 2,6-dimethyl-9-acetylbicyclo[5.3.0]decane (IX), $C_{14}H_{24}O$ (2,4-dinitrophenylhydrazones, m.p. $143\sim 144^\circ$). (IX) was reacted with methylmagnesium iodide to give an alcohol which was found to be identical with (II) in infrared absorption spectrum. These experiments furnished the evidence for the position of the tertiary alcoholic grouping to be in the isopropenyl side-chain. The infrared spectra of (IIIa-c) did not always indicate the strong absorption of vinylidene.

- 1) I. Yosioka, S. Takahashi, H. Iikino, Y. Sasaki : This Bulletin, **7**, 319 (1959).
- 2) I. Yosioka, H. Iikino, Y. Sasaki : *Ibid.*, **7**, 817 (1959).

borohydride. The oxidation of (I) was carried out with potassium permanganate to give the triol (XIV), m.p. 121.5~122°. When oxidized with periodate, (XIV) consumed only one equivalent of the reagent and did not produce acetone. From these facts it follows that (I) possesses the structure of 2,6-dimethyl-9-(1'-methyl-1'-hydroxyethyl)-bicyclo[5.3.0]dec-7-ene. Examinations on the configuration of compounds of this series are in progress.

Faculty of Pharmacy,
University of Osaka,
Toneyama, Toyonaka, Osaka-fu.

Itiro Yosioka (吉岡一郎)
Yasuko Sasaki (佐々木靖子)

Pharmaceutical Institute,
Faculty of Medicine,
Tohoku University,
Kita-4-bancho, Sendai.

Hiroshi Hikino (ヒキノヒロシ)

December 6, 1960.

UDC 577.164.12 : 577.159

On the Mechanism of Enzymic Conversion of 6,7-Dimethylribolumazine to Riboflavin

The writers previously reported that a crude enzyme prepared from the mycelium of *Eremothecium ashbyii* reacted with 6,7-dimethylribolumazine to produce 6-methyl-7-hydroxyribolumazine besides riboflavin¹⁾ and that the reaction yielded a considerable amount of riboflavin without any carbon donor.²⁾ The greater part of the products in this reaction were identified by paper partition chromatography and, of the two unidentified spots, the bluish green fluorescent one at R_f 0.10 was found to resemble the polymer consisting of two moles of 4-ribitylamino-5-aminouracil³⁾ reported by the present writers.

Recently, Plaut⁴⁾ discussed the conversion of 6,7-dimethylribolumazine to riboflavin by means of the enzyme of *Ashbya gossypii*, the gist of which is as follows: 1) The enzyme acts on 6,7-dimethylribolumazine to produce riboflavin without any carbon donor; 2) a C₄ compound is split off from the pyrazine ring of 6,7-dimethylribolumazine and then reacts with another 6,7-dimethylribolumazine to produce riboflavin. In other words, 6,7-dimethylribolumazine acts as a carbon donor as well as a carbon acceptor; 3) one mole of riboflavin is produced from two moles of 6,7-dimethylribolumazine; 4) when the methyl groups at the 6- and 7-positions of 6,7-dimethylribolumazine are labeled with ¹⁴C, the molar specific radioactivity of the resulting riboflavin becomes twice as strong as that of 6,7-dimethylribolumazine; 5) decomposition by the Kuhn-Roth method of the riboflavin thus obtained gives two moles of acetic acid containing ¹⁴C, thereby revealing the distribution of radioactivity in riboflavin. In another paper,⁵⁾ Plaut stated that formation of 1 mole of riboflavin requires 3 moles of 6,7-dimethylribolumazine and that 6-methyl-7-hydroxyribolumazine was a product other than riboflavin, but did not mention the amount of the resulting 6-methyl-7-hydroxyribolumazine in the above-mentioned report.⁴⁾

The writers prepared an enzyme solution from *Er. ashbyii* and allowed the product to react with 6,7-dimethylribolumazine as reported before. A part of the reaction mixture

- 1) S. Kuwada, T. Masuda, T. Kishi, M. Asai : This Bulletin, **6**, 618 (1958).
- 2) S. Kuwada : Vitamins (Kyoto), **14**, 933 (1958).
- 3) S. Kuwada, T. Masuda, T. Kishi, M. Asai : This Bulletin, **8**, 798 (1960).
- 4) G.W.E. Plaut : J. Biol. Chem., **235**, PC 41 (1960).
- 5) *Idem* : Federation Proc., **19**, 312 (1960).