UDC 547.834.3.07

46. Masao Shimizu and Fumihiko Uchimarū : Studies on N-Substituted Nortropane Derivatives. I. Synthesis of N-Substituted 3-Nortropanones by the Robinson-Schöpf Condensation.

(Research Laboratory, Daiichi Seiyaku Co., Ltd.*¹)

3-Tropanone (tropinone*²), the starting material for synthesis of tropane alkaloids was synthesized by Robinson¹⁾ by a one-step process in 1917, and later Schöpf²⁾ modified this method under biological condition. These works were the first of the hypotheses in the field of alkaloid biogenesis. After that, this condensation method was applied to the syntheses of various alkaloids, e.g. meteloidine,³⁾ valeroidine,⁴⁾ etc. The authors have attempted to synthesize various N-substituted nortropane derivatives in order to compare their pharmacological action with that of natural tropane alkaloids. In this paper, the Robinson-Schöpf condensation using the esters of α -amino acids is mainly described.

The Robinson-Schöpf condensation consists in preparing 3-tropanone with succindialdehyde, acetonedicarboxylic acid, and methylamine in a comparatively good yield under biological conditions of temperature, dilution, and pH. It has been reported that alkyl,⁵⁾ hydroxy-alkyl,⁵⁾ benzyl,⁵⁾ and aromatic and heterocyclic amines⁶⁾ are used as the basic component in this condensation, but no case has been known to use the esters of α -amino acids. For the conditions of condensation referring to the descriptions of Keagle and Hartung,⁵⁾ and Stoll, *et al.*,⁶⁾ syntheses of 3-tropanone with methylamine hydrochloride, 3-nortropanone⁷⁾ with ammonium chloride, and 8-(2-hydroxyethyl)-3-nortropanone with ethanolamine were examined. Among the other components, acetonedicarboxylic acid was prepared from citric acid by the usual method.⁸⁾ For preparation of succindialdehyde, the method from pyrrole⁹⁾ and furan⁶⁾ has been reported but the synthesis of its acetal from acetylene was effectively adopted according to the method of Preobrashenski¹⁰⁾ or Wibaut.¹¹⁾ The method is effected in a fairly good yield although the intermediate, acetylenedicarboxaldehyde bis(diethylacetal), is somewhat unstable (Tables I and II). While reduction in the last step was made at 20°

Chart 1.

*¹ Hirakawabashi, Sumida-ku, Tokyo (清水正夫, 内丸文彦).

*² For traditional nomenclature see G. Fodor, K. Nádor : J. Chem. Soc., **1953**, 721. The present one follows the I. U. P. A. C. rules.

- 1) R. Robinson : J. Chem. Soc., **111**, 762, 876 (1917).
- 2) C. Schöpf, G. Lehmann : Ann., **518**, 1 (1935); C. Schöpf : Angew. Chem., **50**, 779, 797.
- 3) C. Schöpf, W. Arnold : Ann., **558**, 109 (1946).
- 4) A. Stoll, B. Becker, E. Jucker : Helv. Chim. Acta, **35**, 1263 (1952).
- 5) L. C. Keagle, W. H. Hartung : J. Am. Chem. Soc., **68**, 1608 (1946).
- 6) A. Stoll, E. Jucker, A. Lindenmann : Helv. Chim. Acta, **37**, 495 (1954).
- 7) A. Lindenmann : *Ibid.*, **42**, 490 (1959).
- 8) Org. Syntheses, **I**, 10.
- 9) G. L. Ciamician, M. Dennstedt : Ber., **17**, 533 (1884); C. Harries, *et al.* : *Ibid.*, **34**, 1448 (1901); **35**, 1183 (1902); **39**, 3670 (1906); **41**, 255 (1908); R. Willstätter, W. Heubner : *Ibid.*, **40**, 3871 (1907); C. Mannich, H. Budde : Arch. Pharm., **270**, 283 (1932).
- 10) N. A. Preobrashenski, I. A. Rubtsov, T. F. Dankova, V. P. Pavlov : J. Gen. Chem. U. S. S. R., **15**, 952 (1945) (C. A., **40**, 6488 (1946)).
- 11) J. P. Wibaut, A. L. van Hulsenbeck, C. M. Siegmans : Proc. Koninkl. Ned. Akad. Wetenschap., **53**, 989 (1950) (C. A., **45**, 5167 (1951)).

TABLE I. Syntheses of Acetylenedicarboxaldehyde Bis(diethylacetal)

Expt. No.	EtMgBr (mole)	Reaction period with		Yield* (%)
		Acetylene (hr.)	HC(OEt) ₃ (hr.)	
1	0.58	11	7.5	34.6
2	1.75	11	6.0	69.7
3	1.76	10	7.0	74.3
4	2.95	10	8.0	83.4
5	2.95	10	8.0	82.8
6	2.95	9	8.5	79.6

* The yield was calculated on the amount of reacted ethyl orthoformate.

TABLE II. Syntheses of Succindialdehyde Bis(diethylacetal)

Expt. No.	Starting material (g.)	H ₂		Raney nickel		Product	
		mole	time (hr.)	(cc.)	(days)*	n _D (°C)	Yield (%)
1	12.6	2.15	8	2	1	1.4180 (26)	84.3
2	13.0	2.08	17	2+2	9	1.4216 (26)	77.9
3	20.7	2.05	19	4+2	2	1.4226 (21)	91.6
4	60.0	2.17	11	12+4	9	1.4230 (16)	91.7
5	111.0	2.25	18	10+4	1	1.4214 (16)	90.2
6	49.1	2.55	10	10	9	1.4224 (16)	88.3

* Number of days after preparation of the catalyst. The speed of hydrogen absorption differs markedly between 1 and 2 moles while the catalyst is fresh.

and 100 atm. pressure by Wibaut¹⁰ the desired product was obtained in about 90% yield using Raney Nickel W-5 as a catalyst at room temperature and normal pressure. The acetal is a stable compound and can be preserved for a long time without any appreciable change.

The condensation reaction was carried out as follows: To a succindialdehyde solution, prepared from the bis-diethylacetal by shaking with 0.5% sulfuric acid solution, acetonedicarboxylic acid and amine hydrochloride were added, and buffered with sodium acetate to pH 4.6~5.2. After dilution with water, the mixture was shaken at room temperature for several days. The longer the shaking time, the better was the yield. The reaction mixture was basified with potassium hydroxide and extracted exhaustively with chloroform. The yield was calculated on the amount of eluate from alumina chromatography. All the products of attempted condensation showed strong infrared absorptions ascribed to a six-membered ring ketone.*³ Examples of condensation carried out are shown in Table III.

Chart 2.

The structures of 8-(1-alkoxycarbonylalkyl)-3-nortropanones obtained in this work were further confirmed by identification with the known compounds derived from 3-tropanone (tropinone). These will be reported in a subsequent paper.

*³ More detailed results will be shown in a subsequent paper.

TABLE III. Robinson-Schöpf Condensation

Expt. No.	Amine R-NH ₂	Acetal (g.)	Shaking time (hr.)	Reaction time (days)	Eluted product	
					(g.)	(%)
1	NH ₄ •Cl	2.4	10	7	0.58	45.2
2		4.8	11	3	1.84	64.6
3	Methylamine hydrochloride	9.6	33	5	4.17	73.2
4		28.8	40	11	11.0	64.3
5		19.2	40	6	8.00	70.2
6		28.8	32	6	14.2	83.2
7	Ethanolamine	2.4	2	3	0.62	36.2
8		4.8	1.5	4	0.87	25.4
9		9.6	19	4	4.74	69.3
10	Glycine methyl ester hydrochloride	9.6	33	6	6.16	76.3
11	Glycine ethyl ester hydrochloride	2.4	0	3	0.64	29.6
12		6.0	14	4	2.25	41.6
13		9.6	22	4	6.00	69.3
14		4.8	22	5	2.73	63.2
15	Glycine benzyl ester hydrochloride	2.4	27	4	1.12	42.2
16	DL-Alanine ethyl ester hydrochloride	2.4	25	4	1.51	65.5
17	DL-Methionine ethyl ester hydrochloride	2.4	19	4	2.70	92.4

Experimental**

Succindialdehyde Bis(diethylacetal)—EtMgBr, prepared from 42 g. of Mg and 192 g. of EtBr in Et₂O, was bubbled with purified acetylene for 11 hr. and 270 g. of ethyl orthoformate was added. The resulting mixture was reacted in benzene for 6 hr. at 55~60° (inner temp.). The reaction product was decomposed by adding cold, satd. solution of AcONH₄, extracted with Et₂O, and the extract was dried over Na₂SO₄ and evaporated. The residue was distilled *in vacuo* and yielded 67 g. (24.8% recovery) of ethyl orthoformate and 110 g. (69.7%) of colorless liquid, b.p.₄ 109~111°. This liquid was immediately hydrogenated over Raney Ni W-5 (14 cc.) in EtOH at room temperature and 2.25 mol. equiv. of H₂ was taken up in 18 hr. After filtration of the catalyst and evaporation of the solvent, the residue was distilled *in vacuo* to leave 101.2 g. of colorless liquid, b.p.₆ 97~101° (90.2%). n_D^{16} 1.4214. IR $\nu_{\max}^{liq.}$ cm⁻¹: 1125, 1063 (ν_{C-O}).

3-Tropanone—To a solution of succindialdehyde, prepared by shaking 9.6 g. of bis-diethylacetal with 0.5% H₂SO₄ for 2 hr., methylamine hydrochloride (5.4 g.), acetonedicarboxylic acid (11.7 g.), and AcONa (21.8 g.) were added, the volume of the mixture was made up to 400 cc. (pH 4.0) by dilution with distilled water, and the mixture was allowed to stand at room temperature for 5 days with occasional shaking (total, 33 hr.) (pH 4.4). The resulting solution was made alkaline with KOH while cooling, saturated with NaCl, and extracted thoroughly with CHCl₃. The extract was dried over Na₂SO₄ and evaporated *in vacuo*. The residue (13.0 g.) was chromatographed on alumina (100 g.) and the fraction (5.75 g.) eluted with Et₂O crystallized to plates (73.2%), m.p. 40~43°, b.p.₉ 97~100°.

Picrate: Yellow needles (from H₂O), m.p. 217~219°. Anal. Calcd. for C₁₄H₁₆O₈N₄: C, 45.65; H, 4.38; N, 15.21. Found: C, 46.17; H, 4.61; N, 15.39.

Methiodide: Colorless prisms (from MeOH-Me₂CO), m.p. 273~274° (decomp.). Anal. Calcd. for C₉H₁₆ONI: C, 38.45; H, 5.74; N, 4.98. Found: C, 38.53; H, 5.61; N, 4.48.

N-Substituted 3-Nortropanones—The 3-tropanones listed in Table III were prepared in accordance with 3-tropanone.

3-Nortropanone—Picrate: Yellow plates (from H₂O), m.p. 167~169°. Anal. Calcd. for C₁₃H₁₄O₈N₄: C, 44.07; H, 3.98; N, 15.82. Found: C, 44.28; H, 4.04; N, 15.89.

8-(2-Hydroxyethyl)-3-nortropanone—Faintly yellow oil, b.p.₄ 150~153°.

Picrate: Yellow prisms (from H₂O), m.p. 172~174°. Anal. Calcd. for C₁₅H₁₈O₉N₄: C, 45.23; H, 4.55; N, 14.07. Found: C, 45.07; H, 4.44; N, 14.14.

Methiodide: Colorless plates (from MeOH-Me₂CO), m.p. 215~217°. Anal. Calcd. for C₁₀H₁₈O₂NI: C, 38.60; H, 5.84; N, 4.50. Found: C, 38.48; H, 5.70; N, 4.66.

** All m.p.s are uncorrected.

Methyl 3-Oxo-8-nortropaneacetate—Picrate : Yellow prisms (from H₂O), m.p. 159~160°(decomp.). *Anal.* Calcd. for C₁₆H₁₈O₁₀N₄ : C, 45.07; H, 4.26; N, 13.14. Found : C, 44.78; H, 4.29; N, 13.52.

Ethyl 3-Oxo-8-nortropaneacetate—Picrate : Yellow prisms (from H₂O), m.p. 146~147°. *Anal.* Calcd. for C₁₇H₂₀O₁₀N₄ : C, 46.36; H, 4.58; N, 12.72. Found : C, 46.52; H, 4.59; N, 12.71.

Benzyl 3-Oxo-8-nortropaneacetate—Picrate : Yellow plates (from iso-PrOH), m.p. 162~163°. *Anal.* Calcd. for C₂₂H₂₂O₁₀N₄ : C, 52.59; H, 4.41; N, 11.15. Found : C, 52.41; H, 4.23; N, 11.34.

Ethyl (±) 3-Oxo-α-methyl-8-nortropaneacetate—Picrate : Yellow needles (from MeOH-Et₂O), m.p. 133~135°. *Anal.* Calcd. for C₁₈H₂₂O₁₀N₄ : C, 47.58; H, 4.80; N, 12.33. Found : C, 47.53; H, 5.07; N, 12.31.

Ethyl (±) 3-Oxo-α-(2-methylthioethyl)-8-nortropanone—Free base : Yellow oil, S, positive, somewhat unstable, and colors on exposure to air. Picrate and methiodide did not crystallize. IR $\nu_{\text{max}}^{\text{liq.}}$ cm⁻¹: 1725 (six-membered ring ketone), 1739 (ester).

The authors express their deep gratitude to Dr. Junzo Shinoda, President of this Company, and to Dr. Takeo Ishiguro, Director of this Laboratory, for kind encouragement. The authors are indebted to Mr. S. Okada for technical assistance and to Messrs. B. Kurihara and K. Abe for elemental analyses.

Summary

Several new N-substituted 3-nortropanones were prepared by the Robinson-Schöpf condensation using the esters of α-amino acids. The compounds obtained were characterized as their picrates after purification by alumina chromatography and their structure confirmed through their infrared spectra.

(Received July 6, 1960)