CHAPTER 14 Concluding Remarks

Mitotic aberrations induced experimentally take precedence over those found in nature: Aberrant mitoses found in nature in special organisms such as parasites, protozoa, some insects or in highly differentiated organs or tissues are described in the work of Bělař (1928), Schrader (1951) and Cleveland (1963). These mitoses are frequently referred to as clues for elucidation of the mechanism of mitosis by animal cytologists.

Such mitotic aberrations are acquired characters in the phylogenetic development of such organisms under special circumstances or under highly specialized habitats utterly unknown at present, and their aberrations might have been induced by mutations as being the most adequate characters for the maintenance of life. Therefore, without analyzing the causes and conditions under which the organisms have to perform peculiar and unique mitoses, it may be of little significance to employ such specialized cell divisions for analyzing the mechanism of ordinary mitosis. In my opinion, in order to understand the behavior of mitosis, the aberrations should be induced artificially by given experimental conditions and be traced by *in vivo* observations.

Expected functions of spindle membrane and spindle poles at the molecular level: Before we consider the function of spindle poles, their anatomical basis must be clarified at the submicroscopic level. Unfortunately, commonly accepted ideas on the mitotic figures in classical cytology, and even current cytology text-books, have disregarded the important biological significance of the intracellular membranous structures. By routine cytological techniques, it is difficult in practice to discern the spindle membrane either in fixed preparations or in *in vivo* observations but it is another story now; by applying improved fixatives the spindle membrane, including the active regions of its poles, is preserved in fixed preparations and gets to be analyzable under the electron microscope.

A new blueprint for mitosis: In this book, based on evidence backed by *in vivo* observations and by improved fixatives, we present a new blueprint for mitosis. Studies concerning Physiology, Biochemistry and Biophysics of mitotic cells, prevalently being undertaken should be re-investigated on the basis of the new blueprint for mitosis as explained in this book.

As already mentioned above, the blueprints hitherto on mitosis have been made on the basis of data depending exclusively on the static morphology of fixed mitotic cells. Therefore, the anatomical basis of mitotic organelles in classical cytology are not only incomplete but also contain gross errors. Moreover, in fixed preparations the dynamic aspects of mitosis are inevitably explained by the researchers' speculations and by

preconceived ideas. In order to analyze the kinetics of each mitotic organelle, researchers should recommence and reinvestigate extensively the new blueprint for mitosis without sticking to the currently common ideas on cell division. At present, intracellular membranous structures, i.e. interior compartmentalization of the cell, present many important biological problems on cellular physiology. Nevertheless, the existence of the spindle membrane in higher organisms is still disregarded even in current cytology text-books. Similarly, the existence of atractoplasm and its entity as distinguished from the cytoplasm which was established by Fujii in 1931, and the existence of the spindle surface membrane by Wada in 1935, have not been generally accepted by cytologists.

Studies on cytomorphology in the future: Concerning the fine structure of the spindle body, especially that of the spindle membrane, investigations have just commenced to take shape in research; solution of many new and important problems will be expected in future research. The effectiveness of fixatives, the reaction of metal ions of various kinds and concentrations should be quantitatively reinvestigated by researchers in order to find out the most adequate fixative specific to the material they use.

Cadmium chloride is effective for generally preserving the spindle membrane in plant mitoses but it is by no means omnipotent for preservation of the spindle membrane of various kinds of organisms. In classical cytology, a tendency to use the fixatives recommended by pioneers, for instance the usage of OsO_4 or KMnO_4 solutions, has in a certain sense been a real obstacle to progress in cytomorphology. In order to judge correctly the truth of fixed preparations, it is my own long-cherished opinion that the effect of reagents or that of ions contained in fixatives against intracellular organelles should be, at first, given preliminary examination by *in vivo* observations. For instance, "*Tradescantia* test *in vivo*" can be used for this purpose and this test also often furnish conclusive results.

In order to analyze mitosis at molecular level: The mechanism of mitosis has been interpreted hitherto by animation of a series of fixed mitotic figures and recently by phase contrast cine-micrographs which only show mainly shadow pictures of chromosomes. These explanations have failed to recognize an important aspect that mitosis is carried out by continuous changes of cause-and-effect relations among mitotic organelles.

Apart from the aspect of metabolism in mitotic cells, the morphological changes of mitotic organelles constitute a "closed system" during a mitotic cycle, i.e. when chromosomes move, the atractoplasm also moves, replacing their positions with each other in the spindle. Therefore, without analyzing the dynamic morphology of each mitotic organelle in living cells, the continuous changes occurring in a mitotic cell cannot be understood correctly. Then, without establishing a sequence of causes and effects among the changes of mitotic organelles, the physiology of the mitotic cell cannot be

elucidated significantly according to physical laws at the submicroscopic level. Finally, without considering physicochemical changes which occur in the intracellular membranous system, especially in the spindle membrane, the solution of the mechanism of mitosis will not be achieved by introducing present knowledge of cellular physiology at the molecular level (Diagram 1).

In the blueprint for mitosis described in this book, some of the common ideas on the mitotic figures described in current cytology text-books have been drastically revised when they are inconsistent with the results confirmed experimentally by *in vivo* observations and by improved fixation techniques. At present, however, the descriptions in this book are only indications of the outline and some incipient steps toward the solution of various problems on mitosis according to laws of physics as well as to modern ideas on cell biology at the molecular level. With regard to further elucidation of the mechanism of mitosis, we expect much from future development of research and to work by many competent cytologists who can grasp the phenomena insofar as they are free from any preconceived ideas.

Addendum: During the proof reading of this book, a monograph written by Went appeared:—Went, H. A. 1966. The Behavior of Centrioles and the Structure and Formation of the Achromatic Figure. Protoplasmatologia VI G1, Springer-Verlag, Wien and New York. This monograph concentrates on descriptions of the structure and formation of the mitotic apparatus. As Went stated in the introduction, the mitotic apparatus is responsible primarily for the equipartition of the hereditary material. Strictly speaking, this event is accomplished in the karyokinetic spindle. The concept of the mitotic apparatus, however, does not define clearly the entity of the karyokinetic spindle as distinguished from the cytokinetic one (See pages 41–42 of this book). The equipartition of the hereditary material is a common event occurring both in higher plants and animals, but the centrioles are specialized cytoplasmic organelles in animal cells and their behavior has nothing to do with plant karyokinesis, and even with animal karyokinesis as well. The role of the centrioles may be restricted to a kinetic center for animal cell cytokinesis (See pages 66–67 of this book).