特別講演 1 DESIGN OF DRUGS INVOLVING THE CONCEPT AND THEORIES OF DRUG METABOLISM AND PHARMACOKINETICS

Dennis A. Smith

*Department of Drug Metabolism, Pfizer Central Research,
Sandwich, Kent CT13 9NJ, U.K.*

Over the last 25 years drug metabolism has developed concepts and theories that predict or rationalise the behaviour of a compound based on physicochemistry and structure. These concepts, that rely heavily on the understanding of free (unbound) drug concentrations and how this relates to drug activity are explored below for existing agents, including beta-adrenoceptor antagonists and calcium channel antagonists, in terms of dissolution and absorption, clearance, half life and potency, and distribution to the target.

Design and Dissolution and Absorption

Absorption across the gastrointestinal tract normally involves the dissolution of a compound in the aqueous contents of the gastrointestinal tract followed by the diffusion of the molecule through the apical and basolateral membranes of the cells lining the gastrointestinal tract (transcellular absorption). To traverse this route, the compound has to have adequate water solubility for the dissolution process to occur and sufficient lipophilicity to 'dissolve' within the lipid core of the cell's membranes. This balance means that many drugs occupy a log D range from 0 to 2 to combine these solubility properties. If a molecule has sufficient lipophilicity, the large surface area of the microvilli means that absorption is rapid. Tight junctions formed by the proteins that act as the junction between cells, offer an aqueous channel (pore) for paracellular absorption. These pores are estimated to be a size of 3 to 10 Å and represent only 0.01% of the total absorptive surface area. Beta-adrenoceptor antagonists being small molecules can make efficient use of this route. Thus atenolol and nadolol are absorbed 56% and 34% respectively in man. More lipophilic compounds such as metoprolol, propranolol, betaxolol and alprenolol or the calcium channel antagonists are able to cross the gastro-intestinal tract by the transcellular route. In traversing by this transcellular route the compounds may encounter the effects of gut extraction due to P-glycoprotein (PgP) and CYP450 enzymes. PgP works as an efflux pump and prevents entry of compounds into the blood stream. The effects of CYP450 enzymes present in the gut may be enhanced by a "recirculation" effect of PgP efflux and reabsorption. It is likely that compounds with relatively lower rates of diffusion across the gastro-intestinal tract will show the effects of PgP and CYP450 metabolism the greatest. Within the beta-adrenoceptor antagonist series talinolol has a considerably reduced bioavailability due to it being an excellent PgP substrate [1]. No simple structure activity relationships exist when the series is examined, although it is noteworthy that para substitution does not confer PgP affinity when the chain is a simple alkoxy as for metoprolol or betaxolol but does when it is a urea as for talinolol. Gut extraction lowers bioavailability and hence increased dose size. It also may be a major factor in non-linear pharmacokinetics due to the high concentrations that occur locally during drug absorption and the likelihood of some degree of saturation occurring over the clinical dose range. Designing molecules for good absorption requires the compound to be small in size or in the mid ground of lipophilicity. The effects of PgP are not highly predictable and are best screened for (2). The experience with talinolol suggests hydrogen bonding functionality that is not essential for selectivity or potency should be avoided.

Design and Clearance, Volume and Potency

Yamada [3] has illustrated how the pharmacokinetics of the beta-adrenergic antagonists combined with daily dose and affinity yield the same degree of receptor occupancy (80%) and hence efficacy. Another feature of this range of compounds is their lipophilicity, as measured by log D_{7.4}

or log P spans four log units. Since lipophilicity is a key parameter in determining affinity for receptors (potency), enzymes (clearance), tissues and membranes (volume) and permeability (tubular reabsorption) it is illuminating to examine the various trends in these parameters changes in lipophilicity invoke. Potency shows a marked correlation with lipophilicity, in that the pA_2 value demonstrates a 300 fold increase in potency with a 10,000 fold increase in lipophilicity (D or P). Such changes can be simply rationalised considering the features of the natural agonists (eg adrenaline) and the key H-bonding and ion pair binding interactions which occur. Any increase in potency in the series is likely to come from non-specific lipophilic interactions.

The major component of volume of distribution is the affinity for the membranes of cells. This affinity is derived from two physiochemical properties: the compounds intrinsic lipophilicity and its basicity. Membranes are comprised of phospholipids with changed head groups, some acidic, and lipid tails organised as a bilayer to form a lipid interior. Lipophilicity determines the compounds ability to interact with the lipid core of the membrane, whilst a basic centre allows ion pair interactions with the ionised head groups. Since the beta-adrenergic antagonists all possess a similar basic secondary amine group (pK_a 9.4), and all are capable of forming ion pairs with the head groups, then lipophilicity becomes the main discriminator in determining volume. A strong correlation between unbound (free) volume and lipophilicity is in keeping with these interactions.

Unbound renal clearance is approximately constant across the beta-adrenergic antagonists. The amount excreted in urine declines dramatically with lipophilicity above $\log D$ 0, due to the increased importance of metabolic clearance, making calculations unreliable. Intrinsic unbound clearance ($Cl_i(u)$) measures the metabolism of the compounds and is derived from the V_{max}/K_m for the enzyme system. With cytochrome P-450 enzymes it is not surprising that K_m shows an inverse relationship to lipophilicity since the binding of the substrate to the enzyme relies largely on hydrophobic forces [4,5]. This is so even for CYP2D6, an enzyme which also interacts by ion pair interactions [5] and is a key enzyme in the clearance of several of the beta-adrenoceptor antagonists. Rate of metabolism is a function of chemical lability and the ability to enter and leave the active site of the enzyme. Actual functionality is important here, rather than just the lipophilicity of the substituent. Comparison can be made between metoprolol and betaxolol. Metoprolol has an ethoxy-methyl para substituent that renders the compound liable to metabolism at both the benzylic and methyl position. In contrast betaxolol has an ethoxy-methyl cyclopropyl substituent that is sufficiently bulky to prevent efficient binding of the compound to the CYP2D6 active site [6]. These changes are reflected in the $Cl_i(u)$ values of metoprolol and betaxolol of 35 and 10 ml/min/kg respectively despite betaxolol being 3-4 fold more lipophilic.

These differences in metabolism are very important since the overall trend is that much of the chemistry around the basic aryloxy-propan-2-ol amino unit has the effect of changing potency and key pharmacokinetic properties in a compensating manner. Increasing lipophilicity raises both V_d and $Cl_i(u)$ effectively cancelling out any changes in half-life. Moreover the daily dose size is governed by the unbound steady state concentration of unbound drug and potency. $Cl_i(u)$ mostly determines the steady state concentration of orally administered drug. Both $Cl_i(u)$ and potency increase with lipophilicity again effectively cancelling each other out. These inter-relationships illustrate why it is bad practice to optimise on only one property, such as in vitro potency. Rationale design, such as moving from the cardio selective, moderately potent (pA_2 , 7.8) but metabolically labile metoprolol, to the metabolically stable betaxolol are highly productive. The advantages of metabolic stability are emphasised because the 3 extra methylene units of the replacement function give the expected increase in lipophilicity and resultant increase in potency (pA_2 , 8.8) and volume [5 cf 16 L/kg; $V_d(u)$]. The metabolic stability of the functionality leads to a reduction in clearance against the expected trend, and increases in volume result in an extension of half-life from 3 to around 16 hours.

The dihydropyridine calcium channel antagonists also show strong trends between increasing lipophilicity and increasing unbound intrinsic clearance ($Cl(u)$) and unbound volume ($Vd(u)$). The relationship for $Cl(u)$ and lipophilicity can be rationalised by oxidation of the dihydropyridine ring, via an electron abstraction process, being a common primary step. This oxidation is conducted by CYP3A4 an enzyme whose interactions are primarily lipophilic (4,5). The relationship with volume and lipophilicity reflects the interaction with the predominantly neutral molecules and the lipid portion of biological membranes. The volume/lipophilicity ratio is lower than the basic β -adrenoceptor antagonists, referred to above, due to the absence of the ion pair interaction. In general most of the dihydropyridine calcium channel antagonists have a relatively short half-life (around 2-5 hours). This is due to the moderate volumes of distribution and high intrinsic clearance, and their close interrelationship with lipophilicity. Most compounds therefore require frequent inconvenient dosage regimens. Amlodipine incorporates a basic centre and the additional interactions with phospholipid (see above for beta-adrenoceptor antagonists) gives the compound a larger volume than expected for neutral dihydropyridines. The combination of this volume increase, plus its low unbound intrinsic clearance gives amlodipine a long half-life (around 40 hours). Again as with beta adrenoceptor antagonists understanding the pharmacokinetic concepts can help to produce a differentiated compound with dramatic advantages. The differentiation is achieved by optimisation of a single property without changing other properties unfavourably.

Design and Distribution to the target

The distribution of drugs to their target has traditionally looked at the total tissue concentration as a measure of a drug reaching a receptor. For instance whole brain concentrations are often cited and compared for CNS drugs. Whole brain partitioning actually represents partitioning into the lipid of the brain, and not actually access to drug receptors. For instance, imipramine and desipramine partition into brain and are distributed unevenly [7]. The distribution corresponds to lipid content of the brain regions and not to specific desipramine binding sites. For seven-trans-membrane spanning receptors like the target(s) for imipramine and desipramine it is the drug in extra-cerebral fluid ECF which is in equilibria with the receptor. Cerebro-spinal fluid (CSF) concentrations can be taken as a reasonable guide of ECF concentrations. The apparent dramatic differences in brain/blood partitioning described for total brain (1000 fold or greater) collapse to a small ratio when free (unbound) concentration of drug in plasma is compared to CSF concentration rather than inert non-specific partitioning. Free unbound drug partitioning actually reflects the drug reaching the receptor and pharmacological effect. Unless active transport systems are invoked the maximum CSF to plasma partition coefficient is 1. The minimum partitioning appears to be 0.1, thus giving only a 10 fold range in CNS partitioning. The CNS in terms of the blood brain barrier is thus more permeable than imagined, allowing drugs like sulpiride (Log D - 1.2) to be used for CNS applications (8). Low lipophilicity compounds have several advantages, including low $Cl(u)$, renal clearance, lack of active metabolites, a low potential for pharmacokinetic drug interactions and non-specific cytotoxic and membrane-ion channel effects. Although potency can be an issue with hydrophilic compounds and therefore dose size, selectivity is not.

Selected References

1. Spahn Langguth, H.; Baktir, G.; Radschuweit, A.; Okyar, A.; Terhaag, B.; Ader, P.; Hanafy, A.; Langguth, P. (1998) P-glycoprotein transporters and the gastrointestinal tract. Evaluation of the potential in vivo relevance of in vitro data employing talinolol as model compound. *Int. J. Clin. Pharmacol. Ther.* 36, 16-24.
2. Smith, D.A. and van de Waterbeemd. (1999) Pharmacokinetics and metabolism in early drug discovery. *Curr. Opinons Chem. Biol.* 3, 373-378
3. Yamada, Y., Ito, K, Nakamura, K., Sawada, Y., Iga, T. (1993) Prediction of therapeutic doses of beta-adrenergic receptor blocking agents based on quantitative structure-pharmacokinetic/pharmacodynamic relationship. *Biol. Pharm. Bull* 16, 1251-9.
4. Smith, D.A., Ackland, M.J. and Jones, B.C. (1997) Properties of cytochrome P-450 and their substrates Part 1: active site characteristics. *Drug Discovery Today*, 2, 406-414.
5. Smith, D.A., Ackland, M.J., Jones, B.C. (1997) Properties of cytochrome P-450 and their substrates Part 2: properties of cytochrome P450 substrates. *Drug Discovery Today*, 2, 479-486
6. Smith, Dennis A.; Jones, Barry. (1999) Variability in drug response as a factor in drug design. *Curr. Opin. Drug Discovery Dev.* 2, 33-41
7. . Daniel, W., Danek, L., Janczar, L., Nocon, H. Melzacka, Mirosława. (1991) Regional distribution of imipramine, desipramine and specific [3H]desipramine binding sites in the rat brain after acute and chronic treatment with imipramine. *J. Pharm. Pharmacol.* 43, 31-5
8. Alfredsson, G.; Bjerkenstedt, L.; Edman, G.; Haernryd, C.; Oxenstierna, G.; Sedvall, G.; Wiesel, F. A. (1984) Relationships between drug concentrations in serum and CSF, clinical effects and monoaminergic variables in schizophrenic patients treated with sulpiride or chlorpromazine. *Acta Psychiatr. Scand.*, Suppl.311, 49-74.