松果腺剔除ノ甲状腺發育ニ及ボス影響ニ就テ

日本大學醫學科解剖學教室

秋 山 成 六 述

松果腺ガ有用ナル内分泌腺タルコトハ、明ナル所ナリ。而シテ松果腺ト、他ノ内分泌腺トノ、相互關係ニ至リテハ、不明ノ點多シ。著者ハ今回松果腺剔除ガ、何等カ甲状腺ニ影響ヲ及ボスモノニ非ザルヤトノ懸念ノモトニ、松果腺剔除白鼠ニ就テ、甲状腺ノ重量計測及ビ組織學的檢索ヲ行ヘリ。

松果腺剔除鼠ハ、伊澤ガ既ニ報告セル所ニシテ、剔出手術及ビ重量計測等ハ、該論文ヲ參考スベシ。鼠ハ二十日ニシテ手術シ、八十五日ニ殺シタリ、故ニ生存期間ハ、六十五日ナリトス。

第一表ハ手術及ビ對照動物ノ甲状腺ノ重量ヲ示シタルモノナルガ、平均重量ニ於テハ、手術動物ノ甲状腺ハ、對照動物ノ其レヨリモ重シ。然レドモ松果腺剔除ハ、身長及ビ體重ノ増大ヲ來スモノニシテ、大ナル白鼠ハ小ナル白鼠ヨリ、大ナル甲状腺ヲ有スベキノ理ナリ。(ドナルドソン氏表) 眞ニ増大シタルモノナリヤ、否ヤニ就テハ、此ヲ統計的ニ計算セザル可ラズ。各平均ノ差ガ少クトモ、算術平均ノ蓋然誤差ノ二乗ノ和ノ平方根ノ三倍ヨリ大ナラザル可ラズ。然ルニ今此表ヲ檢スルニ、P. D. M. (蓋然差異) ノ三倍ハ、平均重量ノ差ヨリ大ナリ。即チ統計的ニ肥大ナリト云フ能ハズ。換言スレバ、甲状腺肥大(松果腺剔除ニヨル)ハ來スモノニ非ズ。

第一表 甲状腺重量ノ統計的觀察

	♂		♀	
	手術	對照	手術	對照
算術平均	0.0121 _瓦	0.0120 _瓦	0.0137 _瓦	0.0123 _瓦
標準誤差	0.0028	0.0024	0.0015	0.0008
算術平均ノ蓋然誤差	0.0005	0.0004	0.0032	0.0001
變異係數	23.3	20.0	11.0	7.0

第二表 濾胞ノ平均直径及上皮細胞ノ高サ

	手術	對照
大濾胞	106.3 ^u	109.0 ^u
中等大濾胞	60.8	52.9
上皮細胞ノ高サ	9.2	9.5

組織的所見

次ニ是等ノ材料ニ就キ、定規ノ方法ノモトニ、パラフィン切片ヲ作製シ、ヘマトキシリン、エオジン染色、組織像ヲ檢スルニ左ノ如シ。

一、大ナル及ビ中等大濾胞ノ直径、並ビニ、腺上皮ノ高サヲ各二十ノ切片ニツキ計測シ、其ノ平均値ヲ求メシモ、手術動物及ビ對照ノ間ニ、格段ノ差ヲ認メズ。(第二表)

二、手術動物ノ甲状腺上皮細胞ノ濾胞内ニ、剝離脫出セルモノ、或ハ濾胞腔内、空虚ニシテ、コロイド物質ヲ缺グモノアルモ、是等ノ變化ハ、對照ニモ見ルコトヲ得タリ。

三、一般ニコロイド物質ハ手術動物ノ甲状腺ニ於テ、

エオジンニ着色セラル程度對照ニ比シテ、強キ傾向ヲ示ス。

長與ハ、七歳ノ兒童松果腺腫瘍ニテ死亡セルヲ、剖見シ、甲状腺濾胞ノ腫大シ、コロイド物質ヲ含有スルコト多カリシト。然レドモ、一次的乃至二次的ナルニモセヨ、毎常松果腺腫瘍ニテ、是等ノ變化ヲ招來シ得ルモノニヤ、尙多

少ノ歟義ヲ存スルモノアリ。余ノ寡聞未他ニ是ニ關スル文献ヲ見出ス能ハズ。

之ヲ要スルニ、白鼠ニ於テハ、松果腺分泌缺損ハ、甲状腺發育ヲ促進セシムルモノニモ非ズシテ、猶ホ顯著ナル甲状腺ノ組織的變化ヲ、惹起セシムルモノニモ非ズ。

出 版 文 献

1) Izawa, Y., 1916

On some anatomical change which follow removal of the pineal body from both sexes of the immature albino rat. *Ann. Jour. Physiol.* Vol. LXXVII, No. 1, P. 126 — 139.

2) 長興又郎, 1919

松果腺ノ病理 神經學雜誌, 第18卷, 第1號.

STUDIES ON THE PINEAL BODY. THE EFFECTS OF PINEALECTOMY ON THE GROWTH OF THYROID GLAND AND SOME OBSERVATION ON ITS HISTOLOGY

By

1 Dr. S. Akiyama

(The Department of Anatomy, School of Medicine of Nippon University.)

This report is concerned with the studies of a lack of pineal body on the growth of the thyroid of the same rats of which Izawa ('26) had already reported.

Izawa subjected these animals to pinealectomy when they were 20 days old and allowed them to grow until 95 days of age. Then they were killed and usual measurement was taken.

The table I, shows the statistical data for the fresh weight of the thyroid gland of experimental and control groups.

A study of this table reveals the following facts:

The mean weight of the thyroid of the operated rats were slightly heavier than that of the control. As already reported by Izawa ('26) pinealectomized rats were heavier than controls. Since the operated rats are larger they would be expected to have heavier absolute weight of organs than those of the controls, and such is really the case. It should be noted, however, that everything depends upon the statistical facts. Determination of this the probable difference of the mean of each organ in controled and tested of both sexes was found by the aid of the formula $P. D. M. = \sqrt{E^2 + E'^2}$.

A difference between the means should be equal to at least three times the probable difference of the mean is necessary to make the observed difference significant. The data for the thyroid in table I, do not stand in this tests, i. e. the observed differences are less than three times the probable difference of the mean. These deviation are, therefore, without significant here.

Table I.
Statistical data of the thyroid body.

	Males		Females	
	Tests	Controls	Tests	Controls
	gram	gram	gram	gram
Mean.....	0.0121	0.0120	0.0137	0.0123
S. D.	0.0028	0.0024	0.0015	0.0008
P. E. M.....	0.0005	0.0004	0.0032	0.0001
C. V.	23.3	20.0	11.0	7.0

S. D.Standart deviation.

P. E. M.Probable error of mean.

C. V.Coefficient of variability.

Histological examination

Materials were fixed in formalin and embedded in paraffin following after the process of general method. Sections were cut in 5 micron thickness and stained with haematoxylin and eosin.

In order to know the histological differences between the thyroid of the operated and controls the following method were employed.

First, the diameter of the follicles was measured. Follicles of large size and follicles of medium size were measured in twenty sections under the microscope and the average was taken. Second, the height of the epithelium was measured in the same sections. The data were given in table II. In this table we find that the differences of the thyroid of tests and controls are not significant.

Table II.
Mean diameter of the follicle and height of the epithelium

	Tests	Controls
	μ	μ
Follicle of large size.....	106.3	109.0
Follicle of medium size.....	60.8	52.9
Height of the epithelium.....	9.2	9.5

In sections of thyroid of both groups there were a few desquamate epithelial cells in follicles and at the same time some of the follicles were found empty. Generally speaking, colloidal substance of the thyroid was coloured deeper with eosin in the pinealectomized rat.

Nagayo ('19) found that there is certain structural change in thyroid in a case of child who died of pineal tumor. He noticed that the follicles were distended, and contained much colloidal substance.

Whether the structural change quoted above though it is a single case found by Nagayo is the consequence of pineal tumor or not is somewhat problematical at our present knowledge and I have so far failed to find any other literature referring to it.

Summary

In the albino rats

1. The growth of the thyroid was not stimulated by the lack of pineal secretion.
2. Pinealectomy did not produce any significant histological change in the thyroid gland.

Literature cited

Izawa, Y., 1926

On some anatomical change which follow removal of the pineal body from both sexes of the immature albino rat.

Am. Jour. Physiol. Vol. LXXVII. No. I. p. 126-139.

Nagayo, M., 1919

Zur Pathologie der Zirbeldrüse

Neurologia. Bd. XVIII. Heft. I. (in Japanese).