Experimental Study on the Mechanism of Hypoglycemic Action of sulfonylurea

By

**Assist.-Professor Hideo YOSHIDA M.D., Sadami YOKOO M.D.
Osamu AOCHI and Tomio UEHIRA**

Medical Department of Kyoto Prefectural Medical School (Director: Professor S. Tateishi M.D.)

By the discovery of insulin¹⁾ in 1922, the therapy of diabetes mellitus was improved very much and the mortality of diabetes mellitus is very few. But we have no view of complete recovering. From this long course per oral medicine was expected very much.

In 1942 Janbon²⁾ discovered the hypoglycemic action of IPTD and now there is no one who suspects the hypoglycemic effect of sulfonylurea. However, about the mechanism of hypoglycemic action of this medicine, there are many different opinions in the researchers throughout the world.

In this congress of Japanese Society of Internal Medicine one of us Yoshida take part in simposium of therapy of diabetes mellitus. So we study on the mechanism of hypoglycemic action of sulfonylurea. We have the following results.

Method

Many rabbits and rats were used. We used Wahrburg's pressure measure method and incubate in 38°C 1 hour long. We used buffer solution pH 7.4 with 0.9% NaCl solution and 0.1 Mol potassium-phosphate dibasic and we added glucose powder in this buffer solution to 150~200 mg/dl. Glucose was measured with Hagedorn-Jensen method.

On the histological study we used pironin-methylgreen, Gomori and Schiff-PAS method besides haematoxylin-eosin and ultra violet microscope (2650 Å) and electronmicroscope.

Results

A) The effect of sulfonylurea on the blood sugar

1) The blood sugar of normal rabbits, which were given BZ55 pro Kg. 0.25g, first a little ascends, next descends and after a long time comes back to original value. (Fig. 1) There are some individual variations of this ascendance and descendance on the time or the value. Many of them ascend after about 15 minutes, and after 30 minutes they show the highest value, namely the original value 80 mg/dl becomes 100 mg/dl. After an hour many of them come back to the original value. After about 3 hours from the beginning the blood sugar descends and from 3 hours to 5 hours we see the lowest value about 50 mg/dl. Then the blood sugar ascends gradually and after 24 hours it shows generally no influence. There is clear difference between natural descendance of normal rabbits and the curve of the rabbits which were given BZ 55.

Fig. 1. The effect of sulfonylurea on the blood sugar

2) When sulfonylurea is given to alloxan rabbits, the blood sugar, first a little ascends, next descends and after a long time comes back to original value as the normal rabbits. But the descendece of the blood sugar might be later than normal. When first the blood sugar shows 150 mg/dl it ascends to 180 mg/dl after about an hour. After 3 hours it descends to 150 mg/dl. Then the blood sugar of alloxan rabbits might descend to nearly normal value. This time too, clear difference is re-

cognised from natural descendece of alloxan rabbits.

When the blood sugar of alloxan rabbits already descends to normal value, they show about the same wave as normal rabbits.

3) While, when every 10 days BZ 55 or D 860 per os 0.05~0.2 g is given, or every 30 days BZ 55 subcutan 0.1 g is given, blood sugar is about normal value after 24 hours of latest medicine.

4) The effect of the sulfonylurea on the blood sugar, when 3g at a time are given to normal person.

At 9.00 a. m. blood sugar is measured in hunger condition and immediately BZ 55 3g are given. After 15 minutes, 30 minutes, 1 hour and 3 hours blood sugar is measured. Next day we measure the same course without medicine as control. On normal persons in hunger condition there is no difference between 9. a. m. and 12 a. m. After 3 hours of BZ 55 blood sugar increases to about 80% (65.9~91.4%) of it at 9 a. m.

But at the time of BZ 55 giving, it becomes about 65% (49.5~76.0%). But at this time we can see rarely a person, who shows no difference with or without medicine. (see Tab. 1, 2)

B) The effect of sulfonylurea on the carbohydrate metabolism in vitro.

1) Carbohydrate metabolism of rabbit which is given sulfonylurea. Carbohydrate metabolism of those rabbits which are given BZ 55 pro Kg. 0.25 g is measured. (see Tab. 3)

On the studium of ascendece of blood sugar, output of liver glucose increases and on the studium of maximum blood sugar or descendece of blood sugar output of liver glucose decreases. On the studium of descendece of blood sugar uptake of diaphragm glucose inclines to increase.

2) The effect of serum of rabbits, which are given sulfonylurea, on output of liver glucose or glucose uptake of diaphragm of normal rats. We measured carbohydrate metabolism of normal rats in the medium of glucose-buffer 1.5 cc and serum 0.5 cc of rats, which are given BZ 55 pro Kg. 0.25 g. Serum of ascendece studium of blood sugar in-

Ph. 1

Ph. 2

Ph. 3

Ph. 4

Ph. 5

Ph. 6

Ph. 7

Ph. 8

Ph. 9

Ph.10

Ph. 11

Ph. 12

Ph. 13

Ph. 14

Ph. 15

Ph. 16

Ph. 17

Ph. 18

Table 1. Blood sugar of normal persons giving BZ 55 3g.

Normal	Medicine	first	15 min.	30 min.	1 hour	3 hours	3 hours first	minimum first
1	BZ 55 3g	91	75	57	67	63	69.2%	62.1%
2	"	75	76	62	57	61	81.3%	76.0%
3	"	76	73	62	66	65	85.5%	81.6%
4	"	69	69	64	59	64	92.7%	85.5%
5	"	112	87	74	77	69	61.6%	61.6%
6	"	95	83	92	95	92	96.8%	96.8%
7	"	81	80	81	73	74	91.3%	90.1%
8	"	79	70	46	54	75	94.9%	58.2%
9	"	83	83	76	78	72	86.7%	86.7%
Medium	BZ 55 3g	85	78	69	70	71	84.4%	77.6%
1	0	74	71	71	78	73	98.6%	95.9%
2	0	57	63	63	63	64	112.2%	110.5%
3	0	68	72	72	68	68	100.0%	100.0%
4	0	70	65	73	70	73	104.2%	92.8%
5	0	90	87	76	87	90	100.0%	84.4%
6	0	87	83	90	90	95	109.1%	95.4%
7	0	44	47	49	53	49	111.3%	106.8%
8	0	61	57	56	61	66	108.2%	91.8%
9	0	76	78	74	81	74	97.4%	97.4%
Medium	0	70	69	69	72	72	104.5%	97.2%

creases the glucose output of liver of other animals. Serum of descendance studium increases the glucose uptake of diaphragm. Each method can be thought a quantitative analysis of glucagon^{3) 4) 5) 6)} or insulin. (see Tab. 4)

3) The effect of sulfonylurea in vitro on the glucose uptake of diaphragm.

When glucose uptake of diaphragm is measured, if insulin in various concentration is added, glucose uptake of diaphragm rises including insulin concentration.⁷⁾ (Fig. 2) When glucose uptake of diaphragm is 17 mg/dry weight 1 g without insulin, this uptake will be 40 mg/dry weight 1 g with 1 u/cc insulin.

But the glucose uptake of diaphragm does not rise with sulfonylurea in vitro. (see Tab. 5)

4) The influence of sulfonylurea in vitro on the effect of insulin on the carbohydrate metabolism of diaphragm.

Fig. 2. The effect of insulin on the uptake of diaphragm

Table 2. Blood sugar of diabetics giving BZ 55 3g.

Diabetics	Medicine	first	15 min.	30 min.	1 hour	3 hours	$\frac{3 \text{ hours}}{\text{first}}$
1	BZ 55 3g	188	178	172	172	100	53.2%
2	"	265	269	261	253	216	81.5%
3	"	116	111	111	94	78	67.2%
4	"	354	345	353	331	270	76.2%
5	"	322	312	302	284	234	72.6%
6	"	150	150	150	146	107	71.3%
7	"	137	133	84	77	68	49.6%
8	"	368	367	353	326	209	56.8%
9	"	141	134	115	96	60	42.6%
Medium	BZ 55 3g	227	222	211	198	149	63.4%
1	0	146	142	142	135	124	65.9%
2	0	266	262	257	251	214	80.4%
3	0	82	75	77	82	75	91.4%
4	0	379	465	441	434	334	88.1%
5	0	336	338	336	322	273	81.2%
6	0	149	142	136	136	117	78.5%
7	0	129	126	124	98	89	69.0%
8	0	262	250	239	234	198	75.5%
9	0	83	79	74	83	81	97.6%
Medium	0	204	209	203	197	167	80.8%

When glucose uptake of diaphragm is measured, insulin 0.1 u/cc, BZ 55 1 mg/cc and both were added in vitro. The effect of insulin on the glucose up take of diaphragm does not rise by the addition of sulfonylurea in vitro. (see Tab. 6)

C) The influence of sulfonylurea on the liver glycogen of rabbits.

With Schiff-PAS method we observed liver glycogen of rabbits, which were given per os BZ 55 pro Kg. 0.25 g, at 15 minutes, 30 minutes, 1.5 hours and 3 hours after giving.

On the studium of ascendance of blood sugar liver glycogen decreases and there is almost no liver glycogen on the studium of hyperglycemy.

On the studium of descendance of blood sugar liver glycogen increases gradually. This decrease of liver glycogen begins on periphery of liver acinus and the increase begins on central part. (see Ph. 1. normal, liverglycogen much; Ph. 2. after 15m., decrease; Ph. 3. after 30m., about no glycogen; Ph. 4. after. 1.5h., a little recover; Ph. 5. after 3h., almost recover)

D) The influence of sulfonylurea on Langerhans islet of rabbits.

We observed histology of Langerhans islet of rabbits, which were given sulfonylurea per os pro Ng. 0.25 g. Already from 15 minutes after giving medicine, stimulated status of beta cells of Langerhans islet was observed. This stimulated status is most distinct after

Table 3. Carbohydrate metabolism of rabbits which are given sulfonylurea
(BZ 55 pro Kg 0.25g) (mg/dry weight 1g.)

No.	Time (min.) from medicate to kill	Blood sugar (mg/dl)	Liver	Kidney	Spleen	Lung	Brain	Muscle
8	15	0min. 15min. 85 93	+85.47	-21.70		-25.43	-89.86	- 7.69
5	15	0min. 15min. 71 110	+47.53	-79.31	-88.39	-10.71	-87.37	- 27.32
13	20	0min. 20min. 82 91	+58.90	- 5.21				+ 17.98
12	20	0min. 20min. 83 102	+40.37	-49.76			-95.23	+ 8.56
6	30	0min. 30min. 63 79	+32.65	-32.96	-39.47		-79.91	- 13.61
2	30	0min. 20min. 83 105	+ 4.77	-29.02	-14.81	-12.84	-78.10	- 2.83
3	40	0min. 40min. 85 70	+20.63	-55.55	-45.21	-88.88	-149.10	- 9.13
4	60	0min. 60min. 87 61	+31.13	-20.68		-58.82	-113.30	-111.11
11	60	0min. 15min. 30min. 45min. 60min. 92 83 87 77 84	+24.45	0			-27.47	+ 13.64
9	60	0min. 45min. 79 118	-35.71	+ 3.81	+ 8.10	+32.25	-38.40	+ 6.48
7	80	0min. 80min. 89 80	0	-46.03	-14.87	0	-90.62	+ 12.48
10	90	0min. 40min. 60min. 90min. 87 95 100 95	+18.18	-43.82		-21.35	-72.57	- 18.00
1	180	0min. 20min. 30min. 60min. 90min. 120min. 180min. 87 105 98 94 93 87 85	+12.40	-57.97	-50.73	-56.41	-82.85	- 10.44
Control (6 cases)			+42.84	-67.03	-38.32	-36.05	-85.79	- 14.43

Table 4. The effect of serum of rabbit, which is given sulfonylurea on the carbohydrate metabolism of rats.

	No.	no serum	with normal serum				with serum at 15min after medicate			with serum at 1.5hours after medicate		
uptake of diaphragm	1	-40.42	-45.57				-38.76			-71.35		
	2	-11.47	-16.22				-28.55			-18.69		
	3	-50.08	-58.15				-52.07			-63.52		
output of Liver	1	+17.00	+14.68				+23.29			+21.80		
	2	+ 4.52	+ 6.52				+35.39			+ 1.78		
	3	+67.88	+57.42				+81.13			+55.55		
	No.	0	15'	30'	60'	90'	2°	3°	4°	5°	7°	(time)
Blood sugar (mg/dl)	1	103	131	126	128	121						
	2	77	88	120	113	100						
	3	81	98	90	125	86	83	92	77	75	73	

Table 5. The effect of insulin on the diaphragm.

		uptake of glucose (mg/lg)				tissue respiration (QO ₂)			
Insulin No.		0	u/10 ²	u/10	1u	0	u/10 ²	u/10	1u
1		-23.96		-24.42	-23.00	6.34		6.53	5.74
4		-27.66	-31.69		-69.80	5.25	6.07		8.11
5		- 5.15	-15.02		-31.93	4.95	5.66		6.32
9		-20.95	-21.51	-35.32	-22.74	4.99	6.03	6.10	4.64
10		- 8.98		-10.35		4.77		5.08	
11		-14.97		-19.88		6.19		5.74	
12		-14.02		-27.20		6.12		6.33	
13		-28.82		-36.69		7.25		6.35	
14		-15.00		-30.03		4.18		6.03	
15		-19.55		-45.71		6.20		8.44	
18		-19.77	-25.96	-35.43	-38.78	5.78	6.55	6.79	6.17
19		-15.89	-29.85	-35.02	-40.98	6.91	6.94	7.08	6.93
20		-17.00	-17.60	-32.90	-53.23	7.64	8.47	7.97	8.81
Medium		-17.82	-23.60	-30.26	-40.06	5.89	6.62	6.60	6.67

1.5 or 3 hours. These changes are observed clearly with Gomori, pyronin-methylgreen staining method, ultraviolet microscope and electron microscope.

1) Gomori staining (CHP method)

At 15 minutes after giving medicine partly, at 30 minutes after giving in almost all Langerhans islets, escape of granule of beta cells⁸⁾ and also enlargement of nucleus of beta cells are observed. These changes are observed most clearly at 3 hours after giving medicine and are observed still after 6 hours. (Ph. 6. normal; Ph. 7. after 3 hours)

Table 6. The effect of sulfonylurea (1mg/cc) in vitro on the tissue of rats.

		on the carbohydrate metabolism						on the tissue respiration					
Organ		Diaphragm			Liver			Diaphragm			Liver		
Medicine	No.	0	BZ 55	D 860	0	BZ 55	D 860	0	EZ 55	D 860	0	BZ 55	D 860
	2				109.74	108.14					9.62	10.47	
	3				120.52	95.47					12.30	9.67	
	6	-18.16	-21.13	-25.28	50.38	45.91	22.76	4.55	5.60	4.97	9.29	11.48	9.40
	7	-10.06	-11.46	-17.04	60.12	93.61	142.13	6.13	5.34	6.15	9.50	10.27	11.33
	8	-22.48	-10.84	-11.57	84.38	108.46	95.42	7.08	7.35	7.44	10.88	12.68	10.88
	10	-8.98	0		81.49	104.60		4.77	5.24		10.37	9.33	
	11	-14.97	0		105.79	142.54		6.19	5.87		11.06	10.87	
	12	-14.02	-14.33		72.22	108.64		6.12	6.09		9.03	9.03	
	13	-28.82	-27.21		96.29	84.61		7.25	5.88		11.73	10.56	
	14	-15.00	-39.44		57.85	57.85		4.18	5.07		12.31	10.12	
	15	-19.55	-19.35		72.23	57.75		6.20	6.21		10.40	9.67	

Table 7. The influence of sulfonylurea (mg/cc) in vitro on the effect of insulin on the carbohydrate metabolism of diaphragm of rats.

		on the carbohydrate metabolism				on the tissue respiration			
Medicine	No.	0	u/10 Insulin	BZ 55	u/10 Ins. + BZ55	0	u/10 Insulin	BZ 55	u/10 Ins. + BZ55
	10	-8.98	-10.35	0	0	4.77	5.08	5.24	5.72
	11	-14.97	-19.88	0	-17.35	6.19	5.74	5.87	6.32
	12	-14.02	-27.20	-14.33	-19.37	6.12	6.33	6.09	6.25
	13	-28.82	-36.69	-27.21	-39.54	7.25	6.35	5.88	6.91
	14	-15.00	-30.03	-39.44	-23.63	4.18	6.03	5.07	5.01
	15	-19.55	-45.71	-19.35	-20.93	6.20	8.44	6.21	7.44

2) Pironin-methylgreen staining.

The more intensive staining of nucleus of beta cells with methylgreen and of protoplasm with pyronin are observed from 15~30 minutes and clearly at 1.5 hours and less at 3~6 hours. (Ph. 8. normal; Ph. 9. after 15 m.; Ph. 10. after 1.5 hours)

3) Observation by ultra-violet microscope. ($\times 1200$)

Bausch-Lomb optical Company's ultra-violet microscope was used with 2650 Å to observe nucleic acids absorption zone. Spectrum purity of this microscope is ± 30 Å.

There was remarkable reduction of nucleic acids in Langerhans' islets than that of exocrine parts and two different types of islet cells were observed. By examining the same cells in a section stained with Gomori's CHP method, their identity was established as alpha and beta cells. The nuclei of the alpha cells were dark, and those of the beta cells were pale.

By the administration of BZ 55, after 15~30 minutes, increase inclination of ultra-violet absorption was found at beta cells' nuclei and cytoplasm. After 1~1.5 hours absorption got maximum, and a little decreased after 3~6 hours.

The same inclination was also observed by the continued administration of BZ 55. (Ph. 11. normal; Ph. 12. after 15m.; Ph. 13. after 1.5 h.; Ph. 14. continued administ)

4) Observation by electron microscope.

Fixation and embedding : general method.

Microtome : Sjöstrand's thermo-expansion type

Electron microscope : Akashi's TRS-50 B type

a) On the normal rabbits

The alpha cells have a pale background cytoplasm, which contains numerous, dense, round granules. The pale nuclei are usually ovoid. One nucleolus is seen distinctly in a nucleus. (See Ph. 15×15000)

The beta cells are darker and contain granules which are less dense and smaller than those of the alpha cells. The darker appearance of the beta cells is apparently due to an abundance of ergastoplasm in their cytoplasm. The nuclei are dark and are ovoid in shape. The other components of the cytoplasm are small, rod shaped mitochondria and a prominent Golgi complex and a fine, fibrillar substance adjacent to the nucleus. (See Ph. 16×21000)

b) On the rabbits administrated sulfonylurea BZ 55 (after 15 minutes).

A remarkable change is not observed in alpha cells.

In beta cells the cytoplasm and the nuclei swell. The chromatine congregates now and then and becomes to some gross mass in swollen nuclei. The beta granules decrease distinctly and are found a few at periphery of the cytoplasm. A remarkable change is not observed on the mitochondria and the other components of the cytoplasm. (See Ph. 18×9000) There is such a change in individual cells in various degrees on the same section. (See Ph. 17×4500)

We appreciated very much for Dr. K. Kimura's cooperation on this work.

Discussion

From our results we recognized 1)~6) about rabbits giving sulfonylurea.

1) The blood sugar first a little ascends, next descends and after a long time comes back to original value.

2) The descendance of the blood sugar of alloxan rabbits might be late.⁹⁾¹⁰⁾¹¹⁾¹²⁾

3) On the studium of ascendance of blood sugar output of liver glucose increases and on the studium of descendance of blood sugar glucose output of liver decreases and glucose uptake of diaphragm increases.

4) Serum of ascendance studium of blood sugar increases the glucose output of liver of other animals. Serum of descendance studium of blood sugar increases the glucose uptake of diaphragm of other animals.

5) On the studium of ascendance of blood sugar liver glycogen decreases and there is almost no liver glycogen on the studium of hyperglycemy and on the studium of descendance of blood sugar liver glycogen increases gradually.¹³⁾¹⁴⁾¹⁵⁾¹⁶⁾¹⁷⁾

6) Already from 15 minutes after giving medicine, stimulated status of beta cells of Langerhans islet was observed. This stimulated status is most distinct after 1.5 or 3 hours. These changes are observed clearly with Gomori, pyronin-methylgreen staining method, ultraviolet ray microscope and electron microscope.

From our results we recognized 7)-8) by the addition of sulfonylurea in vitro.

7) The glucose uptake of diaphragm does not rise by the addition of sulfonylurea in vitro.¹⁸⁾¹⁹⁾²⁰⁾²¹⁾

8) The effect of insulin on the glucose uptake of diaphragm does not rise by the addition of sulfonylurea in vitro.

In addition to the above-mentioned results, we have the following reports :

9) There is no effect of sulfonylurea on the pancreatectomized dog.²²⁾⁻²⁷⁾²⁷⁾

10) By the big dose of sulfonylurea the necrosis of alpha cell of Langerhans islet was recognized.²⁸⁾⁻³⁴⁾

11) We can save insulin clinically and experimentally by sulfonylurea.

12) We can not think commonly, that an artificial substance as sulfonylurea is useful on many different points of therapy of diabetes mellitus as an angel on the battle against diabetes mellitus.

By above-mentioned 12 points we have come to the following belief.

Sulfonylurea work on the Langerhans islet and let the alpha cells having glucagon to put out and a rest to take³⁵⁾, and bring the hyper function of beta cells and so is useful on the therapy of diabetes mellitus.

If our belief is right, we think that the examination of islet function is possible from our result of A) 4). And on the result of A) 4) we can not see the studium of ascendance of blood sugar by a little dose.

Up to this time there are many inverted results of the researchers throughout the world on the various studies of animals administrated sulfonylurea. We think it becomes by no attention to the up and down curve of blood sugar that changes by dose of medicine.

We have no result and can say nothing about the influence of sulfonylurea on the destruction of insulin⁹⁾³⁶⁾.

Conclusion

we think that sulfonylurea are useful on the therapy of diabetes mellitus, by the working on Langerhans islet. And at the same time we advance the possibility of examination of islet function.

References

- 1) Banting, G. and Best, H. : J. Lab. Clin. Med. 1922, 464.
- 2) Janbon, M. et al : Montpellier Med. 21-22; 489, 1942.
- 3) Murlin, J. R. et al : J.B.C. 15; 365, 1913. ; 56 ; 253, 1923.
- 4) Bürger, M., Kramer, H. : Z. ges. exp. Med. 65 ; 487, 1929.
- 5) Bürger, M., Brandt, W. : Z. ges. exp. Med. 67; 441, 1929 ; 69 ; 57, 1930.
- 6) Sutherland, E.W. & Cori, C.F. : J.B.C. 172 ; 137, 1948.
- 7) Gemill C.L. : Bull. Johns Hopkins Hosp. 66 ; 232, 1940.
- 8) Loubatieres, A. : Compt rend. Soc. biol. 138 ; 766, 1944.
- 9) Mirsky, I.A. et al : Metabolism 5 ; 138 156, 1956.
- 10) Beringer, A. u Lindner, A. : Wien Med. Wschr. 68 ; 516, 1956.
- 11) Chen, K.K., Anderson, R.C. and Maze, N. : proc. Soc. Exp. Biol. & Med. 63 ; 483, 1946.
- 12) Baender, A. : Deut. Med. Wschr. 81 ; 889, 1956.
- 13) Tyberghein, J.M. et al : Proc. Soc. exp. Biol. & Med. 92 ; 322, 1956.
- 14) Lang, S. et al : Metabolism 5 ; 733, 1956.
- 15) Dulin, E.W. et al : Ann N.Y. Acad. Sci. 71 ; 177, 1957.
- 16) Beringer, A. u Keibl, E. : Wien. med. Wschr. 38-39; 792, 1956.
- 17) Miller, W. L. Jr. and Dulin W.E. : Science 123 ; 584, 1956.
- 18) Levine, R. : Diabetes 5 ; 375, 1956.
- 19) Levine, R. et al : Metabolism 5; 721, 1956.
- 20) Fritz, I.B. et al : Metabolism 5 ; 744, 1956.
- 21) Wick, A.N., Britton, B. and Grabowski, R. : Metabolism 5 ; 739, 1956.
- 22) Loubatieres, A. : Presse med. 63 ; 1701, 1728, 1955.
- 23) Sirek, A. : Canad. M.A.J. 74 ; 960, 1956.
- 24) Campbell, J. : Canad. M.A.J. 74 ; 962, 1956.
- 25) Schambye, P. : Diabetes 6 ; 146, 1957.
- 26) Loubatieres, A. et al : Ann. N.Y. Acad. Sci. 71 ; 192, 1957.
- 27) Houssay, B.A. et al : Ann. N.Y. Acad. Sci. 71 ; 12, 25, 1957.
- 28) Von Holt, C. et al : Arch. Exp. path. pharmac. 224 ; 78, 1955.
- 29) Ferner, H. : Deut. gesam. Pharmakol. 1955.
- 30) Bertram, F. et al : Deut. med. Wschr. 80 ; 1455, 1955.
- 31) Achelis, J.D. et al : Deut. med. Wschr. 80 ; 1452, 1955.
- 32) Yoshioka, H. : Naibunpi 3; 97, 1956.
- 33) Creutzfeldt, W. : Diabetes 6; 135, 1957.
- 34) Colwell, A.R. and Colwell, J.A. : Metabolism 5; 749, 1956.
- 35) Anderson, G.E. et al : Proc. Soc. exp. Biol. 92; 340, 1956.
- 36) Volk, B.W. et al : Ann. N.Y. Acad. Sci. 71; 141, 1957.
- 37) Yoshida, H. et al : will be published on Fol. Endocri. Jap.