REVIEW

The roles of kisspeptin and gonadotropin inhibitory hormone in stress-induced reproductive disorders

Takeshi Iwasa, Toshiya Matsuzaki, Kiyohito Yano, Yiliyasi Mayila and Minoru Irahara

Department of Obstetrics and Gynecology, Institute of Biomedical Sciences, Tokushima University Graduate School, Tokushima 770-8503, Japan

Abstract. Several kinds of stress suppress the hypothalamic-pituitary-gonadal (HPG) axis and reproductive behavior in humans and animals. These changes can eventually cause diseases and disorders, such as amenorrhea and infertility. In previous studies, it has been shown that stress-related factors, *e.g.*, corticotropin-releasing hormone, cortisol, and pro-inflammatory cytokines, promote the stress-induced suppression of the HPG axis. However, these mechanisms are not sufficient to explain how stress suppresses HPG axis activity, and it has been suggested that some other factors might also be involved. In the early 21st century, novel neuroendocrine peptides, kisspeptin and gonadotropin inhibitory hormone (GnIH)/RFamide-related peptide 3 (RFRP-3), which directly regulate GnRH/gonadotropin synthesis and secretion, were newly discovered. Growing evidence indicates that kisspeptin and GnIH/RFRP-3 play pivotal roles in the stress-induced disruption of the HPG axis and reproductive behavior in addition to their physiological functions. This review summarizes what is currently known about the roles of kisspeptin and GnIH/RFRP-3 in stress-induced reproductive disorders.

Key words: Kisspeptin, Gonadotropin inhibitory hormone, Stress, Gonadotropin

Introduction

Reproductive functions and stress regulation systems are closely linked in most species. Generally, reproductive functions are temporarily suppressed by various kinds of stress, such as infection, psychological burdens, excess exercise, and undernutrition, in humans and animals because reproduction, which requires a large amount of energy, is not essential for survival [1-4]. Such changes are considered to play important roles in the regulation of homeostasis; *i.e.*, in the appropriate allocation of finite energy to prioritized activities. Although reproductive functions are usually restored immediately after stress levels reduce [5, 6], prolonged or excessive stress can sometimes have negative long-term consequences [7].

Reproductive functions are mainly regulated by the hypothalamic-pituitary-gonadal (HPG) axis in humans

and animals. Physical and psychological stressors suppress HPG axis activity, mainly by inhibiting gonadotropin-releasing hormone (GnRH) activity in both males and females, which results in decreased gonadotropin levels [8-13]. Previous studies by our group and others have shown that alterations in the activity levels of some stress-related factors reduce GnRH and gonadotropin secretion and concomitantly promote the stress response [14-16]. However, these mechanisms are not sufficient to explain how stress suppresses HPG axis activity, and it has been suggested that some other factors may also be involved. A breakthrough occurred in the early 21st century, when novel neuroendocrine peptides that directly regulate GnRH/gonadotropin synthesis and secretion were discovered [17-19]. Kisspeptin, which is encoded by the *Kiss1* gene, is a hypothalamic peptide that directly stimulates GnRH synthesis and release [20, 21], whereas gonadotropin inhibitory hormone (GnIH)/RFamide-related peptide 3 (RFRP-3) acts within the hypothalamus and pituitary to suppress the release and synthesis of GnRH and gonadotropins [22-27]. Some studies have shown that these two factors are involved in stress-induced reproductive disorders as well as the regulation of physiological HPG axis activity. In this review,

Submitted Jan. 16, 2018; Accepted Jan. 16, 2018 as EJ18-0026
Released online in J-STAGE as advance publication Jan. 26, 2018
Correspondence to: Takeshi Iwasa, Department of Obstetrics and Gynecology, Institute of Biomedical Sciences, Tokushima University Graduate School, 3-18-15 Kuramoto-Cho, Tokushima 770-8503, Japan.
E-mail: iwasa.takeshi@tokushima-u.ac.jp

we discuss the neuroendocrine mechanisms responsible for stress-induced reproductive disorders, mainly focusing on the relationships between kisspeptin, GnIH/RFRP-3, and HPG axis dysfunction. We also discuss the relationship between GnIH/RFRP-3 and dysfunctional reproductive behavior.

The Roles of Stress-Related Factors in Stress-Induced HPG Axis Dysfunction

As noted above, alterations in the activity levels of some stress-related factors decrease GnRH and gonadotropin secretion and concomitantly promote the stress response (Fig. 1). For example, hypothalamic corticotropin-releasing hormone (CRH), which is the principal driving factor of the HPA axis during stress, is a potent inhibitor of GnRH secretion. In experimental animals, the central injection of CRH suppressed the luteinizing hormone (LH) pulse [28-30], which is an indicator of GnRH pulse activity, and various types of stress-induced LH suppression were reversed by the administration of CRH antagonists [30-34]. Similarly, in women the peripheral

infusion of CRH decreases gonadotropin secretion, and GnRH administration prevents this alteration, indicating that CRH inhibits GnRH secretion in humans [35, 36]. In addition, some studies have reported that cortisol, which is produced by the adrenal gland, suppresses GnRH secretion in humans and monkeys [37-40]; however, some other studies obtained contradictory results [9, 41]. Pro-inflammatory cytokines; *i.e.*, interleukin-1 β and tumor necrosis factor- α , also play pivotal roles in the stress-induced suppression of the HPG axis. The expression levels of these cytokines are upregulated by immune stress, and GnRH release and serum gonadotropin levels are decreased by their central administration [42-47]. Although the primary roles of these stress-related factors are to promote appropriate responses to exogenous and endogenous stressors, they also modulate HPG axis activity in order to save energy. The stress-related factors described above are promptly activated by stress, and they might play roles in the acute phases of stress-induced reproductive disorders. Whereas, it has been assumed that some other factors might also be involved in such disorders, especially in chronic or severe stress

Fig. 1 The roles of kisspeptin, gonadotropin-inhibitory hormone (GnIH)/RFamide-related peptide 3 (RFRP-3), and stress-related factors under physiological and stress conditions. Kisspeptin and GnIH/RFRP-3 participate in the control of physiological GnRH/luteinizing hormone (LH) regulation. The activity levels of kisspeptin and GnIH/RFRP-3 are increased and decreased, respectively, by high estrogen concentrations at the time of the GnRH/LH surge. These alterations promote both ovulation and reproductive behaviors, and consequently, increase the chance of pregnancy. On the contrary, kisspeptin, GnIH, and stress-related factors also play roles in stress-induced reproductive disorders. Kisspeptin and GnIH expression and activity are decreased and increased by stress, respectively, and these changes contribute to suppressing the activity of the hypothalamic-pituitary-gonadal axis. The upregulation of GnIH/RFRP-3 expression decreases reproductive behavior, and consequently, reduces the chances of pregnancy and causes infertility.

conditions. Growing evidence suggests that kisspeptin and GnIH/RFRP-3 might contribute to reproductive disorders in such adverse physiological states. Thus, the physiological and pathological roles of kisspeptin and GnIH/RFRP-3 are discussed in the following sections.

Physiological Roles of Kisspeptin and GnIH/RFRP-3

In 2003, two independent groups reported that a mutation in the G protein-coupled receptor (GPR54); *i.e.*, the kisspeptin receptor, induced hypogonadotropic hypogonadism and pubertal failure in humans and mice [18, 19]. Thus, kisspeptin was assumed to be a key factor for gonadotropin secretion and a modulator of HPG axis activity. Indeed, subsequent studies have shown that kisspeptin stimulates GnRH secretion *via* GPR54 on GnRH neurons [48-50]. The amino acid sequence of kisspeptin, especially the C-terminal 10-amino acid sequence, is well conserved in most mammals [51-57]. Hypothalamic kisspeptin neurons are mainly located in the anterior and posterior nuclei; *i.e.*, the anteroventral periventricular nucleus (AVPV)/preoptic area (POA) and arcuate nucleus (ARC) [55, 56, 58-64]. *Kiss1* mRNA expression in the AVPV/POA is increased in the afternoon during proestrus and is also upregulated by the administration of estrogen, whereas *Kiss1* mRNA expression in the ARC is decreased by the administration of estrogen [61, 63, 65-67]. These results indicate that kisspeptin plays pivotal roles in the regulation of the negative and positive feedback effects of estrogen and that it is indispensable for the maintenance of the ovulation cycle (Fig. 1).

In 2000, Tsutsui *et al.* discovered a novel neuropeptide that suppresses the release of gonadotropins from cultured avian pituitary cells [17]. As this neuropeptide was the first hypothalamic factor that was found to suppress gonadotropin release, it was named GnIH based on its biological activity. GnIH molecules were subsequently identified in other vertebrates, mammals, primates, and humans [68-70]. The mammalian GnIH orthologous gene and peptide are named *Rfrp* and RFRP-3, respectively, because the neuropeptide possesses the LPXRFamide (X = L or Q) motif in its C-terminus. Hypothalamic GnIH/RFRP-3 neurons are mainly located in the paraventricular nucleus in birds and the dorsomedial hypothalamic area in mammals [71]. GnIH/RFRP-3 neurons project to the median eminence in birds and female sheep, and suppress the synthesis and secretion of gonadotropins at the

pituitary level *via* the GnIH/RFRP-3 receptor GPR147 under both *in vivo* and *in vitro* conditions in male birds, female rats, and female sheep [17, 26, 69, 72]. In addition, GnIH/RFRP-3 neurons project to GnRH neurons in the hypothalamus and suppress their activity *via* GPR147 in mammals and birds [23, 26, 73]. GnIH/RFRP-3 neurons were activated by the injection of estradiol in ovariectomized female hamsters [23]. However, estradiol has no such effect and GnIH/RFRP-3 neurons are inactivated during the LH surge that occurs during the estrous stage [26, 74]. These results indicate that GnIH/RFRP-3, as well as kisspeptin, plays important roles in the regulation of the negative and positive feedback effects of estrogen, and hence, contributes to the maintenance of the ovulation cycle (Fig. 1).

The Roles of Kisspeptin in Stress-Induced HPG Axis Dysfunction

As noted above, kisspeptin plays pivotal roles in the regulation of HPG axis activity and the maintenance of a regular ovulation cycle in many species. Thus, it had been assumed that the disruption of kisspeptin activity would decrease HPG axis activity and cause reproductive disorders (Fig. 1). In 2008, we reported that immune stress induced by the administration of lipopolysaccharide (LPS) suppressed hypothalamic *Kiss1* mRNA expression and the serum LH level in female rats [75]. In latter study, we also showed that the LPS-induced LH suppression was completely reversed by the co-administration of kisspeptin. As far as we know, our study was the first to describe the effects of stress on the kisspeptin system and the gonadotropin secretion disorders they cause. Thereafter, we and other groups have vigorously evaluated the effects of stress on the kisspeptin system and the neuroendocrine mechanisms that underlie these effects. In agreement with the findings of our aforementioned study, the administration of LPS decreased hypothalamic *Kiss1* mRNA expression and kisspeptin immunoreactivity, and the co-administration of kisspeptin partially restored serum LH levels in female rats [76]. In addition, other kinds of stress, such as psychosocial, unpredictable chronic (six randomly assigned stressors), and hypoglycemic stress, also reduced hypothalamic *Kiss1* mRNA expression and kisspeptin neuron activity [77-79]. Some studies have also evaluated the intermediate factors that transfer stress signals to kisspeptin neurons. As a result, it was found that both the central administration of CRH and the peripheral administration of corticosterone

reduced hypothalamic *Kiss1* mRNA expression and kisspeptin neuron activity in female rats and mice, indicating that activation of the HPA axis is involved in the stress-induced suppression of the kisspeptin system [77, 80]. Interestingly, relatively severe stress protocols were used in these studies; *i.e.*, high-dose LPS (1–5 mg/kg) or repeated LPS administration protocols (three consecutive injections at 24-h intervals), to evaluate the effects of immune stress [75, 76, 81, 82], and chronic (once per day for four weeks) or combined (restraint and isolation) protocols were used to evaluate the effects of unpredictable and psychosocial stress [78, 79]. Similarly, we found that a subseptic dose (500 µg/kg) of LPS did not affect hypothalamic *Kiss1* mRNA expression in female rats, indicating that only chronic or severe stress affects the kisspeptin system [81, 82]. As kisspeptin is indispensable for the regulation of HPG axis activity and maintaining fecundity, it might be rigidly maintained even under stressful conditions. In other words, reproductive disorders caused by disruption of the kisspeptin system might be more serious than those induced by stress-related factors alone.

The Roles of GnIH/RFRP-3 in Stress-Induced HPG Axis Dysfunction

As described above, GnIH/RFRP-3 has suppressive effects on GnRH and gonadotropin activity in many species and plays roles in the regulation of HPG axis activity, especially during the GnRH/LH surge. Recently, it has been shown that some kinds of acute and chronic stress result in increases in the number of GnIH/RFRP-3-immunoreactive cells and *GnIH/Rfrp* mRNA expression in the hypothalamus. It has also been reported that these changes in GnIH/RFRP-3 expression disrupt HPG axis activity and suppress reproductive ability (Fig. 1). Acute (3 h) and chronic (14 days, 3 h/day) psychological (immobilization) stress lead to the upregulation of hypothalamic *GnIH/Rfrp* mRNA expression, and *GnIH/Rfrp* mRNA expression levels are negatively correlated with serum LH levels in male rats [83]. In addition, half of GnIH/RFRP-3 neurons express the glucocorticoid receptor (GR), and adrenalectomy abolishes the increase in *GnIH/Rfrp* mRNA expression seen under chronic psychological stress conditions [83]. Similarly, the administration of corticosterone increases *GnIH/Rfrp* mRNA expression levels in rHypoE23, an *Rfrp*-expressing cell line [84], and a GR antagonist blocked this effect of corticosterone [85, 86]. In addition, the administration of

cortisol increased *GnIH/Rfrp* mRNA expression and reduced GnRH mRNA and serum LH levels in fish [87]. These findings suggest that GnIH/RFRP-3 mediates the suppressive effects of glucocorticoids on the HPG axis under stress conditions. Interestingly, a recent study has shown that GnIH/RFRP-3 suppressed sexual maturation in socially non-dominant female rats living in colonies, indicating that GnIH/RFRP-3 is related to social stress-induced reproductive disorders [88]. As has been found for kisspeptin, relatively severe stress protocols are needed to affect the hypothalamic GnIH/RFRP-3 system; *e.g.*, high-dose LPS (2–5 mg/kg) was used to evaluate the effects of immune stress [81, 89], and a repeated immobilization protocol was used to evaluate the effects of psychological stress [83]. As GnIH/RFRP-3 is also essential for the regulation of HPG axis activity, GnIH/RFRP-3 expression might be rigidly maintained even under stressful conditions. We suggest that the neuroendocrine mechanisms responsible for stress-induced reproductive disorders might differ according to the severity of stress, and that GnIH/RFRP-3 and kisspeptin might only contribute to such disorders in severe physiological and/or psychological conditions.

The Roles of GnIH/RFRP-3 in Stress-Induced Reproductive Behavior Disorders

Recently, it has been shown that GnIH/RFRP-3 plays roles in the regulation of reproductive behavior in rodents (Fig. 1). The central administration of GnIH/RFRP-3 decreased sexual behavior in male rats [90], and it reduced sexual motivation in female hamsters [71, 91]. GnIH/RFRP-3 affects neuronal activity in some hypothalamic nuclei (the POA, medial amygdala, and the bed nucleus of the stria terminalis) that are related to female sexual behavior [91]. Thus, it has been speculated that stress-induced upregulation of GnIH/RFRP-3 expression not only disrupts the HPG axis, but also suppresses reproductive behavior and promotes infertility or subfertility. An excellent report about this topic was published in 2017 [92]. In the latter study, it was shown that chronic immobilization stress increased hypothalamic *GnIH/Rfrp* mRNA expression and reduced sexual behavior, the frequency of pregnancy, and litter size in female rats. It was also demonstrated that genetic silencing of *GnIH/Rfrp* with short hairpin RNA during stress led to the normalization of these parameters. These results indicate that GnIH/RFRP-3 is involved in the stress-induced disruption of reproductive behavior, as well as in HPG

axis disorders and that GnIH/RFRP-3 might be a useful clinical target for preventing stress-induced infertility.

Conclusion

Growing evidence indicates that kisspeptin and GnIH/RFRP-3, as well as stress-related factors, play pivotal roles in stress-induced reproductive disorders, such as

disruption of the HPG axis and reproductive behavior. Thus, kisspeptin and GnIH/RFRP-3 might be useful clinical targets for the treatment of stress-induced reproductive disorders, such as menstrual dysfunction and infertility. As the amount of human data available is limited, more evaluations are needed before kisspeptin or GnIH/RFRP-3-based treatments can be developed.

References

- Hart BL (1988) Biological basis of the behavior of sick animals. *Neurosci Biobehav Rev* 12: 123–137.
- Yirmiya R, Avitsur R, Donchin O, Cohen E (1995) Interleukin-1 inhibits sexual behavior in female but not in male-rats. *Brain Behav Immun* 9: 220–233.
- Bonneaud C, Mazuc J, Gonzalez G, Haussy C, Chastel O, *et al.* (2003) Assessing the cost of mounting an immune response. *Am Nat* 161: 367–379.
- Pauli SA, Berga SL (2010) Athletic amenorrhea: energy deficit or psychogenic challenge? *Ann N Y Acad Sci* 1205: 33–38.
- Perkins RB, Hall JE, Martin KA (2001) Aetiology, previous menstrual function and patterns of neuro-endocrine disturbance as prognostic indicators in hypothalamic amenorrhea. *Hum Reprod* 16: 2198–2205.
- Falsetti L, Gambera A, Barbetti L, Specchia C (2002) Long-term follow-up of functional hypothalamic amenorrhea and prognostic factors. *J Clin Endocrinol Metab* 87: 500–505.
- Jacoangeli F, Masala S, Staar Mezzasalma F, Fiori R, Martinetti A, *et al.* (2006) Amenorrhea after weight recover in anorexia nervosa: role of body composition and endocrine abnormalities. *Eat Weight Disord* 11: e20–e26.
- Rivier C, Rivest S (1991) Effect of stress on the activity of the hypothalamic-pituitary-gonadal axis: peripheral and central mechanisms. *Biol Reprod* 45: 523–532.
- Matsuwaki T, Kayasuga Y, Yamanouchi K, Nishihara M (2006) Maintenance of gonadotropin secretion by glucocorticoids under stress conditions through the inhibition of prostaglandin synthesis in the brain. *Endocrinology* 147: 1087–1093.
- Du Ruisseau P, Tache Y, Brazeau P, Collu R (1979) Effects of chronic immobilization stress on pituitary hormone secretion, on hypothalamic factor levels, and on pituitary responsiveness to LHRH and TRH in female rats. *Neuroendocrinology* 29: 90–99.
- Gonzalez-Quijano MI, Ariznavarreta C, Martin AI, Treguerres JA, Lopez-Calderon A (1991) Naltrexone does not reverse the inhibitory effect of chronic restraint on gonadotropin secretion in the intact male rat. *Neuroendocrinology* 54: 447–453.
- Santoro N, Filicori M, Crowley WF Jr (1986) Hypogonadotropic disorders in men and women: diagnosis and therapy with pulsatile gonadotropin-releasing hormone. *Endocr Rev* 7: 11–23.
- Reame NE, Sauder SE, Case GD, Kelch RP, Marshall JC (1985) Pulsatile gonadotropin secretion in women with hypothalamic amenorrhea: evidence that reduced frequency of gonadotropin-releasing hormone secretion is the mechanism of persistent anovulation. *J Clin Endocrinol Metab* 61: 851–858.
- Watanobe H, Hayakawa Y (2003) Hypothalamic interleukin-1b and tumor necrosis factor- α , but not interleukin-6, mediate the endotoxin-induced suppression of the reproductive axis in rats. *Endocrinology* 144: 4868–4875.
- Mitchell JC, Li XF, Breen L, Thalabard JC, O'Byrne KT (2005) The role of the locus coeruleus in corticotropin-releasing hormone and stress-induced suppression of pulsatile luteinizing hormone secretion in the female rat. *Endocrinology* 146: 323–331.
- Iwasa T, Matsuzaki T, Kiyokawa M, Shimizu F, Minakuchi M, *et al.* (2007) The type 2 corticotrophin-releasing hormone receptor mediates orexin A-induced luteinising hormone suppression in ovariectomised rats. *J Neuroendocrinol* 19: 732–738.
- Tsutsui K, Saigoh E, Ukena K, Teranishi H, Fujisawa Y, *et al.* (2000) A novel avian hypothalamic peptide inhibiting gonadotropin release. *Biochem Biophys Res Commun* 275: 661–667.
- de Roux N, Genin E, Carel JC, Matsuda F, Chaussain JL, *et al.* (2003) Hypogonadotropic hypogonadism due to loss of function of the KiSS-1-derived peptide receptor GPR54. *Proc Natl Acad Sci USA* 100: 10972–10976.
- Seminara SB, Messenger S, Chatzidaki EE, Thresher RR, Acierno JS Jr, *et al.* (2003) The GPR54 gene as a regulator of puberty. *N Engl J Med* 349: 1614–1627.
- Roa J, Navarro VM, Tena-Sempere M (2011) Kisspeptins in reproductive biology: consensus knowledge and recent developments. *Biol Reprod* 85: 650–660.

21. Terasaka T, Otsuka F, Tsukamoto N, Nakamura E, Inagaki K, *et al.* (2013) Mutual interaction of kisspeptin, estrogen and bone morphogenetic protein-4 activity in GnRH regulation by GT1-7 cells. *Mol Cell Endocrinol* 381: 8–15.
22. Ducret E, Anderson GM, Herbison AE (2009) RFamide-related peptide-3, a mammalian gonadotropin-inhibitory hormone ortholog, regulates gonadotropin-releasing hormone neuron firing in the mouse. *Endocrinology* 150: 2799–2804.
23. Kriegsfeld LJ, Mei DF, Bentley GE, Ubuka T, Mason AO, *et al.* (2006) Identification and characterization of a gonadotropin-inhibitory system in the brain of mammals. *Proc Natl Acad Sci USA* 103: 2410–2415.
24. Smith JT, Coolen LM, Kriegsfeld LJ, Sari IP, Jaafarzadehshirazi MR, *et al.* (2008) Variation in kisspeptin and RFamide-related peptide (RFRP) expression and terminal connections to gonadotropin-releasing hormone neurons in the brain: a novel medium for seasonal breeding in the sheep. *Endocrinology* 149: 5770–5782.
25. Wu M, Dumalska I, Morozova E, van den Pol AN, Alreja M (2009) Gonadotropin inhibitory hormone inhibits basal forebrain vGlu T2-gonadotropin-releasing hormone neurons via a direct postsynaptic mechanism. *J Physiol* 587: 1401–1411.
26. Murakami M, Matsuzaki T, Iwasa T, Yasui T, Irahara M, *et al.* (2008) Hypophysiotropic role of RFamide-related peptide-3 in the inhibition of LH secretion in female rats. *J Endocrinol* 199: 105–112.
27. Sari IP, Rao A, Smith JT, Tilbrook AJ, Clarke IJ (2009) Effect of RF-amide-related peptide-3 on luteinizing hormone and follicle-stimulating hormone synthesis and secretion in ovine pituitary gonadotropes. *Endocrinology* 150: 5549–5556.
28. Rivier C, Vale W (1984) Influence of corticotropin-releasing factor on reproductive functions in the rat. *Endocrinology* 114: 914–921.
29. Williams CL, Nishihara M, Thalabard JC, Grosser PM, Hotchkiss J, *et al.* (1990) Corticotropin-releasing factor and gonadotropin-releasing hormone pulse generator activity in the rhesus monkey. Electrophysiological studies. *Neuroendocrinology* 52: 133–137.
30. Cates PS, Li XF, O'Byrne KT (2004) The influence of 17 β -oestradiol on corticotrophin-releasing hormone induced suppression of luteinizing hormone pulses and the role of CRH in hypoglycaemic stress-induced suppression of pulsatile LH secretion in the female rat. *Stress* 7: 113–118.
31. Chen MD, Ordog T, O'Byrne KT, Goldsmith JR, Connaughton MA, *et al.* (1996) The insulin hypoglycemia-induced inhibition of gonadotropin-releasing hormone pulse generator activity in the rhesus monkey: roles of vasopressin and corticotropin-releasing factor. *Endocrinology* 137: 2012–2021.
32. Tsukahara S, Tsukamura H, Foster DL, Maeda KI (1999) Effect of corticotropin-releasing hormone antagonist on oestrogen-dependent glucoprivic suppression of luteinizing hormone secretion in female rats. *J Neuroendocrinol* 11: 101–105.
33. Battaglia DF, Brown ME, Krasa HB, Thrun LA, Viguie C, *et al.* (1998) Systemic challenge with endotoxin stimulates corticotropin-releasing hormone and arginine vasopressin secretion into hypophyseal portal blood: coincidence with gonadotropin-releasing hormone suppression. *Endocrinology* 139: 4175–4181.
34. Li XF, Bowe JE, Lightman SL, O'Byrne KT (2005) Role of corticotropin-releasing factor receptor-2 in stress-induced suppression of pulsatile luteinizing hormone secretion in the rat. *Endocrinology* 146: 318–322.
35. Barbarino A, De Marinis L, Tofani A, Della Casa S, D'Amico C, *et al.* (1989) Corticotropin-releasing hormone inhibition of gonadotropin release and the effect of opioid blockade. *J Clin Endocrinol Metab* 68: 523–528.
36. Fourman LT, Fazeli PK (2015) Neuroendocrine causes of amenorrhea—an update. *J Clin Endocrinol Metab* 100: 812–824.
37. Biller BM, Federoff HJ, Koenig JI, Klibanski A (1990) Abnormal cortisol secretion and responses to corticotropin-releasing hormone in women with hypothalamic amenorrhea. *J Clin Endocrinol Metab* 70: 311–317.
38. Dubey AK, Plant TM (1985) A suppression of gonadotropin secretion by cortisol in castrated male rhesus monkeys (*Macaca mulatta*) mediated by the interruption of hypothalamic gonadotropin-releasing hormone release. *Biol Reprod* 33: 423–431.
39. Boccuzzi G, Angeli A, Bisbocci D, Fonzo D, Giadano GP, *et al.* (1975) Effect of synthetic luteinizing hormone releasing hormone (LH-RH) on the release of gonadotropins in Cushing's disease. *J Clin Endocrinol Metab* 40: 892–895.
40. Sakakura M, Takebe K, Nakagawa S (1975) Inhibition of luteinizing hormone secretion induced by synthetic LRH by long-term treatment with glucocorticoids in human subjects. *J Clin Endocrinol Metab* 40: 774–779.
41. Matsuwaki T, Watanabe E, Suzuki M, Yamanouchi K, Nishihara M (2003) Glucocorticoid maintains pulsatile secretion of luteinizing hormone under infectious stress condition. *Endocrinology* 144: 3477–3482.
42. Kalra PS, Sahu A, Kalra SP (1990) Interleukin-1 inhibits the ovarian steroid-induced luteinizing hormone surge and release of hypothalamic luteinizing hormone-releasing hormone in rats. *Endocrinology* 126: 2145–2152.
43. Rivier C, Vale W (1990) Cytokines act within the brain to inhibit luteinizing hormone secretion and ovulation in the rat. *Endocrinology* 127: 849–856.
44. Ebisui O, Fukata J, Tominaga T, Murakami N, Kobayashi

- H, *et al.* (1992) Roles of interleukin-1 α and -1 β in endotoxin-induced suppression of plasma gonadotropin levels in rats. *Endocrinology* 130: 3307–3313.
45. Rivest S, Lee S, Attardi B, Rivier C (1993) The chronic intracerebroventricular infusion of interleukin-1 β alters the activity of the hypothalamic-pituitary-gonadal axis of cycling rats. I. Effect on LHRH and gonadotropin biosynthesis and secretion. *Endocrinology* 133: 2424–2430.
 46. Yoo MJ, Nishihara M, Takahashi M (1997) Involvement of prostaglandins in suppression of gonadotropin-releasing hormone pulse generator activity by tumor necrosis factor- α . *J Reprod Dev* 43: 181–187.
 47. Yoo MJ, Nishihara M, Takahashi M (1997) Tumor necrosis factor- α mediates endotoxin induced suppression of gonadotropin-releasing hormone pulse generator activity in the rat. *Endocr J* 44: 141–148.
 48. Irwig MS, Fraley GS, Smith JT, Acohido BV, Popa SM, *et al.* (2004) Kisspeptin activation of gonadotropin releasing hormone neurons and regulation of KiSS-1 mRNA in the male rat. *Neuroendocrinology* 80: 264–272.
 49. Matsui H, Takatsu Y, Kumano S, Matsumoto H, Ohtaki T (2004) Peripheral administration of metastin induces marked gonadotropin release and ovulation in the rat. *Biochem Biophys Res Commun* 320: 383–388.
 50. Shahab M, Mastronardi C, Seminara SB, Crowley WF, Ojeda SR, *et al.* (2005) Increased hypothalamic GPR54 signaling: a potential mechanism for initiation of puberty in primates. *Proc Natl Acad Sci USA* 102: 2129–2134.
 51. Ohtaki T, Shintani Y, Honda S, Matsumoto H, Hori A, *et al.* (2001) Metastasis suppressor gene KiSS-1 encodes peptide ligand of a G-protein-coupled receptor. *Nature* 411: 613–617.
 52. Kotani M, Dethieux M, Vandenbogaerde A, Communi D, Vanderwinden JM, *et al.* (2001) The metastasis suppressor gene KiSS-1 encodes kisspeptins, the natural ligands of the orphan G protein-coupled receptor GPR54. *J Biol Chem* 276: 34631–34636.
 53. Terao Y, Kumano S, Takatsu Y, Hattori M, Nishimura A, *et al.* (2004) Expression of KiSS-1, a metastasis suppressor gene, in trophoblast giant cells of the rat placenta. *Biochim Biophys Acta* 1678: 102–110.
 54. Ohkura S, Takase K, Matsuyama S, Mogi K, Ichimaru T, *et al.* (2009) Gonadotrophin-releasing hormone pulse generator activity in the hypothalamus of the goat. *J Neuroendocrinol* 21: 813–821.
 55. Tomikawa J, Homma T, Tajima S, Shibata T, Inamoto Y, *et al.* (2010) Molecular characterization and estrogen regulation of hypothalamic KISS1 gene in the pig. *Biol Reprod* 82: 313–319.
 56. Inoue N, Sasagawa K, Ikai K, Sasaki Y, Tomikawa J, *et al.* (2011) Kisspeptin neurons mediate reflex ovulation in the musk shrew (*Suncus murinus*). *Proc Natl Acad Sci USA* 108: 17527–17532.
 57. Naniwa Y, Nakatsukasa K, Setsuda S, Oishi S, Fujii N, *et al.* (2013) Effects of full-length kisspeptin administration on follicular development in Japanese Black beef cows. *J Reprod Dev* 59: 588–594.
 58. Watanabe Y, Uenoyama Y, Suzuki J, Takase K, Suetomi Y, *et al.* (2014) Oestrogen-induced activation of preoptic kisspeptin neurones may be involved in the luteinising hormone surge in male and female Japanese monkeys. *J Neuroendocrinol* 26: 909–917.
 59. Matsuda F, Nakatsukasa K, Suetomi Y, Naniwa Y, Ito D, *et al.* (2014) The luteinising hormone surge-generating system is functional in male goats as in females: involvement of kisspeptin neurones in the medial preoptic area. *J Neuroendocrinol* 27: 57–65.
 60. Gottsch ML, Cunningham MJ, Smith JT, Popa SM, Acohido BV, *et al.* (2004) A role for kisspeptins in the regulation of gonadotropin secretion in the mouse. *Endocrinology* 145: 4073–4077.
 61. Smith JT, Cunningham MJ, Rissman EF, Clifton DK, Steiner RA (2005) Regulation of Kiss1 gene expression in the brain of the female mouse. *Endocrinology* 146: 3686–3692.
 62. Clarkson J, Herbison AE (2006) Postnatal development of kisspeptin neurons in mouse hypothalamus; sexual dimorphism and projections to gonadotropin-releasing hormone neurons. *Endocrinology* 147: 5817–5825.
 63. Adachi S, Yamada S, Takatsu Y, Matsui H, Kinoshita M, *et al.* (2007) Involvement of anteroventral periventricular metastin/kisspeptin neurons in estrogen positive feedback action on luteinizing hormone release in female rats. *J Reprod Dev* 53: 367–378.
 64. Smith JT, Li Q, Pereira A, Clarke IJ (2009) Kisspeptin neurons in the ovine arcuate nucleus and preoptic area are involved in the preovulatory luteinizing hormone surge. *Endocrinology* 150: 5530–5538.
 65. Kinoshita M, Tsukamura H, Adachi S, Matsui H, Uenoyama Y, *et al.* (2005) Involvement of central metastin in the regulation of preovulatory luteinizing hormone surge and estrous cyclicity in female rats. *Endocrinology* 146: 4431–4436.
 66. Smith JT, Popa SM, Clifton DK, Hoffman GE, Steiner RA (2006) Kiss1 neurons in the forebrain as central processors for generating the preovulatory luteinizing hormone surge. *J Neurosci* 26: 6687–6694.
 67. Uenoyama Y, Pheng V, Tsukamura H, Maeda KI (2016) The roles of kisspeptin revisited: inside and outside the hypothalamus. *J Reprod Dev* 62: 537–545.
 68. Tsutsui K, Ubuka T, Son YL, Bentley GE, Kriegsfeld LJ (2015) Contribution of GnIH research to the progress of reproductive neuroendocrinology. *Front Endocrinol (Lausanne)* 6: 179.
 69. Tsutsui K, Ubuka T (2016) GnIH control of feeding and reproductive behaviors. *Front Endocrinol (Lausanne)* 7:

- 170.
70. Parhar I, Ogawa S, Ubuka T (2016) Reproductive neuroendocrine pathways of social behavior. *Front Endocrinol (Lausanne)* 7: 28.
71. Ubuka T, Son YL, Tobari Y, Tsutsui K (2012) Gonadotropin-inhibitory hormone action in the brain and pituitary. *Front Endocrinol (Lausanne)* 3: 148.
72. Clarke IJ, Sari IP, Qi Y, Smith JT, Parkington HC, *et al.* (2008) Potent action of RFamide-related peptide-3 on pituitary gonadotropes indicative of a hypophysiotropic role in the negative regulation of gonadotropin secretion. *Endocrinology* 149: 5811–5821.
73. Ubuka T, Kim S, Huang YC, Reid J, Jiang J, *et al.* (2008) Gonadotropin-inhibitory hormone neurons interact directly with gonadotropin-releasing hormone-I and -II neurons in European starling brain. *Endocrinology* 149: 268–278.
74. Gibson EM, Humber SA, Jain S, Williams WP III, Zhao S, *et al.* (2008) Alterations in RFamide-related peptide expression are coordinated with the preovulatory luteinizing hormone surge. *Endocrinology* 149: 4958–4969.
75. Iwasa T, Matsuzaki T, Murakami M, Shimizu F, Kuwahara A, *et al.* (2008) Decreased expression of kisspeptin mediates acute immune/inflammatory stress-induced suppression of gonadotropin secretion in female rat. *J Endocrinol Invest* 31: 656–659.
76. Castellano JM, Bentsen AH, Romero M, Pineda R, Ruiz-Pino F, *et al.* (2010) Acute inflammation reduces kisspeptin immunoreactivity at the arcuate nucleus and decreases responsiveness to kisspeptin independently of its anorectic effects. *Am J Physiol Endocrinol Metab* 299: E54–E61.
77. Kinsey-Jones JS, Li XF, Knox AMI, Wilkinson ES, Zhu L, *et al.* (2009) Down-regulation of hypothalamic kisspeptin and its receptor, Kiss1r, mRNA expression is associated with stress-induced suppression of luteinising hormone secretion in the female rat. *J Neuroendocrinol* 21: 20–29.
78. Hirano T, Kobayashi Y, Omotehara T, Tatsumi A, Hashimoto R, *et al.* (2014) Unpredictable chronic stress-induced reproductive suppression associated with the decrease of kisspeptin immunoreactivity in male mice. *J Vet Med Sci* 76: 1201–1208.
79. Yang JA, Song CI, Hughes JK, Kreisman MJ, Parra RA, *et al.* (2017) Acute psychosocial stress inhibits LH pulsatility and Kiss1 neuronal activation in female mice. *Endocrinology* 158: 3716–3723.
80. Luo E, Stephens SB, Chaing S, Munaganuru N, Kauffman AS, *et al.* (2016) Corticosterone blocks ovarian cyclicity and the LH surge via decreased kisspeptin neuron activation in female mice. *Endocrinology* 157: 1187–1199.
81. Iwasa T, Matsuzaki T, Tungalagsuvd A, Munkhzaya M, Kawami T, *et al.* (2004) Hypothalamic Kiss1 and RFRP gene expressions are changed by a high dose of lipopolysaccharide in female rats. *Horm Behav* 66: 309–316.
82. Iwasa T, Matsuzaki T, Yano K, Irahara M (2017) Gonadotropin-inhibitory hormone plays roles in stress-induced reproductive dysfunction. *Front Endocrinol (Lausanne)* 8: 62.
83. Kirby ED, Geraghty AC, Ubuka T, Bentley GE, Kaufer D (2009) Stress increases putative gonadotropin inhibitory hormone and decreases luteinizing hormone in male rats. *Proc Natl Acad Sci USA* (2009) 106: 11324–11329.
84. Gingerich S, Wang X, Lee P, Dhillon S, Chalmers J, *et al.* (2009) The generation of an array of clonal, immortalized cell models from the rat hypothalamus: analysis of melatonin effects on kisspeptin and gonadotropin-inhibitory hormone neurons. *Neuroscience* 162: 1134–1140.
85. Gojska NM, Belsham DD (2014) Glucocorticoid receptor-mediated regulation of Rfrp (GnIH) and Gpr147 (GnIH-R) synthesis in immortalized hypothalamic neurons. *Mol Cell Endocrinol* 384: 23–31.
86. Son YL, Ubuka T, Narihiro M, Fukuda Y, Hasunuma I, *et al.* (2014) Molecular basis for the activation of gonadotropin-inhibitory hormone gene transcription by corticosterone. *Endocrinology* 155: 1817–1826.
87. Choi YJ, Habibi HR, Kil GS, Jung MM, Choi CY (2017) Effect of cortisol on gonadotropin inhibitory hormone (GnIH) in the cinnamon clownfish, *Amphiprion melanopus*. *Biochem Biophys Res Commun* 485: 342–348.
88. Peragine DE, Pokarowski M, Mendoza-Viveros L, Swift-Gallant A, Cheng HM, *et al.* (2017) RFamide-related peptide-3 (RFRP-3) suppresses sexual maturation in a eusocial mammal. *Proc Natl Acad Sci USA* 114: 1207–1212.
89. Lopes PC, Wingfield JC, Bentley GE (2012) Lipopolysaccharide injection induces rapid decrease of hypothalamic GnRH mRNA and peptide, but does not affect GnIH in zebra finches. *Horm Behav* 62: 173–179.
90. Johnson MA, Tsutsui K, Fraley GS (2007) Rat RFamide-related peptide-3 stimulates GH secretion, inhibits LH secretion, and has variable effects on sex behavior in the adult male rat. *Horm Behav* 51: 171–180.
91. Piekarski DJ, Zhao S, Jennings KJ, Iwasa T, Legan SJ, *et al.* (2013) Gonadotropin-inhibitory hormone reduces sexual motivation but not lordosis behavior in female Syrian hamsters (*Mesocricetus auratus*). *Horm Behav* 64: 501–510.
92. Geraghty AC, Muroy SE, Zhao S, Bentley GE, Kriegsfeld LJ, *et al.* (2015) Knockdown of hypothalamic RFRP3 prevents chronic stress-induced infertility and embryo resorption. *Elife* 4: e04316.