THE INTERNAL SECRETION OF SALIVARY GLAND

TOMOSABURO OGATA*

SEVEN FIGURES

This communication presents a resume of a series of investigations on the internal secretion of the salivary gland, which were carried out mostly in the Department of Pathology, Tokyo University, since the year 1928 under my supervision. The substance of the conclusion summarized here was given to my students in a farewell address, on March 18, 1943 on the occasion of my retirement from active service (Ogata, T. 1943).

The story of this research begins with the work of my associates in the Department of Pathology on the possible endocrine function of the salivary gland with special reference to the role of the internal secretion in diabetes (Hiki, *et al.* 1929, 1930, 1933, 1934. Ogata, T. 1931).

The results of studies in my department at that time have convinced me that the salivary gland might be an organ of internal secretion related to the metabolism of carbohydrates.

A new group of investigators were added my study group in 1934, and our attention was directed to the relation of the salivary secretion to the nutrition and development of the skeletal system (Takizawa, *et al.* 1935, 1936, 1937, 1938, 1939). There new studies have since clearly brought out the existence of a close relation between them and new knowledge regarding the true function of the internal secretion of the salivary gland has been added. These new observations have led us further to investigation of the nutrition and development of another hard tissue, namely, the tooth. This again has revealed the fact that there is a similarly close relation between them, as evidenced by a striking change in the incisor teeth of rats and mice following the deficiency of the internal secretion of the salivary gland (Fukushima, 1940, 1942).

With the assurance that these experimental animals, in which the effect of the salivary hormone manifested in a relatively short period of observation, could be utilized for isolating the active principle, I enlisted the cooperation of Akira Ogata, who was then professor of the Hormone Chemistry in Tokyo University for the isolation of the salivary hormone. The material prepared by him was then injected into young rats and the changes produced in the dental tissues were histologically followed by Ishii (1944) in the Department of Pathology. It was further observed by A. Ogata that the injection of the same fraction, which accelerated the growth of hard tissues, caused also a marked diminution in the

Received for publication Dec. 15, 1955.

* Professor Emeritus of Pathology, Tokyo University Medical School, and Member of the Japan Academy.

serum calcium content of the rabbit. Although, at that time, the chemical nature of this hormone was not yet elucidated, we obtained a protein substance, which is soluble in water at pH 8.0 and precipitable at pH 5.4, having a supplementary effect in cases of salivary hormone deficiency. We named this isolated salivary hormone "Parotin" (Ogata, *et al.* 1944, 1945). Furthermore Prof. Y. Ito, the successor of Prof. A. Ogata, succeeded to crystalize this hormone and described its chemical characters (Ito, *et al.* 1952).

We regret extremely, that, on account of the confusion attending the world war and the surrender of our country, it has been impossible to give this material a proper clinical trial. Since 1951, however, we have been able to organize a group of skilled clinicians, with the subsidy from the Scientific Research Fund of Education Ministry, for the purpose of giving the active substance a thorough clinical test, even though the amount available was very limited. The work which this new research group now sets out accomplish is, first of all, to elucidate the chemical structure of this hormone, and then to ascertain the biological action of the principle through its application to asialadenism (salivary hormone deficiency) and certain other diseases.

The review of the literature dealing with the internal secretion of the salivary glands is here purposely omitted, since the principal aim of the present series of investigation has very little to do with the points brought out in older literatures. Our investigations endeavor to demonstrate the endocrine function of the salivary gland concerned chiefly with the development and nutrition of the skeletal system including the tooth.

I wish to discuss first the endocrine mechanism of the salivary glands (parotid and submaxillary). These glands are not specifically provided with any endocrine tissue, comparable for instance to the islands of Langerhans of the pancreas. This function is performed, as I shall later illustrate, by the resorption of the saliva, which takes place in the striated tubule (Streifenstück, Zimmermann) of the efferent system of the salivary glands.

The striated tubules are situated between the small ducts the isthmus which are directly connected with acini and the true efferent ducts, and have relatively wide lumina as compared to the other portions of the system. Figuratively speaking, the striated tubule is a reservoir in the passage of the water-way, where the saliva is temporarily held prior to excretion into the buccal cavity. As a remarkable fact, the epithelial layer of the striated tubules is not fixed upon the basal membrane, but freely exposed to the surrounding lymph spaces (Fig. 1). It is common knowledge to regard the basal membrane as a stabilizing apparatus of the epithelial layer. The question how the striated tubules lacking in the basal membrane can support a physiological structure without the stabilizing apparatus, can best be answered by proposing the compensatory function of the firm network (Schlussleistennetz) in the inner side. Thus the epithelial layer comes in direct contact with the saliva internally and with the lymph externally.

A more important morphological characteristic is to be found in the minute structure of the epithelial cells of the striated tubules. The cytoplasm is provided with the granular system concerned with secretion and the canalicular system which effects resorption. The granular system (Iwasaki, 1937, 1938, 1939; Yuasa, 1940,

1941) resembles that of the epithelium of the main renal tubules. The mitochondria are arranged longitudinally at the base of the cell and the metachondria, presumably derived from mitochondria and regarded as secretory granules, are

Fig. 1. Schematic diagram of the striated tubule of the salivary gland. GS=granular system, M=mitochondria, S=secretory granules (metachondria), CS=canalicular system, V=basal vacuoles, C=intracellular canaliculi, N=fasten frame net (Schlussleistennetz), L=Lymph-space.

distributed above the nucleus in the cytoplasm. This system undergoes remarkable changes during the process of secretion. The secretory action of these tubules has been recognised for many years, since such distinct names as "Speichelrohr" (Pflüger) or "Sekretrohr" (Stöhr) have been proposed to designate this special function. In every instance, the secretory function of the striated tubules can be demonstrated to be most marked in the submaxillary glands, the parotids showing always a less pronounced secretory action.

As is well known, the secretion of saliva is mediated by the stimulation of either the sympathetic or the parasympathetic nerves. The saliva secreted by the stimulation of the sympathetic nerve is called sympathetic saliva, which is characterized by its small amount and thick consistency, whereas that which is secreted by the stimulation of the chorda tympani, a nerve of the parasympathetic system, is known as chorda saliva characterized by great bulk and thin consistency. We demonstrated clearly that the sympathetic saliva originates from the striated tubules, while the chorda saliva from the glandular cells of acini. Thus we know the striated tubules perform their secretory function under the influence of the sympathetic nervous system.

The canalicular system (Hiki, *et al.* 1933, 1934; Yuasa, 1940, 1941) in the cytoplasm can be demonstrated by filling the lumina of canaliculi with dye solu-

tion. This may be accomplished by examining the histological sections of the tissue after injection into the efferent ducts of dogs such dye solutions as lithium carmine or trypan blue or emulsions of sepia for 30 minutes at a low pressure (10 cm. water), taking care that the salivary gland tissue is not damaged. The dye solution injected slowly into the lumina of the striated tubules is rapidly carried into the canalicular system, just like a liquid aspirated into a pipet by suction. This resorption picture is quite different from any other previously observed in the epithelial cells of other portions, and it may be described as follows:

The canalicular system consists of basal vacuoles and intracellular canaliculi. The basal vacuoles are elongated slit-like structures arranged vertically near the cell. The position of these vacuoles corresponds to the protoplasmic space between those occupied by the vertically arranged structures belonging to the granular system. Because of the presence of these vacuoles, the so-called rod-like structure of the striated tubules can be demonstrated more clearly than that of the epithelial cells of the main renal tubules. The intracellular canaliculi originate from the uppermost portions of the basal vacuoles and course toward the internal lumina of the striated tubules in a parallel fashion. The boundary between the basal vacuoles and the emerging canaliculi may become somewhat obscure if the vacuoles are not distended, and the entire picture appears to be but one small canal. In some instances the canaliculi appears to be interlaced during their course. The canaliculi avoid in their course the nucleus of the cell and hence they seem to be closely packed around the nucleus. There are many canaliculi, the terminal portions of which are not clearly demonstrable, although there are some that are clearly seen to reach the tubular lumina. I am convinced that all canaliculi have their openings on the surface of cells. Furthermore, I am inclined to imagine that the basal vacuoles also open at the base of the cells. The orifices open as occasions demand, and expel the contents into the lymph space surrounding the striated tubule immediately below. This conclusion is verified by the existence of the same dyes in that lymph space.

The presence of the dye solution in the canalicular system is a result of active resorptive function of the epithelial cells and not of passive introduction due to pressure, as proven by the following observations. In spite of the presence of the dye in the lumina of the striated tubules, it is sometimes not resorbed into the epithelial cells of that portion. Generally speaking, the epithelial cells do not show resorption in the same degree, some cells containing large amounts while others showing no evidence of the dye.

There are other experiments (Yuasa 1940, 1941) in which a 5% egg albumin instead of a dye solution was injected under a similarly low pressure into the efferent ducts of the salivary gland in dogs. In these experiments 3.8 cc. of the solution were injected in 15 minutes into the parotid gland and 4.0 cc. in 20 minutes into the submaxillary. The appearance of the injected albumin solution in the blood stream was followed by estimating the final dilution value, or antigen titer, by the precipitin reaction of the serum. The test was carried out with samples of the dog blood removed at ten minutes intervals following the injection, and we found that the egg albumin could be definitely demonstrated within 10 minutes in the blood after injection and a considerable amount after 40 minutes.

When the same egg albumin was injected intramuscularly, the first weak reaction was obtained within 40 minutes, but when it is given by subcutaneous injection a trace could be demonstrated only after 60 minutes. These observations prove that the resorptive power of the striated tubules in the salivary glands is both rapid and active.

As these experiments were repeated, the belief that the saliva itself is resorbed as a whole in the striated tubules has been strengthened more and more. Thus the saliva, resorbed into the epithelial cells because of their specific resorptive effect, is first pushed into the lymph spaces surrounding the striated tubules and then emptied into the blood. If all the constituents of the saliva enter the blood stream, the customary idea about the saliva as a simple digestive juice must be revised. The salivary hormone (parotin) is secreted with other salivary constituents (ptyalin *etc.*) chiefly from the parotid gland, as first clearly demonsted by Ishii (1943). In the digestive tract, it is entirely inactive, but, when it enters the blood stream after resorption, it exhibits its true function as a hormone. Since this hormone is indispensable in the normal development and nutrition of the skeletal system, particularly of cartilaginous tissue, its endocrine dysfunction will lead to the pathological changes of the cartilage and bone. At the same time specific changes will also appear in the dental structure.

The endocrine dysfunction of the salivary gland leads, just as in any other instances of internal secretion, to either deficiency (asialadenism), or excess

Fig. 2. Asialadenism (rat).

Remarkable degenerative atrophy of the epiphyseal cartilage,
thin bony layer of the metaphyseal end of the diaphysis
(Osteoporosis)

(hypersialadenism). Asialadenism results from lack or deficiency of the salivary hormone, which can be produced experimentally by the extirpation of all the

salivary glands, particularly the parotids and the submaxillaries. Hypersialadenism, on the other hand, results when the animal is injected repeatedly with the gland or the salivary hormone itself. When the submaxillary glands only are removed, we have noted a peculiarly paradoxical hypersialadenism to appear instead of a relative asialadenism. This can be explained on the basis of the actual compensatory hypertrophy (hyperfunction) of the intact parotids which, as we have already mentioned, are the most important source of the salivary hormone.

According to our experimental observations, in the case of asialadenism young growing animals, as compared with controls, showed arrested development of limbs and tended to be micromelic. Histologically disturbance of the intracartilaginous ossification is demonstrated. In the epiphyseal cartilage (Fig. 2) not only is there to be seen delayed or arrested growth with insufficient provisional calcification, but degenerative changes in the cartilage known as chondromalacia is recognizable. This degeneration has been described hitherto as specific to the articular cartilage in arthritis deformans, which is characterized by the appearance of fibrous structure with the loss of homogeneous nature of the ground substance in the hyaline cartilage. Where this degeneration is most marked, there occurs a degenerative change characterized by the necrosis of all cartilaginous tissues, forming what may be termed a malatic or more correctly a liquefaction focus.

In experimental animals the chondromacia has been demonstrated only in the epiphyseal cartilage, but not in the articular cartilage, which has become generally thin as a result of hypoplasia. This is explainable on the basis of the fact that, during the developmental stage in animals, the epiphyseal cartilage undergoes proliferation much more vigorously than the articular cartilage, and the effect of hormonal deficiency appears more clearly in the form of chondromalacia at

Fig. 3. Hypersialadenism (rat).
Remarkable hyperplasia of the epiphyseal cartilage, thickened bony layer of the metaphyseal end of the diaphysis.

these locations. Beside the changes of the epiphyseal cartilage the formation of trabeculae in the ossification zone is disturbed and osteoporosis of the epiphysis is clearly demonstrable.

The skeletal picture in hypersialadenism is characterized by changes exactly opposite those seen in asialadenism, the bone and cartilage showing remarkable hyperplasia giving rise to abnormal growth in the length accompanied by osteosclerosis (Fig. 3).

The extirpation of the salivary glands (asialadenism) in albino rats leads to increased frequency in the appearance of dental caries. This high incidence was demonstrated to be not due merely to the lack of cleaning effect of the decreased flow of saliva. In one experiment, the control group showed an incidence of 30% of dental caries, while in the animals with asialadenism it reached 100%. When the saliva was injected subcutaneously into animals in which both salivary glands were removed, dental caries occurred only to the extent of 62.5%. Moreover, inflammatory changes of the gingiva are also frequently seen in asialadenism. Since such changes are almost absent in cases of hypersialadenism, the deficiency of the salivary hormone undoubtedly plays an important role in this connection.

The histological changes (Fig. 4) in asialadenism (albino rats) are characterized by both hypoplasia and absorption of dental tissues generally. The changes in the enamel organ are mostly apparent in the enamel cells below the center of the labial surface of the incisor tooth where the enamel formation occurs

Fig. 4. Asialadenism (rat).

Enamel (right-upper) and dentin (left lower). Remarkable thickened non-calcified dentin.

most vigorously. These changes consist of the irregularity, atrophy, hydropic degeneration, vacuolization, pycnosis, abnormality in nuclear locations, separation

from the enamel substance, and increase in the substance which gives a positive iron reaction. Consequently it is natural that a marked hypoplasia and at some time deficiency of the calcium deposition of enamel substance occurs in the pro-

Fig. 5. Hypersialadenism (rat).

The teeth and their supporting structures. Thickened calcified dentin and thickened alveolar processes (Osteosclerosis).

pinquity of the degenerated enamel organ. A similar hypoplasia is also seen in the dentine with widening of the predentine layer and irregular boundaries between this layer and the dentine layer proper, because of the incomplete deposition of calcium. The cement substance is also atrophic, particularly the cellular cement substance near the dental root, and the new formation of cement is completely lacking. The marked absorption picture of the cement substance and of the dentine performed by the root membrane is especially noteworthy. This process begins as a depression but some show perforation reaching the dental cavity. The pulp cavity is generally broadened as a result of decreased new formation of the dentine. The pulp tissue is also atrophic and occasionally even fibrous. The odontoblast are generally atrophic and they may even be definitely absent in the walls of the root canal. The alveolar bone is hypoplastic and often absorbed, while the root membrane is also atrophic.

In hypersialadenism the dental changes (Fig. 5) paralleling the skeletal changes exactly opposite those of asialadenism were found. In cases of hypersecretion of the salivary hormone, the entire dental tissue including the alveolar bone showed a pathologic acceleration in development. Naturally no absorption picture was encountered, and the enamel development is prominent and the dentine hypertrophic, showing even a secondary growth of the dentine. The formation of the cement substance was so active that, at the termination of the dental root, globular enlargement, resembling onion bulb in shape, was noted.

We have found moreover marked changes in other endocrine organs. Having ascertained this interrelationship between the salivary hormone and other internal secretions, we have also found an antithetical relation between asialadenism and hypersialadenism, just as in the skeletal system and the tooth. This proved the

Fig. 6. The parotid salivary gland in chondrodystrophia fetalis, 4 yrs. female.

Remarkable disappearance and atrophy of the acini with remarkable lipomatosis, isolated atrophic striated tubules of the salivary duct and the infiltration of round cells in the loose connective tissue.

correctness of our theory in regard to the endocrine dysfunction of the salivary glands.

In asialadenism the hypophysis was obviously enlarged and indicated an evident increase of the eosinophile cells, while in hypersialadenism a decrease of the eosinophile cells recognized. The thyroid gland in asialadenism showed a picture of hyperfunction, as hypertrophy of the follicular epithelial cells and liquefaction of the colloid substance, whereas in hypersialadenism the picture was that of

Table 1. Antithesis between asialadenism and hypersialadenism.

	Hypophysis	Thyroid	Pancreatic islets	Adrenal cortex	Para-thyroid	Gonads (especially female)
Asialadenism	+	+	+	-	-	±
Hypersialadenism	-	-	-	+	+	-

Signs in bold face indicate marked changes.

hypofunction. The functional appearance of other endocrine organs likewise indicate a slight antithesis between asialadenism and hypersialadenism as shown in the following table.

We have also succeeded in inhibiting the effects of asialadenism in experimental animals by continued injections of either an extract of the parotid glands or of the salivary hormone. These observations prove further the fact that the salivary glands are the organs of internal secretion.

As to the question what types of diseases are produced in man by the dysfunction of the salivary glands, we have made careful comparative studies not only of the osseous and articular changes, but also of the visceral changes, especially of the endocrine organs, between man and animal. From these observations we have been able to correlate the clinical forms due to salivary dysfunction, as tabulated in the following table.

Diseases due to dysfunction of salivary hormone.	{	Asialadenism (chondromalatic disease of bone and joints).	{	Fetal stage: Fetal chondrodystrophy (except hyperplastic type).
				Growth stage: Kaschin-Beck disease.
	{	Hypersialadenism.	{	Adult stage: Arthritis deformans Spondylitis deformans.
				Growth stage: Eunuchoidism (giant type?).

According to the findings in the experimental animals, the effect of asialadenism begin with specific degenerative changes in the cartilage called chondromalacia, especially in the epiphyseal cartilage. Hence in man the diseases must also be chronic osteoarticular disease with chondromalacia as the specific lesion.

Thus I have considered it proper to designate these conditions by the generic term "osteoarthrosis chondromalacica" (Ogata, T. 1949a, 1949b). To this group of diseases belong (1) chondrodystrophia foetalis, except the hyperplastic tipe, in the fetal stage, (2) Kaschin-Beck disease, in the juvenile (growth) stage, and (3) arthritis deformans and spondylitis deformans in the adult or senile stage.

That these diseases are all different clinical forms of asialadenism have been proved by the demonstration of pathological changes in the salivary gland, particularly in the parotid gland, in addition to the skeletal changes.

The observations on chondrodystrophia foetalis (Nozaki, *et al.* 1951; Matsueda 1952; Kohno 1941) are based on histological studies of the salivary gland in 6 cases (1 case of 8 months fetus, 2 cases of 10 months fetus, 1 cases of 11 months infant, 1 cases of 4 years child, and 1 child of 12 years and 4 months). In every instance, there was developmental arrest in the striated tubules and acinar tissues of the salivary gland.

In the fetal specimens, the development of the acini showed a developmental delay of at least 3 months in degree, while the sections of postnatal cases showed marked insufficiency in acinar formation and pronounced lipomatosis, the developmental delay being the more remarkable the younger the postnatal age of the infant studied (Fig. 6). The pathological changes were especially noticeable in the striated tubules where striation was almost absent, the nuclei of the epithelial cells being at the base, with no basal vacuolar formation, and the formation of the pericanalicular lymph spaces was indistinct. These changes are most pronounced in the case of 11 months old infant.

In every instance the changes were distinct especially in the parotid gland which clearly indicated disturbances in both salivary secretion from the acinar tissue and in resorption in the striated tubules. All these findings constitute definite evidence of this disease being an instance of asialadenism.

Fig. 7. The submaxillary gland in the Kaschin-Beck disease, 34 ys. male. Remarkable degeneration of the striated tubules (right and upper) of the salivary duct with dilatation of the pericanalicular lymphatic space of the striated tubule (middle upper).

In Kaschin-Beck disease characteristic lesions in the striated tubules of the salivary gland were demonstrated by Ogata and Takizawa (Ogata, T. *et al.* 1942, 1943).

In young patients the enlargement of the parotids and increased secretion of saliva into the oral cavity were noted during the early stage of the disease, with accompanying hypertrophy of the acinar tissue of gland. This hypertrophy may be interpreted to be compensatory in nature so that the acini may secrete saliva in excess of normal, because of the deficiency in salivary hormone (asialadenism) resulting from the decreased salivary resorption at the damaged striated tubules.

The pathological lesions of the striated tubules should be regarded as those of degenerative atrophy, as the cells are small and the striated structure is extremely indistinct. In instances showing marked skeletal change, the epithelium in the striated tubule near the isthmus showed granular destruction as well as karyorrhexis, and flattening, and disappearance of basal vacuoles, while the thick striated tubules showed here and there atrophy of epithelium and disappearance of striated structure (Fig. 7). The precipitation of proteid material in pericanalicular lymphatic spaces, the increase in the connective tissue with hyalinization around the striated tubules as well as round cell infiltrations were also noted.

These lesions as observed in Kaschin-Beck disease are ample proof that this disease is a type of asialadenism due to decreased salivary absorption in the striated tubules (Fig. 7).

Fig. 8. The parotid salivary gland in arthropathia deformans senilis, 53 yrs. male. Remarkable degeneration of the striated tubules of the salivary duct with the infiltration of round cells in the loose connective tissue. Hydropic degeneration of serous cells.

Finally, in arthritis deformans the following facts have been brought out by Takizawa (1949), Ide (1950), and Matsuyeda (1941). In patients during the first stage of this disease, in which the changes are present only in the deep layers of the articular cartilage, the weight of the salivary gland is either normal or somewhat increased and the acinar tissue shows a rather compensatory hypertrophy.

The striated tubules, however, show a characteristic degenerative atrophy in the parts near the isthmus. The epithelial cells are flattened, the striation indistinct and the lumina are dilated. During the second stage, in which ulceration appears on the surface of the articular cartilage, especially in acutely progressing cases, there occur atrophy of acinar tissue, granular degeneration and vacuolization of the epithelium of the striated tubules accumulation of proteid materials in the pericanalicular lymphatic spaces, and proliferation of the epithelium of isthmus. In more chronic cases, there appear atrophy of acinar tissue, localized disappearance of the acinar tissue, and round cell infiltration. The epithelium of the striated tubule is atrophic and flattened, with the nuclei moved toward the base, and the mitochondria disappeared, frequently there is noted cystic dilatation of the lumina of the striated tubules. During the third (last) stage, the atrophy of the acinar tissue is extreme, the epithelium of the striated tubule greatly flat-

tened, the lumina showing cystic dilatation, and pericanalicular connective tissue proliferation and hyalinization. These changes are most marked in the parotid glands but they are also seen in the submaxillary. These findings all indicate the diminution in the salivary resorption in the striated tubule, and this disease is undoubtedly due to asialadenism (Fig. 8).

Spondylitis deformans, which is identical with arthritis deformans in its nature, and frequently coexisting, has not been investigated especially, but the changes in this disease may also be safely assumed to be identical with those of arthritis deformans.

As a clinical form of hypersialadenism we have included the giant type of eunuchoidism. Upon examination of all autopsy records in cases of eunuchoidism found in the literature no reference to the salivary glands has been made. Hence, the inclusion of eunuchoidism is here merely hypothetical.

I shall finally consider the question of salivary enzyme ptyalin. Our observations clearly indicate that ptyalin as well as the salivary hormone enters the blood stream. This fact will offer an important clue as to the origin of blood amylase. Although the relation between the amylase in the blood and that which is present in visceral tissues has not been elucidated, it may be assumed that ptyalin which is constantly being resorbed into the blood from the salivary glands may play an important role in carbohydrate metabolism particularly in tissue glycogenolysis. The following facts may be offered as evidence for the validity of this assumption.

As stated at the onset, in the first stage of our research (Hiki, *et al.* 1930, 1933, 1934, Ogata, T., *et al.* 1934) the main effort was directed toward the relationship between internal secretion of the salivary gland and carbohydrate metabolism. The observation in those experimental studies may here be briefly summarized.

Using dogs as experimental animals, asialadenism was produced by total removal of the salivary glands. In such animals, upon cessation of the absorption of the saliva into the blood, a tendency for slight hypoglycemia was noted and sugar tolerance was increased. On the contrary in hypersialadenism an excess of saliva was permitted to enter the blood stream, with the result that the fasting blood sugar tended to be high and sugar tolerance somewhat lowered.

In pancreatectomized dogs asialadenism produced by the above mentioned method lessened the decrease in body weight due to diabetes, and the survival period was prolonged with relatively good nutritive state. The blood sugar remained somewhat low throughout the course, and the injection of insulin even in small doses caused a marked decrease in blood sugar. In hypersialadenism on the contrary an exactly opposite effect was noted.

When we observed these facts, we considered at first that this depended on either deficiency or excess of salivary hormone. Later, however, when the salivary hormone was found to have an action directed towards the development and nutrition of the bone and the tooth, we came to suspect this effect to be due to the action of ptyalin in the resorbed saliva.

This paper was already performed several years ago. So it does not include our recent results on the study of endocrine of salivary gland, especially on the relationship to the gerontological observations.

REFERENCE

- Fukushima, M.: The effect of salivary hormone upon the development of dental tissue, Transact. Soc. Path. Japonicae **30**: 245~259, 1940; **32**: 293~295, 1942.
- Hiki, Y., K. Akazaki, T. Ban, Y. Miyazaki, N. Takizawa and T. Yoshida: Experimental studies on diabetes mellitus, Transact. Soc. Path. Japonicae **19**: 49~56, 1929.
- Hiki, Y., T. Ban, K. Akazaki, N. Takizawa and Y. Miyazaki: Experimental studies on the salivary gland, Transact. Soc. Path. Japonicae **20**: 130~133, 1930.
- Hiki, Y., K. Akazaki, Y. Miyazaki, N. Takizawa and T. Ban, T. Yoshida and S. Nishimura: On the internal secretion of the salivary gland, Tokyo Igakukai Zasshi **47**: 773~776, 777~814, 1164~1183, 1199~1248, 1485~1546, 1814~1871, 2153~2235, 2283~2436, 1933; *ibid.* **48**: 856~878, 1934.
- Ide, G.: The pathologic anatomical and histological study of ulcerous arthropathy, Transact. Soc. Path. Japonicae **39**: editio generalis, 219~220, 1950.
- Ishii, Z.: Changes in the dental and peridental tissues in albino rats injected with parotid and submaxillary salivas of dog, Transact. Soc. Path. Japonicae **33**: 303~305, 1943.
- Ishii, Z.: Histological studies on the isolation experiment of salivary hormone, Byorigaku Zasshi **3**: 103~115, 1944.
- Ito, Y. and A. Mizutani: Studies on the salivary gland hormones, Report XIII Purification of parotin, the pH 5.4 precipitate from bovine parotid glands by fractional precipitation with sodium sulfate, J. Pharm. Soc. Japan **72**: 239~244, 244~248, 1952; Report XIV Isolation of crystalline parotin by fractional precipitation with ammonium sulfate, *ibid.* **72**: 244~248, 1952.
- Iwasaki, T.: On the secretory function of the striated tubule cells of the salivary gland, Transact. Soc. Path. Japonicae **27**: 559~562, 1937; *ibid.* **28**: 451~454, 1938.
- Iwasaki, T.: Ueber die sekretorische Funktion der Streifenstückepithelzellen der Mundspeicheldrüsen, ein Beitrag zum Wesen des Sympathicusspeichels, Japanese J. Med. Sci., V. Pathology **4**: 55~89, 1939. (In German)
- Kohn, G. and F. Kubota: A Case of chondrodystrophia foetalis in which a biopsy of the parotid gland was made, Rinshô Naika Shônica **6**: 258~261, 1941.
- Matsueda, K.: Pathologic anatomical studies on arthropathia ulcerosa.-Supp.-, Transact. Soc. Path. Japonicae **40**: editio regionalis, 32~34, 1951.
- Matsueda, K.: Pathologic anatomical studies on the changes of the organs of the internal secretion, especially of the salivary glands, in the chondrodystrophia foetalis, Transact. Soc. Path. Japonicae **41**: editio generalis, to be published, abstracted, Forty-First Annual Meeting of the Japanese pathological Society, Okayama City, April 6, 1952.
- Nozaki, K., S. Ikeda and Z. Fujita: Histological studies of the salivary gland in a 4 years old boy of typical chondrodystrophia foetalis, Transact. Soc. Path. Japonicae **40**: editio regionalis, 133~135, 1951.
- Ogata, A., Y. Ito, Y. Nozaki, S. Okabe, T. Ogata and Z. Ishii: Chemical and pathological studies on the isolation of salivary hormone, Igaku-to-Seibutsugaku **5**: 253~257, 1944.
- Ogata, A., Y. Ito, Y. Nozaki and S. Okabe: Studies on the salivary gland hormones, Reports I~XII, J. Pharm. Soc. Japan **64**: 79~88, 114~126, 146~153, 325~331, 332~336, 337~340, 1944; *ibid.* **65**: 9~13, 20~26, 27~31, 32~35, 36~42, 43~47, 1945.
- Ogata, T.: Ueber die innere Sekretion der Mundspeicheldrüsen, Transaction of the Nineth Congr. Far Eastern Ass. of Trop. Med., Nanking, No. 2, 709~713, 1934. (In German)
- Ogata, T.: Researches on the salivary hormone in the Department of Pathology, Tokyo University (Farewell Lecture, March 7, 1943), Byorigaku Zasshi **2**: 332~353, 1943; Tetsumon (Senji Tokushu-go), No. 5, 2~15, 1943; Ikai Shuho, No. 425, pp. 3~8, 1943; *ibid.* No. 426, pp. 3~7, 1943.

- Ogata, T. : On osteoarthritis chondromalacia, Niigata Med. Soc. **63** : 181~188, 1949 a.
- Ogata, T. : Diseases caused by asialadenism (osteoarthritis chondromalacia), Proc. Japan. Acad. **25** : 182~184, 1949 b. (English).
- Ogata, T., N. Takizawa, O. Wakabayashi, K. Ohta, S. Karakasa, T. Takeuchi, T. Hashimoto, M. Fukushima, M. Hirai, G. Kawase, H. Sun and F. Kuo : Studies on Kaschin-Beck disease, Report III Addendum to the Pathological observations on Kaschin-Beck disease, I. The salivary gland. Transact. Soc. Path. Japonicae **32** : 448~452, 1942.
- Ogata, T., N. Takizawa, O. Wakabayashi, K. Ohta, T. Takeuchi, K. Hashimoto, M. Fukushima, K. Suzuki, M. Hirai, G. Kawase and H. Sun : Studies on Kaschin-Beck disease, Report V. Pathological observations in Kaschin-Beck disease, III. On the changes in the viscera, particularly in the endocrine organs including the salivary gland, Transact. Soc. Path. Japonicae **33** : 325~328, 1943.
- Takizawa, N., T. Ban, N. Nagao, N. Kusano, H. Hayami, J. Hirayama, Z. Ishii, T. Matsumoto and K. Endo : Relation of salivary hormone to the development of the bone, Reports I-V, Transact. Soc. Path. Japonicae **25** : 510~511, 1935 ; *ibid.* **26** : 529~530, 1936 : *ibid.* **27** : 549~551, 1937 ; *ibid.* **28** : 454~456, 1938 ; *ibid.* **29** : 577~580, 1939.
- Takizawa, N. and G. Ide : On the changes in the organs of internal secretion, especially the salivary gland, in ulcerous arthropathy, Transact. Soc. Path. Japonicae **38** : 163~165, 1949.
- Yuasa, H. : Experimental studies on the absorptive function of the striated tubules of the salivary gland, Transact. Soc. Path. Japonicae **30** : 244, 1940 ; *ibid.* **31** : 728~730, 1941.