Fig. 1. Mud deposited by the 2011 Tohoku-oki tsunami. The mud thickness was 2 cm, and no sand layer formed at this site. Cracks developed on the dried mud layer. Photo taken at Ido, Wakabayashi, Sendai City, Miyagi Prefecture, on June 6th, 2011 (38.203760°N, 140.962993°E).

日本各地の多様な津波堆積物
Various tsunami deposits in Japan

高清水康博¹・後藤和久²・菅原大助³・藤原 治⁴・川上源太郎⁵

Yasuhiro Takashimizu¹, Kazuhisa Goto², Daisuke Sugawara³, Osamu Fujiwara⁴ and Gentaro Kawakami⁵

2017年9月17日受付。2017年10月6日受理。

¹ 新潟大学人文社会・教育科学系自然情報講座。Mathematical and Natural Sciences Course, Faculty of Education, Institute of Humanities, Social Sciences and Education, Niigata University, Niigata 950-2181, Japan

² 東北大学災害科学国際研究所。International Research Institute of Disaster Science, Tohoku University, Sendai 980-0845, Japan

³ ふじのくに地球環境史ミュージアム。Museum of Natural and Environmental History, Shizuoka, Shizuoka 422-8017, Japan

⁴ 産業技術総合研究所地質調査総合センター。Geological Survey of Japan, AIST, Tsukuba 305-8567, Japan

⁵ 北海道立総合研究機構地質研究所。Geological Survey of Hokkaido, Hokkaido Research Organization, Sapporo 060-0819, Japan

Corresponding author: takashimi@ed.niigata-u.ac.jp

日本列島の各地から報告される津波堆積物は一様ではなく、津波の性質、保存される場の状況や起源物質の違い等に支配されて多様な層相を示す。また、平成23年(2011年)東北地方太平洋沖地震による巨大津波(以下、3・11津波)は、複雑な侵食・堆積作用によって実に豊富な種類の津波堆積物が形成されることを私たちに示した。本口絵では、この特集号「日本の古津波」で紹介した日本各地の多様な津波堆積物の一部を紹介する。Fig. 1には3・11津波が形成した泥層とその乾裂の様子、Fig. 2には砂質津波堆積物の例、Fig. 3には粗粒な津波堆積物、そしてFig. 4は津波堆積物調査でよく行われるトレンチの様子を示す。

Fig. 2. Sandy tsunami deposits. (a) A late-Holocene tsunami sand bed deposited in a coastal pond on the Hamamatsu plain, Hamamatsu City, Shizuoka Prefecture, facing the eastern Nankai Trough (34.686428°N, 137.678156°E). The sand bed had clear and sharp contacts with the underlying and overlying peat, and had a multiple-layered structure, each consisting of well-sorted medium- to fine-grained sand with plant debris. Scale: 5 cm. The picture was modified from Fig. 4B of Fujiwara and Sawai (2014). (b) Tsunami deposits on the rice paddy at Arahama, Wakabayashi, Sendai City, Miyagi Prefecture (38.231269°N, 140.977507°E). The shore-normal distance of the site is 1.5 km. Medium-grained sand was deposited on the erosional surface of the soil with a thickness of 20 cm. There were faint laminations in the sand layer with small rip-up clasts. Photo taken on May 11, 2011. (c) A 17th-century tsunami deposit overlying peat with erosional contact at Yufutsu, Tomakomai City, Hokkaido Prefecture (42.628900°N, 141.710511°E). A thick (12–18 cm) medium-grained tsunami sand bed was deposited below the AD 1663 Us-b tephra, separated by a very thin lenticular layer of peat (arrows). Scale: 20 cm.

Fig. 3. Coarse-grained tsunami deposits. (a) Tsunami debris, sand, and fallen power poles (left) and iron poles (right) in the residential area of Arahama, Wakabayashi, Sendai City, Miyagi Prefecture (38.219908°N, 140.976444°E). Pine trees were also transported from the beach area by the tsunami. Photo taken on April 20, 2011. (b) A large *Porites* boulder deposited by the AD 1771 Meiwa tsunami on east coast of Ishigaki Island, Okinawa Prefecture, Japan. (24.519883°N, 124.299417°E) (c) Cobble layer (arrow) intercalated in the organic muddy deposits of a coastal marsh environment, Sado Island, Niigata Prefecture (38.319417°N, 138.463158°E). The layer may have been formed by a tsunami of 9th or 11th century. Scale: 1 m.

Fig. 4. Outcrop of the trench survey of paleotsunami deposits at Noda Village, Iwate Prefecture (40.116017°N, 141.824617°E). The possible AD 869 Jogan tsunami bed (arrow) was intercalated in organic-rich mud layers (Takada et al., 2016). Scale: 1 m.

文 献

- 藤原 治・澤井祐紀(Fujiwara, O. and Sawai Y.), 2014, 静岡県沿岸の古地震・津波堆積物調査(Paleoearthquake and paleotsunami surveys in the Shizuoka Prefecture). 地質分野研究企画室編, 巨大地震による複合的地質災害に関する調査・研究報告書(*In Research Planning Office for Geological Survey and Applied Geoscience, ed., Reports of Research and Investigation on Multiple Geological Hazards Caused by Huge Earthquakes*), 産総研地質調査総合センター (Geol. Surv. Japan, AIST), **66**, 39–48.
- 高田圭太・宍倉正展・今井健太郎・蛭名裕一・後藤和久・越谷 信.....岩手県県土整備部河川課(Takada, K., Shishikura, M., Imai, K., Ebina, Y., Goto, K., Koshiya, S.,... River Division Department of prefectural Land Development, Iwate Prefecture Government), 2016, 岩手県沿岸における津波堆積物の分布とその年代 (Distribution and ages of tsunami deposits along the Pacific Coast of the Iwate Prefecture). 活断層・古地震研報(*Annu. Rep. Active Fault Paleoseismicity Res.*), **16**, 1–52.