The Malampaya Sound Group in the Calamian Islands, North Palawan Block (Philippines)

Lawrence R. Zamoras* and
Atsushi Matsuoka**

* Graduate School of Science and Technology, Niigata University, Niigata 950-2181, Japan.

** Department of Geology, Faculty of Science, Niigata University, Niigata 950-2181, Japan.

Received 23, January, 2001

Accepted 24, March, 2001

Fig. 1. The linear topographic expression of the chert-clastic sequences showing the chert mountain ranges (Balnek Mountains) with intervals of narrow valleys representing the clastics.

The Malampaya Sound Group is predominantly composed of chert (Liminangcong Formation), clastics (Guinlo Formation) and scattered limestone units that characterize the North Palawan Block (Zamoras and Matsuoka, 2001). In the Calamian Islands (grouping Busuanga, Coron, Culion and other smaller islands) the Liminangcong Formation is represented by Permian to Late Jurassic chert; the Guinlo Formation is represented by Middle-Late Jurassic to Early Cretaceous clastics.

The common succession from chert to clastics with generally conformable contact is thus referred to as a chert-clastic sequence. In Busuanga Island, linear topographic patterns reveal the parallel distribution of chert-clastic sequences dipping generally NW in the west, and NE in the central and eastern portions. This chert-clastic repetition indicates an imbricate structure typical to offscrape accretion complexes.

Fig. 2. Geologic map of the Calamian Islands (Busuanga, Culion, Coron and smaller islands).

Fig. 3. The Tulbuan Plain composed of Jurassic (late Callovian-Oxfordian) clastics (Guinlo Formation).

Fig. 4. Typical thinly bedded Triassic chert sequence (part of the Liminancong Formation).

Fig. 5. Late Triassic Coron Limestone viewed in Cayangan Lake in Coron Island.

Fig. 6. Molluscan assemblages reflective of lagoonal environmental setting of the Coron Limestone. The pelecypod fossils probably include *Paramegalodus* which was reported by Wolfart et al. (1986).

Reference

- Wolfart, R., Cepek, P., Gramann, F., Kemper, E. and Porth, H., 1986, Stratigraphy of Palawan Island, Philippines. *Newsl. Stratigr.*, **16**, 19–48.
- Zamoras, L.R., and Matsuoka, A., 2001, Malampaya Sound Group: a Jurassic-Early Cretaceous accretionary complex in Busuanga Island, North Palawan Block (Philippines). *Jour. Geol. Soc. Japan*, **107**, 316–336.