On the Occurrence of the Lower Tertiary Formation in the Province of Hōki.

By

M. Yokoyama.

In my brief note on the distribution of the Lower and Upper Tertiary Formations in Japan published in the May number of this year of the Journal of Geography of the Tokyo Geographical Society, I referred to a find of a specimen of *Crassatella fusca* Yok., a Japanese Palaeogene fossil, near Tari in the county of Hino, Hōki Province. The specimen was obtained by Mr. S. Kitamura, a geologist, at the chromite mine of Nogumi where he was told that it had been picked up somewhere in the neighbourhood long previously. This firmly establishes the occurrence of the Lower Tertiary in Hōki, further confirmed by another shell which the same geologist found among the fossils kept in the primary school of Tari and which is said to have been collected at Araya, a place not far from the school. This shell is a large and splendid *Pholadomya* much elongated in shape, pointed at beak, gradually dilated towards behind and ornamented with many radiating ribs on the surface. On comparing it with the already known species of the genus, it shows the closest resemblance to *Pholadomya puschi* Goldf. and *Pholadomya weissii* Phil., both found in the Oligocene of Germany. The chief difference between these two latter lies in the presence or absence of ribs on the posterior surface; namely, in *P. weissii* they are present, while in *P. puschi* they are absent on the posterior one-third of the surface. The Japanese form, so far as can be ascertained, also lacks them like *P. puschi* on the posterior surface, although

fourth of the whole surface. It is very likely that it is to be compared with *P. puschi* rather than with *P. weissii*.

The occurrence of the Lower Tertiary fossils in Hōki is quite interesting, as up to this time, so far as I know, they have never been met with outside Kyushu.

Explanation of Plate

Pholadomya cf. *puschi* Goldf. in nat. size. from Araya, Hōki.