THE SARUKUBO CONGLOMERATE

KAZUYOSHI OKAMI*

(Received June 14, 1972)

Introduction

The occurrence of orthoquartzite pebbles hitherto regarded as of Precambrian origin has been recently reported from several localities in Japan, such as the Shimanto terrain (TOKUOKA, 1967, 1970; THE KISHU SHIMANTO RESEARCH GROUP, 1968), Tanba belt (RESEARCH GROUP OF THE TANBA BELT, 1971) and the Mino belt (ADACHI, 1971). The precambrian terrain from which the orthoquartzite pebbles are inferred to have been derived, is postulated by THE KISHU SHIMANTO RESEARCH GROUP and ADACHI to have existed under the oldest dated rocks, the Silurian System, and to have extensive distribution southeastwards (Pacific side) and also northwestwards (Japan sea side) beyond the area of the present Japanese Islands. The radiometric dating of the silimanite gneiss pebbles of the Kamiaso Conglomerate of the Mino belt indicates the Precambrian age (SHIBATA *et al.*, 1971), and may support the postulation expressed by THE KISHU SHIMANTO RESEARCH GROUP (1968) and ADACHI (1971).

The orthoquartzite pebbles newly discovered from the Mesozoic Sarukubo Conglomerate distributed in the Torinoko mountain block of the Yamizo Mountains, Ibaraki Prefecture, Northeast Japan form the subject of the present article. The occurrence of the orthoquartzite pebbles from Northeast Japan may be important for consideration of the geotectonic relationship between Southwest and Northeast Japan because the hitherto known orthoquartzite pebbles of Japan have been restricted to Southwest Japan. The pebbles may also serve for the reconstruction of the geotectonic history of the Abukuma Plateau and the Yamizo Mountains.

In this paper the Sarukubo Conglomerate is described and discussed mainly from the view point of sedimentary petrology with special reference to the geotectonic history of the southern Abukuma Plateau.

* Department of Mining and Civil Engineering, Faculty of Technology, Iwate University.

Acknowledgments

I wish to express my thanks to Professor Kotora HATAI for his critical reading of the manuscript and Dr. Masafumi MURATA for his valuable advices, both of the Institute of Geology and Paleontology of the Tohoku University. Thanks are due to Professor Toshiji OYAMA and Drs. Toshio SAITO and Haruyuki TAKAHASHI of the Department of Geology, Ibaraki University for their kind advices and suggestions. Appreciation is also expressed to Mr. Kentaro OKAWA and some students of the Ibaraki University for their kind advices in many different ways.

Outline of the Geology

The geology of the Yamizo Mountains that include the Yamizo (strict sense), Torinoko and Toriashi (Keisoku of some authors) mountain blocks from north to south have been made by FUJIMOTO and HATAKEYAMA (1938), KAWADA and KANOMATA (1948), KAWADA (1953), KANO (1960), KANOMATA (1961), KASAI and KIMURA (1971) and KASAI and OYAMA (1972).

Those mountain blocks consist mainly of sandstone and shale with intercalated layers of chert and small limestone lenses of Mesozoic and Paleozoic age. According to KANOMATA (1961), the rocks can be classified into the Paleozoic Kasama Group and the Mesozoic Yamizo Group. The latter group is divided into the five formations, namely Kunimiyama, Toriashisan, Yamizosan, Torinokosan and Karasuyama formations in descending order superposed with conformity from east to west (Table 1). The Mesozoic age of the Yamizo Group was determined by the radiolaria assemblage from the chert and siliceous shale of various localities and a fossil wood *Dadoxylon* (*Araucarioxylon*?) (KANOMATA, 1961). Recently, from the Torinokosan Formation *Kranaosphinctes matsushimai* (YOKOYAMA) (Jurassic Oxfordian age) has been found (SUZUKI and SATO, 1972).

The brief descriptions, according to KANOMATA (1961), of each formation of the Yamizo Group

Fig. 1. Geologic map of the Sarukubo area, Ibaraki Prefecture.

Table 1. Stratigraphic unit of the Yamizo Mountains (KANOMATA, 1961).

YAMIZO GROUP	Karasuyama F.
	Torinokosan F.
	Yamizosan F.
	Toriashisan F.
	Kunimiyama F.
	Fault
KASAMA GROUP	

are as follows;

The Kunimiyama Formation with a thickness of about 3000 m, distributed in the Kunimiyama area of Toriashi mountain block, consists of sandstone and shale and their alternation. In the

southern part of Toriashi mountain block the Kunimiyama Formation is in contact with the Paleozoic Kasama Group by fault.

The Toriashisan Formation consists of sandstone and shale with three layers of chert that can be traced throughout the whole area of the Yamizo Mountains. The thickness of the formation is estimated to be about 4000 m.

The Yamizosan Formation about 2000 m in thickness distributed typically in the central part of the Torinoko mountain block is composed of sandstone and shale; the sandstone is dominant.

The Torinokosan Formation is estimated to be about 2000 m in thickness. It consists of sand-

stone and shale similar to the subjacent Yamizosan Formation. The distinction between the two formations resides in the fact that shale is dominant in the Torinokosan Formation and sandstone is dominant in the Yamizosan Formation.

The Karasuyama Formation, the uppermost unit of the Yamizo Group, consists of sandstone and shale intercalated with a thin layer of chert and conglomerate (Sarukubo Conglomerate). The thickness of the formation is about 2500 m.

A brief description of the Yamizo Group was given by KANOMATA as mentioned above, but since then, with the progress of study, different opinions have been introduced, particularly concerning the geostructure of the Yamizo Mountains (KASAI and KIMURA, 1971). The current view denies the westward dipping homoclinal sequence of the strata, and expresses the repetition of strata in three times caused by reverse faults trending parallel to the strike of the beds. Because of the repetition of the strata by reverse faults, the nomenclature of the formation becomes complicated. For example in the Toriashi mountain block, KANOMATA's Kunimiyama Formation is divided into the Kunimiyama (lower) and Keisokusan (upper) formations, and Toriashisan, Yamizosan and Torinokosan formations of KANOMATA (the Karasuyama Formation does not exist in this block) are reclassified into the Kasama (Paleozoic), Kunimiyama and Keisokusan formations (OYAMA *et al.*, MS). In the Torinoko and Yamizo mountain blocks, the geostructure and stratigraphy is more complicated than in the Toriashi mountain block. Further discussions are not present here because it is out of the scope of the present article.

The area dealt with in this article is situated at the southern periphery of the Torinoko mountain (Fig. 1). The Yamizo Group of this area has a general strike and dip of N 20°W, 50°W, and is structurally homoclinal, the strata superposed with conformity from east to west. The typical distribution area of the Sarukubo Conglomerate is situated in a narrow area, along the road leading from Sarukubo to Yoshida in Naka County, Ibaraki Prefecture. The area is cut by two faults each of N-S trend (Fig. 1). The Sarukubo Conglomerate in the southern part of this area is covered by the Tertiary rocks with unconformity. The Sarukubo Conglomerate continues northward to the adjacent area. The rocks in the Sarukubo area have a general strike and dip of N 50°E, 45°W, which is different from the trend in the adjacent area.

Sedimentary Petrology of the Sarukubo Conglomerate

1. Sarukubo Conglomerate (General Setting)

The Sarukubo Conglomerate (KAWADA and KANOMATA, 1948) is typically developed in the cliff of the road leading from Sarukubo to Yoshida

Fig. 2. Route map along the road leading from Sarukubo to Yoshida.

Fig. 3. Columnar section of the Sarukubo area, and the paleocurrent direction (arrow represents azimuthal arithmetic mean).

in Naka County, Ibaraki Prefecture. Its distribution continues to the summit of the mountain in the adjacent area. The outcrops of the Conglomerate in the Sarukubo and adjacent areas is small as shown in the route map (Fig. 2). The sequence of the Sarukubo Conglomerate observed along the road mentioned above is, from the lower, cobble size conglomerate intercalated with thin sandstone and shale, thin laminated sandy shale, alternation of fine grained sandstone and sandy shale intercalating coarse grained sandstone (3—4 m). The lower part of the Sarukubo Conglomerate is unknown because it is overlain by the Tertiary rock with unconformity. The Sarukubo Conglomerate mentioned above represents the lower conglomerate unit of the above sequence, and has a thickness of about 60 m (Fig.

3). The Sarukubo Conglomerate consists of three cobble size conglomerate layers, two pebble size conglomerate layers, with thin sandstone and shale layers. The thickest layer of cobble size conglomerate is the uppermost one which has a thickness about 20 m including a thin sandy shale layer in the middle part. The alternation layers in the upper part of the Sarukubo Conglomerate include the abundant plant fragments.

The Sarukubo Conglomerate was considered to be the basal conglomerate of the Torinokosan Formation by KAWADA (1953), and subsequently, KANOMATA (1961) regarded this conglomerate to be an intraformational conglomerate of the Karasuyama Formation. However, recently KASAI and OYAMA (1972) suggested that the Sarukubo Conglomerate represents the basal conglomerate of the Keisokusan Formation (same as Karasuyama Formation) as is evident from the sharp difference in the lithofacies of the superjacent and subjacent strata. The writer holds the view that the Sarukubo Conglomerate is the basal conglomerate of the Keisokusan Formation (KASAI and OYAMA, 1972), because the lithofacies of the superjacent sandstone changes abruptly to arkosic and the because of the discordant distribution of the Sarukubo Conglomerate to the subjacent strata (Fig. 1). The alternation layers of the strata superjacent to the conglomerate contain abundant plant leaves and fragments and the internal structure of the alternation layers does not show turbidite facies. The alternation layers are thus considered to represent the mixed depositional environment.

The Sarukubo Conglomerate is composed dominantly of cobble size pebbles and sporadically of boulder size pebbles attaining as much as 40 cm in maximum diameter (Pl. 1). The pebbles are shaped variously from well rounded to angular (POWERS, 1952). The matrix consists of coarse grained sandstone and the conglomerate is loosely packed. The sandy matrix of the conglomerate is very arkosic, consisting of quartz 55%, feldspars 22%, mica (biotite) 0.4%, rock fragments (sandstone, quartz porphyry, aplite, clayey material and so on) 10% and matrix (under 5 ϕ) about 5% (Table 3). The grains are angular and not beyond the subangular grade. The textural parameters of the sandstone constituting the matrix of the conglomerate are of mean size 1.6 ϕ , sorting degree 1.17, skewness 0.25 (Table 4).

2. Constituents and Composition of the Sarukubo Conglomerate

The Sarukubo Conglomerate comprises pebbles of sandstone, slate, chert, limestone, clayey rocks,

Table 2. Composition of the pebble species of the Sarukubo Conglomerate.
N: total_number_of counted.

	Sarukubo		Kitada No. 1		Kitada No. 2	
chert	38	40.0%	52	50.5%	35	48.6%
sandstone	11	11.5	24	23.3	14	19.4
slate	11	11.5	11	10.7	9	12.4
orthoquartzite	4	4.2	1	0.9	5	6.9
limestone	1	1.0	1	0.9	1	1.4
vein quartz	8	8.4	1	0.9	1	1.4
granite	4	4.2	2	1.8	1	1.4
porphyrite	2	2.1	—	—	—	—
porphyry	1	1.0	1	0.9	—	—
others	15	15.8	10	9.7	6	8.3
	N=95	99.7	N=103	99.6	N=72	99.8

Fig. 4. Composition of the pebbles in the Sarukubo Conglomerate.

orthoquartzite, porphyrite, granitic rocks and hornfels. Pebbles of sedimentary rocks predominate. The sandstone pebbles consist of those various sizes. The pebbles of clayey rocks may be the result of contemporaneous erosion. The pebbles of chert are blue to gray in color and angular in shape; they resemble in all features the chert of the underlying bed. According to KANO (1960) the granitic rocks consist of biotite granite, porphyroidal two mica granite and biotite syenite, and all of them are characterized by their

sheared structure. The limestone pebbles are fossiliferous, and KAWADA and KANOMATA (1948) reported from them such Permian fossils as *Neoschwagerina* sp., *Pseudofusulina* cf. *krafti* (SCHELLWIEN), *Schwagerina* sp., *Triticites* sp., and *Martina* sp. The pebbles of limestone are weathered easily and in the outcrop commonly only their cast are found. Except for the orthoquartzite pebble, the pebbles of the sedimentary rocks resemble the underlying sedimentary rocks. Igneous rocks and orthoquartzite pebbles are exotic.

3. Composition of the Sarukubo Conglomerate

The composition of the pebbles of the Sarukubo Conglomerate was analysed by the method of counting about 100 pieces larger than 3 cm in diameter. The composition is shown by the percentage of each species to the total number counted (HIRABAYASHI, 1969). As shown in Table 2 and Fig. 4, the Sarukubo Conglomerate contains abundant chert pebbles, more than 40 percent. The pebbles of sandstone are next abundant; both hard and soft ones occur in amount of about 15 percent. Slate pebbles amount to about 10 percent of the total. The composition of the Conglomerate at two places, Sarukubo and Kitada, resemble each other but differ in the frequency of the orthoquartzite pebbles as shown in the data of Kitada. The difference in the frequency may be due in part to the comparatively small pebble size. In general the frequency of the orthoquartzite pebbles is about 4 percent.

4. Description of the Orthoquartzite

The orthoquartzite pebble is pale chocolate to whitish gray in color. The chocolate color is due to weathering. The size of the orthoquartzite pebble is variable, the largest pebble attains 25 cm in diameter and the smallest measures about 3 cm in diameter. The orthoquartzite pebbles are well rounded and thus may be distinguished readily from other kind of pebbles of sedimentary rocks. The orthoquartzite pebbles consist essentially of rounded detrital quartz grains, with small amounts of feldspars and tourmaline grains. The opaque minerals are contained in a traceable amount. The grains are cemented tightly by secondary quartz. Clayey material is also contained in a very small amount in the interstitial pore space. On the quartz grains overgrowth is observed in optical continuity and the original grains are fringed by dust ring (Pl. 2). The quartz grains consist of monocrystalline quartz without undulate extinction and polycrystalline undulate extinctions are not found.

Feldspars are K-feldspar and plagioclase. K-feldspar shows micropertite structure with albite and plagioclase is albite twinned. The composition of the orthoquartzite is quartz 85—90%, feldspars 0.1—2.7%, tourmaline 1% to traceable, secondary quartz 8—10% and matrix 2% (Table 3). All of the analysed samples are plotted into the orthoquartzite clan (Fig. 5).

The textural parameters of the grain size distribution of the orthoquartzite, calculated according to FOLK's formula (1965), are as follows;

Fig. 5. Composition of the orthoquartzite pebbles of the Sarukubo Conglomerate according to FOLK's classification (1965). All of them are plotted in the orthoquartzite clan. Q; all types of quartz except stretched metaquartzite; plus chert, F; single grains of all type of feldspars; igneous rock fragments both plutonic and extrusive, M; metamorphic rock fragments and mica.

the mean size range -0.19 to 0.55ϕ , sorting 0.41 to 0.69 , skewness about 0.10 and kurtosis about 0.11 ; so the orthoquartzite is texturally coarse grained, well to moderately well sorted, normal distributed sandstone. The sandstone possessing such textural parameters and composition may be considered to have been deposited through the heavy and or long abrasion process as in an eolian environment (FOLK, 1965). The characters of the orthoquartzite pebble from the Sarukubo Conglomerate well resemble the orthoquartzite pebbles recorded from the Shimanto terrain (THE KISHU SHIMANTO RESEARCH GROUP, 1968; TOKUOKA, 1970), Kamiaso Conglomerate (ADACHI, 1971) and Tanba belt (RESEARCH GROUP OF TANBA BELT, 1971). They are considered to have been derived from Precambrian rocks because of the non-development of the orthoquartzite in the geological systems of the Japanese Islands. The data of the textural parameters and composition are shown in Table 3 and 4.

The roundness of the grains was estimated by the method of CAILLEUX. The grains are mostly spheroidal, pollard grains are rare (Pl. 2), the value of the roundness is about 0.60 and the highest value obtained is 0.78 (Table 4).

5. Roundness and Sphericity of the Pebbles

Table 3. Mineral composition of the orthoquartzite pebbles and the matrix of the Sarukubo Conglomerate.

	Q	F	M	T	H	RF	UK	OV	Matrix
A-1	84.1	2.7	*	1.9	tr.	*	0.1	8.5	2.5
A-2	84.1	1.9	*	tr.	*	*	0.1	12.1	1.7
B-2	90.0	0.1	*	tr.	*	*	0.1	8.7	1.1
B-3	89.4	1.5	*	tr.	*	0.04	0.1	7.8	2.4
C-1	85.0	1.3	*	1.1	*	0.13	tr.	11.0	1.3
Matrix	55.0	22.1	0.4	*	0.2	17.6	0.4	*	4.7

Q; quartz, F; feldspars, M; mica, T; tourmaline, H; heavy minerals, RF; rock fragments, OV; overgrowth, A, B and C; sample number of the orthoquartzite pebble of 72116-7A, B and C, and 1, 2 and 3 is the section no. made from each pebble.

Table 4. Grain size distribution and roundness of consisting grains of the orthoquartzite pebbles and the matrix of conglomerate of the Sarukubo Conglomerate; definition of parameters after FOLK (1965) and CAILLEUX (1947). Sample nos. are same to the nos. in Table 3.

	Mean size	Sorting	Skewness	Kurtosis	Roundness
A-1	0.51	0.43	0.30	1.17	0.63-0.60
A-2	-0.19	0.71	0.11	1.23	0.78-0.69
B-2	0.30	0.62	0.05	1.47	0.57-0.51
B-3	0.23	0.63	0.22	1.12	0.54-0.50
C-1	0.28	0.69	0.02	1.23	0.55-0.45
Matrix	1.60	1.17	0.25	1.53	*

To determine the provenance of the pebbles included in the Sarukubo Conglomerate, analyses were made concerning the difference of the abrasion process, and the roundness and sphericity of every rock species by the method of DOBKINS and FOLK (1970). The data obtained from the analyses are listed in Table 5. The hardness and foliation (structure) of the rocks are closely related to the shape of the pebbles and also to their environment (KUENEN, 1956; DRAKE, 1970; DOBKINS and FOLK, 1970). The influences of the foliation of the rocks may be represented in the parameter of the sphericity (DRAKE, 1970). The values of the sphericity of the pebbles are mostly above 0.60, so the pebbles are spheroidal in shape and this suggests that strongly foliated to platy rocks do not exist.

The degree of the roundness ranges from 0.10 to 0.77. However there exists clear difference in roundness according to the rock species (Table 5). The pebbles in the Sarukubo Conglomerate are classifiable into two groups of low and high roundness (Fig. 6). Chert, slate, hornfels and sandstone pebbles belong to the former category, porphyrite, granite and orthoquartzite pebbles to the latter (Pl. 2). These differences are distinct in the outcrops. The roundness of each pebble species is, 0.10 to 0.35 for chert, 0.35 to 0.53 for sand-

stone, 0.58 to 0.62 for porphyrite, 0.56 to 0.84 for granite and 0.37 to 0.68 for orthoquartzite. KUENEN (1956) studied experimentally the abrasion process and the relationships between the roundness and distance of the transportation and hardness of the pebbles. He found that long distance transportation makes the value of the roundness increase and the degree of the increase of the roundness value is corresponding to the hardness of each pebble species. From the roundness, chert pebbles are inferred not to have suffered hard abrasion and it is suggested that they may have been supplied from rather short distance and or not through heavy abrasion environment. The granite pebbles are well rounded, and the high value of the roundness may be due to the moderate hardness (KUENEN, 1956), or long distance transportation. From the resemblance of the characters of the pebbles of the sedimentary rocks to the underlying sedimentary rocks, the pebbles except for the orthoquartzite ones, show roundness of low value and thus may have been derived directly from an adjacent area. The chert pebbles also show low value of roundness.

The granite and orthoquartzite pebbles show high roundness. According to KUENEN (1956), granite pebbles can gain a rounded shape (roundness 0.40 to 0.60) by moderate distance transporta-

Table 5. Sphericity and roundness of the pebbles of the Sarukubo Conglomerate; definition of parameter after DOBKINS and FOLK (1970).

	L	S	I	SP	Di	Dk	R
Orthoquartzite	125.0	56.0	91.2	0.65	45.0	23.0	0.51
	37.7	24.3	32.8	0.78	16.0	6.0	0.37
	39.5	24.5	32.2	0.72	15.0	6.0	0.40
	*113.0	54.0	72.0	0.65	57.0	16.0	0.42
	*55.0	36.0	39.0	0.54	19.0	12.0	0.63
	*90.0	45.0	80.0	0.65	40.0	27.0	0.67
	*34.0	28.0	31.0	0.89	13.0	6.0	0.46
	125.0	80.0	102.0	0.79	51.0	35.0	0.68
	120.0	68.0	92.6	0.74	36.0	24.0	0.67
Sandstone	51.1	19.0	36.2	0.58	18.0	9.0	0.47
	56.5	39.6	47.0	0.83	23.0	6.0	0.26
	38.3	27.2	35.0	0.81	17.0	9.0	0.53
	73.5	37.9	55.0	0.71	27.0	10.0	0.37
	*45.0	25.0	30.0	0.77	15.0	6.0	0.40
Chert	79.3	48.8	74.0	0.74	34.0	12.0	0.35
	36.0	21.5	31.2	0.74	14.0	3.0	0.21
	52.2	30.0	36.0	0.78	18.0	3.0	0.16
	50.0	27.3	48.0	0.68	19.0	2.0	0.11
	41.0	22.0	38.0	0.68	15.0	4.0	0.26
	36.0	22.5	29.0	0.84	13.0	3.0	0.22
	34.8	19.3	31.0	0.70	12.0	4.0	0.33
	*52.0	31.0	48.0	0.38	23.0	4.0	0.17
	*57.0	45.0	46.0	0.77	17.0	5.0	0.29
Granite	96.2	57.8	72.2	0.77	35.0	27.0	0.77
	*65.0	20.0	54.0	0.48	27.0	15.0	0.56
	70.0	58.0	64.0	0.90	32.0	27.0	0.84
Porphyrite	38.8	19.2	32.0	0.66	16.0	10.0	0.62
	*30.0	23.0	27.0	0.86	12.0	7.0	0.58

* collected from Kitada, size in mm., L, S and I; long, short and intermediate axis, SP; sphericity, Di; radius of the largest inscribed circle, Dk; radius of the sharpest developed corner, R; roundness

tion of about 30 to 50 Km. He stated from the relationship between roundness and weight loss, that the weight loss of the granite pebble is ten times that of the chert pebble. Therefore it may be said that the granite pebble can obtain a rounded form by short distance transportation. Since the orthoquartzite pebble is very hard like that of the chert pebble, and possesses high roundness, the long distance transportation and or hard abrasion is necessary to obtain a rounded shape. This is an important factor for the consideration of the provenance of the orthoquartzite pebbles.

Discussion (Provenance Problem)

The direction of the Paleocurrent during deposition of the conglomerate is judged to be from SE to NW as determined from the current ripple marks developed in the alternation of the sandstone and shale overlying the Sarukubo Conglomerate (Fig. 3, Pl. 1). The paleocurrent direction supports the KANO's suggestion (1960) that the granite pebbles in the Sarukubo Con-

glomerate, characterized by the large amount of microcline and sheared structure, are exotic in the Yamizo Mountains and may have been derived from the Hitachi sheared granite, about 35 Km east of Sarukubo (Fig. 1). KANO (1960) also stated that the geologic age of the Sarukubo Conglomerate, from its characteristics and resemblance with known granites in Japan, may be Mesozoic (Jurassic) in age.

The high roundness of the pebbles of the granite and orthoquartzite is the reflection of the abrasion process during transportation. The chert pebbles as hard as orthoquartzite pebbles have angular shape, do not seem to have suffered strong abrasion, resemble the underlying chert bed, and are inferred to have been supplied from the underlying bed. And other pebbles of the sedimentary rocks are similarly considered. The orthoquartzite pebble can not be considered to have originated from some subjacent unit as chert, because of high roundness, which suggests long distance transportation. But the sedimentary

the orthoquartzite pebbles discovered from the Shimanto terrain (THE KISHU SHIMANTO RESEARCH GROUP, 1968; TOKUOKA, 1970), Tanba belt (RESEARCH GROUP OF THE TANBA BELT, 1971) and Kamiaso Conglomerate (ADACHI, 1971). All have been considered to have been derived from the Precambrian. The orthoquartzite pebbles of the Sarukubo Conglomerate are not considered to have been derived from the Precambrian directly but to have been reworked from a Permian conglomerate now not represented in the terrain. The possibility of the existence of the Precambrian beneath the southern Abukuma Plateau is reasonable, and perhaps rocks of such age make the basement of the Paleozoic rocks in this area.

5). The Permian limestone pebbles included in the Sarukubo Conglomerate (KAWADA and KANOMATA, 1948) and Momiya Conglomerate (SUZUKI and OMORI, 1953) suggest the former existence of the Permian System in the southern Abukuma Plateau.

References

- ADACHI, M. (1971), Permian intraformational conglomerate at Kamiaso, Gifu Prefecture, Central Japan. *Jour. Geol. Soc. Japan*, vol. 77, p. 471—482.
- DOBKINS, J. E. and FOLK, R. L. (1970), Shape development of Tahitinui. *Jour. Sed. Petr.*, vol. 40, p. 1167—1203.
- DRAKE, L. D. (1970), Rock texture: An important factor for clast shape studies. *Jour. Sed. Petr.*, vol. 40, p. 1356—1361.
- FOLK, R. L. (1965), Petrology of Sedimentary rocks. *Hemphill's Austin*, 159p.
- FUJIMOTO, H. and HATAKEYAMA, H. (1938), Upper Paleozoic fossils from the Toriashi mountain block, Ibaraki-Ken (preliminarily report). *Jour. Geol. Soc. Japan*, vol. 45, p. 377—378. (in Japanese).
- HIRABAYASHI, T. (1966), Neogene conglomerate and Cenozoic geohistory of the northern Fossa Magna in Central Japan (part 1). *Earth Science*, vol. 24, p. 79—92.
- KANO, H. (1960), On a granite-bearing conglomerate from the Torinoko massif and its geologic significance — Studies on the granite bearing conglomerate in Japan, no. 8. *Sci. Rep. Tohoku Univ.*, 2nd. ser. (Geol.), Special Volume no. 4, p. 476—481 (in Japanese with English abstract).
- KANOMATA, N. (1961), The geology of the Yamizo, Torinoko and Toriashi mountain blocks and their geologic age. *Jour. Fac. Art. and Sci., Chiba Univ.*, vol. 3, p. 351—367.
- KASAI, K. and KIMURA, K. (1971), Stratigraphic study on the chert bed of the Yamizo Mountains. *Abst. Program, 1971 Annual Meeting Geol. Soc. Japan*, p. 466 (In Japanese).
- and OYAMA, T. (1972), On the Sarukubo Conglomerate of the Torinoko mountain block, Ibaraki Prefecture. *Abst. Program, 1972 Annual Meeting Geol. Soc. Japan*, p. 137 (in Japanese).
- KAWADA, K. (1953), Geological studies on the Yamizo, Torinoko and Toriashi mountain blocks and their neighbourhood in the Northeastern Kanto district. *Sci. Rep. Tokyo Bunrika Daigaku, sec. C*, vol. 2, p. 217—307.
- and KANOMATA, N. (1948), On the Paleozoic fossils from pebbles in the conglomerate found in the southern foot of Torinoko mountain, Ibaraki Prefecture. *Jour. Geol. Soc. Japan*, vol. 54, p. 17—18 (in Japanese).
- KUENEN, Ph. H. (1956), Experimental abrasion of pebbles, 2 rolling by current. *Jour. Geol.*, vol. 64, p. 336—368.
- KURODA, Y. (1963), Mutual relations between the metamorphic rocks in the Northeast Japan. *Earth Science*, no. 67, p. 21—29 (in Japanese with English abstract).
- MINATO, M. (1968), Basement complex and Paleozoic orogeny in Japan. *Pacific Geology*, vol. 1, p. 85—96, *Tsukiji Shokan, Tokyo*.
- OYAMA, T., SAITO, T. and TAKAHASHI, H. (1972), subsurface geological map of Ibaraki Prefecture, 1:200,000 (MS).
- POWERS, M. C. (1952), A new roundness scale for sedimentary particles. *Jour. Sed. Petr.*, vol. 23, p. 117—119.
- RESEARCH GROUP OF THE ABUKUMA PLATEAU (1969), Polymetamorphism in the Abukuma Plateau with special regards to the discovery of staurolite and kyanite. *Mem. Geol. Soc. Japan*, no. 4, p. 83—98 (in Japanese with English abstract).
- SHIBATA, K., ADACHI, M. and MIZUTANI, S. (1971), Precambrian rocks in Permian conglomerate from Central Japan. *Jour. Geol. Soc. Japan*, vol. 77, p. 507—514.
- SUZUKI, A. and SATO, T. (1972), Discovery of Jurassic Ammonite from Toriashi mountain. *Jour. Geol. Soc. Japan*, vol. 78, p. 123—215 (in Japanese).
- SUZUKI, K. and OMORI, M. (1953), Etude de la géologie stratigraphique de la système Tertiaire dans le district de Ota-machi, Ibaraki Pref. (Etude de la géologie historique de la système Néozoïque au Bord de S-W du Plateau Abukuma part II). *Jour. Geol. Soc. Japan*, vol. 59, p. 35—46 (in Japanese with French abstract).
- THE KISHU SHIMANTO RESEARCH GROUP (1968), The study of the Shimanto terrain in the Kii Peninsula, Southwest Japan (part 2) — The present status of the research and southern Formerland in the Pacific Ocean —. *Earth Science*, vol. 22, p. 224—231 (in Japanese with English abstract).
- TOKUOKA, T. (1967), The Shimanto terrain in the Kii Peninsula, Southwest Japan, — With reference to its geologic development viewed from coarser clastic sediments. *Mem. Fac. Sci., Kyoto Univ., ser. Geol. and Mineral.*, vol. 34, p. 35—74.
- (1970), Orthoquartzite gravels in the Paleogene Muro Group, Southwest Japan. *ibid.*, vol. 37, p. 113—132.
- WATANABE, J. and BIKERMAN, M. (1971), K-Ar age

of the Nishidohira gneiss complex in the Abukuma mountains, Japan. *Earth Science*, vol. 25, p. 23—26

(in Japanese with English abstract).

猿 久 保 礫 岩

大 上 和 良

(要 旨)

近年、先カンブリア系起源と推定されるオルソコーツアイト礫の産出が各地から報告され、日本列島の基盤に関する新たな知見を与えている。しかしながら、このオルソコーツアイト礫の産出はフォッサ・マグナ以西の西南日本に限られていた。

筆者は今回、八溝山系に分布するジュラ系の基底礫岩、猿久保礫岩中に東北日本では初めてのオルソコーツアイト礫が多量に含有されていることを発見した。このオルソコーツアイト礫は円磨度が非常に高く、鏡下観察の結果西南日本で報告されたものに酷似している。さらに、同礫岩に対して詳細な堆積岩石学的検討を加えた結果、このオルソコーツアイト礫は再食礫であると結論され、供給源は礫岩層の上位の地層中に発達する澱痕および共存する花崗岩礫の性質、二畳紀化石を含む石灰岩礫等から南部阿武隈高原に発達していた二畳系礫岩と推定される。現在、同高原南部には二畳系の分布が知られていないが、同高原南端に発達する鮮新統茂宮礫岩は二畳紀化

石を含む石灰岩の巨礫を多数含み、二畳系が阿武隈高原南部に発達していたことを裏づけている。

さらに、このオルソコーツアイト礫の発見は南部阿武隈高原に分布する西堂平片麻岩に認められる複変成作用の存在事実とあわせ、同地に先カンブリア系の存在を、さらに同系が阿武隈および八溝山系の基盤を形成することを示唆する根拠を与えた。

地 名	
Karasuyama ……鳥山	Ogawa ……緒川
Kasama ……笠間	Sarukubo ……猿久保
Keisokusan (Toriashisan)	Toriashisan (Keisokusan)
……………鶏足山	……………鶏足山
Kitada ……北田	Torinokosan ……鷺子山
Kunimiyama ……国見山	Yamizo ……八溝
Momiya ……茂宮	Yoshida ……吉田
Naka ……那珂	

Explanation of Plates

Plate I

- 1 View of the uppermost horizon of the Sarukubo Conglomerate. An outcrop along the road leading from Sarukubo to Yoshida, Ogawa Village, Ibaraki Prefecture.
- 2 Cobble size conglomerate of the Sarukubo Conglomerate.
- 3 Pebble size conglomerate of the Sarukubo Conglomerate; middle horizon.
- 4 Ripple marks developed in the sandstone of the alternation of sandstone and shale in a cliff along the road from Sarukubo to Yoshida.

Plate II

- 1 Photomicrograph of an orthoquartzite pebble (A-2) from the Sarukubo Conglomerate. Note the well-rounded detrital quartz grains cemented by secondary quartz. Crossed nicols.
- 2 Photomicrograph of an orthoquartzite pebble (A-1) from the Sarukubo Conglomerate. Note the well rounded detrital quartz grains cemented by secondary quartz as optically continued and containing a very small amount of clay minerals. Crossed nicols.
- 3 Pebbles of the Sarukubo Conglomerate showing different shapes according to rock species; 1, granite, 2, orthoquartzite, 3, 4 and 5, chert pebbles.