15) A Study on Coronary Damage in SHRSP — The Determination of Medial Tissue Collagen — 1) Kengi Suzuki, and Haruhito Kawashimo 2) Nobuyuki Abe, Kokichi Takeuchi, Mitsuyo Saito, Chikao Arai, and Motoharu Hasegawa 3) Yoshitaka Takayama 4) Hiroyuki Ito, and Kozo Okamoto 1) Epidemic Institute of Arteriosclerosis 2) Department of Clinical Functional Physiology, Faculty of Medicine, Toho University, Tokyo-to 143 3) Department of Medical Engineering, Faculty of Medicine, Toho University 4) Department of Pathology, Kinki University School of Medicine.

The medial collagen tissues of the coronary artery in the SHRSP and WKY were observed histochemically and quantitatively for the comparison. The subjects used in the present study were 60 WKY's and 77 SHRSP's 137 cases in total. Collagen was assayed by the MSP method after Van Gieson's staining. The adventitia in WKY was rich in collagen, but it is less in the media and only fibrous collagen observed. The adventitia was thick in the SHRSP with increase in collagen component, but such increase in collagen was not observed in the media as in the case of the WKY. The mean collagen content in various age population was as follows: 30.2%E in less than 100 days WKY (SHRSP), 29.3(30.3)%E in the 100 days', 30.1 (35.7)%E in the 200 days', 29.1 (33.2)%E in the 300 days', 29.1 (33.2)%E in the 400 days', 38.1(33.7)%E in the 500 days', and 33.5%E in the 600 days'. and 33.5%E in the 600 days'. Collagen content increased slightly with aging, but there was no statistically significant difference among the both strain groups and age groups