11) **Impaired noradrenergic projection via baroreflex to hypothalamus in SHR**

Shingo Kawasaki, Kazuo Takeda, Hiroshi Ito, Masahiro Hirata, Tetsuo Nakata,
Junko Hayashi, Minako Oguro, Susumu Sasaki and Masao Nakagawa

The Second Department of Internal Medicine , Kyoto Prefectural University of Medicine

(Purpose) It is well known that the input of the afferent nerve of baroreflex is located in the nucleus tractus solitarius(NTS), and NTS also connects with other various nucleus, which includes the hypothalamus. But there are few reports to clear this mechanism. In this study, in order to clarify the role of noradrenergic projections to posterior hypothalamus via baroreflex on the development of hypertension, extracellular norepinephrine (NE) content of posterior hypothalamus was measured in both SHR and WKY after sinoaortic denervation(SAD) or SHAM operation.

(Materials and methods) 9 weeks-old male WKY and SHR were used. The rats were divided into two groups. On one group was performed the SAD operation by Kriegar method and on the other group was performed the SHAM operation; all rats were anesthetized with pentobarbital(30mg/kg i.p.). Systolic blood pressure and heart rate were measured by tail cuff plethystogram before the operation and also 24 hours after the operation. After anesthetizing with urethane(1.2g/kg), rats were placed on the stereotaxic apparatus (David-Kopf) and the brain dialysis probe was placed in the left posterior hypothalamus using standard stereotaxic technique coordinates(AP 4.6mm, L 1.0mm, DV 2.5mm), and terminal of tube from brain dialysis probe was connected to the microinfusion pump, and perfused over 90 minutes with artificial cerebrospinal fluid at the rate of 1μ l/minute. Norepinephrines in the perfusate were analyzed by the method of radioenzymatic assay. To confirm whether SAD was successful or not, disappearance of bradycardic response to intravenous NE injection was examined after the experiments.

(Results) In WKY, blood pressure(BP) twenty four hours after SAD were significantly higher than those of SHAM operated rats (SHAM $121 \pm 1 \rightarrow 127 \pm 1$ mmHg , SAD $124 \pm 1 \rightarrow 158 \pm 4$ mmHg : $P < 0.001$). Contrarily, in SHR they were no significant difference between SAD and SHAM groups (SHAM $179 \pm 2 \rightarrow 170 \pm 6$ mmHg , SAD $178 \pm 3 \rightarrow 176 \pm 5$ mmHg : NS). Extracellular NE content after SAD were also significantly higher in WKY than those of SHAM operated rats (SHAM 95 ± 19 pg/ml, SAD 268 ± 38 pg/ml: $P < 0.01$), but in SHR they were no significant difference between two groups (SHAM 206 ± 25 pg/ml, SAD 266 ± 24 pg/ml: NS).

(Conclusion) These results suggest that in SHR noradrenergic pathway via the baroreflex is impaired and this mechanism may play an important role in development and maintenance of hypertension in SHR.