Calcium Antagonists Protect against Nephrosclerosis in Partially Nephrectomized Spontaneously Hypertensive Rats.

Keiji Fujiwara, Koichi Hayashi, Masanori Honda, Yuri Ozawa, Hirobumi Tokuyama, Hiroto Matsuda, and Takao Saruta. Department of Internal Medicine, School of Medicine, Keio University, Tokyo, 160.

It has been well established that systemic hypertension not only causes renal dysfunction but also aggravates nephrosclerosis. Although blood pressure reduction *per se* favors renal protection, there exists some controversy that calcium antagonists may exacerbate renal function in renal disorders. In the present study, we investigated the renal protective effects of newly developed calcium antagonists, pranidipine (OPC-13340), aranidipine (MPC-1304), and its metabolite (MP-3442), on the progression of renal failure in salt-loaded partially nephrectomized spontaneously hypertensive rats (SHR).

Methods: SHR with 5/6 nephrectomy were divided into four groups: group 1 as control (n=11); group 2, pranidipine-treated (3mg/kg/day, n=8); group 3, aranidipine-treated (3mg/kg/day, n=8); and group 4, MP-3442-treated (10mg/kg/day, n=8). The rats were given these drugs and a high-salt diet (5% NaCl) for 8 weeks. During the experiments, systolic blood pressure (SBP), and daily urinary protein excretion (UPE) were measured. At the end of the study, blood was collected and renal tissues were obtained for light microscopic examination.

Results: SBP was markedly reduced by 8-week antihypertensive treatment (control, 264 ± 6 mmHg; pranidipine, 158 ± 8 mmHg; aranidipine, 176 ± 11 mmHg; MP-3442, 181 ± 4 mmHg). Glomerular and arteriolar sclerosis developed markedly in the control group, but were markedly prevented in pranidipine- and aranidipine-treated groups. UPE tended to decrease in pranidipine- and aranidipine-treated groups [233 ± 5 (p=0.07) and 237 ± 9 mg/day (p=0.09), respectively, vs. 322 ± 33 mg/day for control]; MP-3442 (251 ± 18 mg/day) did not suppress the increase in UPE. No difference in histopathological score, serum creatinine concentration, or UPE was noted among these treated groups.

Conclusion: Both pranidipine and aranidipine attenuated SBP increase and ameliorated glomerular and arteriolar injury. In contrast, MP-3442 reduced SBP, but failed to reduce UPE, a finding in accordance with our previous observations that nifedipine did not reduce proteinuria despite a reduction in blood pressure. Thus, the renal protective effects of calcium antagonists may differ, which may be associated with divergent effects of calcium antagonists in renal diseases.