Titanium Dioxide Exposure Induces Acute Eosinophilic Lung Inflammation in Rabbits

Gil Soon CHOI^{1†}, Chulho OAK^{1†}, Bong-Kwon CHUN², Donald WILSON³, Tae Won JANG¹, Hee-Kyoo KIM¹, Mannhong JUNG¹, Engin TUTKUN⁴ and Eun-Kee PARK^{5*}

¹Department of Internal Medicine, Kosin University College of Medicine, Republic of Korea

²Department of Pathology, Kosin University College of Medicine, Republic of Korea

³Department of Occupational Toxicology, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan

⁴Ankara Occupational Diseases Hospital, Ministry of Health, Turkey

⁵Department of Medical Humanities and Social Medicine, Kosin University College of Medicine, Republic of Korea

Received May 22, 2013 and accepted March 7, 2014

Published online in J-STAGE April 4, 2014

Abstract: Titanium dioxide (TiO₂) is increasingly widely used in industrial, commercial and home products. TiO₂ aggravates respiratory symptoms by induction of pulmonary inflammation although the mechanisms have not been well investigated. We aimed to investigate lung inflammation in rabbits after intratracheal instillation of P25 TiO₂. One ml of 10, 50 and 250 µg of P25 TiO₂ was instilled into one of the lungs of rabbits, chest computed-tomography was performed, and bronchoalveolar lavage (BAL) fluid was collected before, at 1 and 24 h after P25 TiO₂ exposure. Changes in inflammatory cells in the BAL fluids were measured. Lung pathological assay was also carried out at 24 h after P25 TiO₂ exposure. Ground glass opacities were noted in both lungs 1 h after P25 TiO₂ and saline (control) instillation. Although the control lung showed complete resolution at 24 h, the lung exposed to P25 TiO₂ showed persistent ground glass opacities at 24 h. The eosinophil counts in BAL fluid were significantly increased after P25 TiO₂ exposure. P25 TiO₂ induced a dose dependent increase of eosinophils in BAL fluid but no significant differences in neutrophil and lymphocyte cell counts were detected. The present findings suggest that P25 TiO₂ induces lung inflammation in rabbits which is associated with eosinophilic inflammation.

Key words: Allergy, Eosinophil, Inflammation, P25 TiO₂ nanoparticle

Introduction

Over the past decades, advances in nanotechnology have led to their rapid applications in the fields of medi-

cine, pharmaceuticals, biotechnology, energy production and environmental sciences¹⁾. The increasing use of nanomaterials in various products at workplaces and in the home setting, including many consumer items such as clothing and plastic wares²⁾ therefore pose an obvious risk to humans.

Titanium dioxide (TiO₂) nanoparticles (NPs) are one of the most abundantly utilized nanomaterials because of their chemical stability, low toxicity and relatively cheap price³⁾. It is used as a white pigment in paint, food color-

*To whom correspondence should be addressed.

E-mail: ekpark@kosin.ac.kr

†These authors contributed equally to this work.

© National Institute of Occupational Safety and Health

ing, as an ultraviolet blocker in cosmetics, disinfectant in environment and wastewater, and as a photosensitizer for photodynamic therapy. Oral ingestion and entry through the dermal route are mainly mediated by therapeutic or cosmetic application^{4, 5}). The respiratory route is most important because the intake of NPs into the body is from atmospheric air via the upper respiratory tract⁶). TiO₂ NPs are increasingly being manufactured, leading to increased occupational exposure and release into the atmospheric environment. Nano-sized particles are generally more toxic to the lung than their larger-sized counterparts⁷) which are why there has recently been increasing concern about the impact of TiO₂ NPs in the lung.

Several epidemiological studies have reported that TiO₂ NPs exposure at the work place aggravate respiratory symptoms^{8, 9}). Besides, earlier studies indicated that inhalation TiO₂ NPs can induce pulmonary response such as inflammation, fibrosis, emphysema-like lung injury, and lung cancer^{6, 10, 11}). However, the pulmonary effects of TiO₂ NPs are not fully understood. Previous animal studies have shown that exposure to TiO₂ NPs causes oxidative stress, induce lung inflammation in the airways and alveolar spaces^{12, 13}). Moreover, it has been reported that TiO₂ NPs are able to induce neutrophilic pulmonary inflammation^{14, 15}). Recent studies found that TiO₂ NPs cause lung inflammation by activation of T-helper 2 cells and that the exposure of high concentration of TiO₂ NPs in the lung induced an innate immune activation^{16, 17}). Although *in vitro* and *in vivo* studies suggest that TiO₂ NPs cause various forms of pulmonary inflammation^{11, 18}), to our knowledge, relatively few studies have investigated the pulmonary effect in a rabbit model, and its pathogenic mechanism. The present study investigated the effect of TiO₂ NPs on rabbit lungs, evaluated by lung image analysis, bronchoalveolar lavage (BAL) fluids examination, and histopathologic analysis.

Materials and Methods

Titanium dioxide (TiO₂)

P25 TiO₂ nanoparticles (Brunauer-Emmett-Teller (BET) specific surface area of 53.8 m²/g) were obtained from Degussa. The particles were suspended as follows: 1.5 g of P25 TiO₂ powder was suspended in 100 ml of distilled water in a pyrex glass beaker, and sonicated for 15 min by a Branson Digital Cell Disruptor Sonifier 250 (Branson, USA) with a double-stepped microtip (3 mm diameter); this process was repeated 3 times. To stabilize temperature during sonication, the beaker was placed in a bucket of ice

throughout the process. The particle suspension was then centrifuged at 3,000 × g for 20 min at 20 °C. The supernatant was carefully collected and filtered through a 1 µm filter to remove the large agglomerates (>1 µm). A given volume of particle suspension was evaporated, after which the weight of the remaining evaporate was measured and P25 TiO₂ concentration determined (w/v; in mg/ml). The hydrodynamic size distribution by number of P25 TiO₂ particles suspended in water was analyzed using a Dynamic Light Scattering Zetasizer Nano (Malvern Instruments, UK), and the average particle size was calculated to be 61.9 ± 5.1 nm.

Animals and study protocol

Nine Male New Zealand white rabbits (Taesung Laboratory Animal Science, Busan, Republic of Korea) weighing 3.0 to 3.5 kg were used for this experiment. The rabbits were housed at 20–25 °C and 50–70% relative humidity with a 12 h light/dark cycle. They had free access to water and diet and were acclimatized for at least 1 wk before starting the experiments. Radiologic image analysis (computer-tomography (CT)) was performed to ascertain lung inflammation at 1 and 24 h after P25 TiO₂ exposure, and also to investigate the pathogenic mechanism, bronchoalveolar lavage (BAL) was performed at before P25 TiO₂ exposure, 1 and 24 h after P25 TiO₂ exposure. For further histological analysis, all rabbits were euthanized using CO₂ gas at 24 h after P25 TiO₂ exposure.

Animal experimental procedure was approved by the Animal Research Ethical Committee in Kosin Gospel Hospital, Busan, Republic of Korea.

P25 TiO₂ nanoparticles exposure

Rabbits were anesthetized by intramuscular injection of ketamine 5 mg/kg (Huons Co., Korea) and xylazine 0.8 mg/kg (Bayer, Republic of Korea). Oxygen saturation was monitored by pulse oxymeter in the ear. Transbronchial P25 TiO₂ instillation was performed using an ultrathin bronchoscope (BF-XP260F, Olympus; Tokyo, Japan). The ultrathin bronchoscope was inserted into the target bronchus as deep as possible under direct vision. The instillation catheter was inserted beyond the visible bronchus through working channel. One ml of 10 µg P25 TiO₂ was once instilled into the right lung through the catheter and 1 ml of normal saline (as control) was instilled into the left lung (N=3). One ml of 50 and 250 µg P25 TiO₂ were instilled in the same way (N=3 in each group).

Fig. 1. Lung inflammatory change by chest CT image analysis. Both lungs were instilled with 10 $\mu\text{g}/\text{ml}$ of P25 TiO₂ in the right lung and normal saline in the left lung. (A) normal lung at baseline; (B) lung inflammation with ground glass opacity in both lung fields 1 h after instillation of P25 TiO₂ NP; (C) after 24 h of P25 TiO₂ NP exposure, ground glass opacities in the left lung (control) disappeared but they remained in the right lower lung field.

Fig. 2. Total cell count change in bronchoalveolar lavage fluid (BALF). Total cell count in BALF was significantly increased at 1 and 24 h after P25 TiO₂ NP challenge. The bars represent mean $\pm$ SE; * $p < 0.05$ significantly different from baseline; Mann-Whitney U test.

Bronchoalveolar Lavage and Cell Counting

Bronchoalveolar lavage (BAL) was performed before P25 TiO₂ exposure, at 1 and 24 h after P25 TiO₂ exposure through an ultrathin bronchoscope, which was wedged into the 1st branch bronchus of the right lung. Sterile saline solution (2 ml) was instilled through the bronchoscope. The fluid was immediately recovered by gentle suction after each instillation. The measurement of recovered fluids showed an approximately 90% recovery. To maximize cell viability, the harvested BALF was immediately placed

on ice and centrifuged at $1,000 \times g$ for 10 min. The supernatants were immediately stored at -80°C for further analysis. The cell pellet was used to prepare slides, which were stained according to the May-Grunwald and Giemsa procedures to morphologically assess the cells in the fluid. The differential cell counts were then counted by hemocytometer.

Lung Pathologic Examination Assay

The lung was harvested for pathologic examination at 24 h after P25 TiO₂ exposure. Tissue pretreatments and preparation of hematoxylin and eosin (H&E) stained slices were carried out as previously described¹⁶. They were evaluated by light microscopy.

Statistical Analysis

Results were expressed as mean $\pm$ standard error (SE). Mann-Whitney U test was used in the case of two independent samples. All analyses were carried out using SPSS 16.0 (SPSS Inc., Chicago, IL, USA). A $p < 0.05$ was considered statistically significant.

Results

Lung image analysis

To ascertain lung inflammation by P25 TiO₂ exposure in the rabbit, chest CT was performed after P25 TiO₂ instillation. Both lungs were clear before the experiment (Fig. 1A), but at 1 h ground glass opacities (GGO) were noted in each lung instilled with P25 TiO₂ (10 $\mu\text{g}/\text{ml}$) and normal saline in Fig. 1B. At 24 h after exposure, persistent lung inflam-

Fig. 3. Eosinophil percentage change in bronchoalveolar lavage fluid (BALF). The increase in eosinophil percentage was noted at 1 and 24 h after P25 TiO₂ NP challenge with dose-dependent responses. (A) the eosinophil count in BALF increased at 1 and 24 h after P25 TiO₂ NP challenges; (B) the photos of eosinophil count in BALF at the dose of 50 µg/ml compared to baseline. The bars represent mean $\pm$ SE; * p <0.05 significantly different from baseline; Mann-Whitney U test.

Fig. 4. Pathological examination of the lungs following P25 TiO₂ NP challenge. After 24 h of P25 TiO₂ instillation, (A) severe eosinophilic inflammation was noted in the lung that was instilled with 50 µg/ml; (B) mild eosinophil inflammation was noted in the lung exposed to the dose of 10 µg/ml; (C) normal baseline. Left pane at magnification of 100 $\times$ and right pane at a magnification of 1,000 $\times$.

mation with GGO was noted in the right lung instilled with P25 TiO₂, while lung inflammation disappeared in the control lung, instilled with normal saline in Fig. 1C. Similar results were obtained from the experiments using 50 and 250 µg/ml of P25 TiO₂ (data not shown).

Inflammatory cell changes in BAL fluids

After one single instillation of 10 µg/ml P25 TiO₂ in the rabbit lung, the total cell count in BAL fluids was increased at 1 h ($p=0.24$) and 24 h ($p=0.42$) (Fig. 2). At 50 and 250 µg/ml of P25 TiO₂ the total cell count in BAL fluids was high at 1 h and 24 h compared to the baseline.

There was an especially marked increase in eosinophil percentage at 1 h ($4.0 \pm 0.4\%$) and 24 h ($17.5 \pm 5.5\%$) after 10 µg/ml of P25 TiO₂ ($p<0.05$, Fig. 3A). When exposed to higher concentrations of P25 TiO₂ (50 and 250 µg/ml), a dose-dependent increase in eosinophil percentage was detected at 1 h and 24 h (Fig. 3A). The eosinophil count on BAL fluids increased at 1 and 24 h after 50 µg/ml P25 TiO₂ exposure when compared to baseline (Fig. 3B).

Pathogenic changes after P25 TiO2 exposure

Analysis of BAL fluids confirmed that P25 TiO₂ induced inflammation in lung tissues. After 24 h of P25 TiO₂ instillation (50 µg/ml), severe eosinophilic inflammation was noted in the alveolar, peribronchial and perivascular regions, with moderate hemorrhage (Fig. 4A). Mild eosinophilic inflammation was noted in lung tissues exposed to 10 µg/ml P25 TiO₂ (Fig. 4B). On the other hand, in the control lung (not exposed to P25 TiO₂), there was no eosinophilic inflammation (Fig. 4C).

Discussion

Exposure to TiO₂ NPs during production or use is most likely to occur via different routes such as skin penetration, ingestion, or inhalation, but, it is believed that the lung is

the most important target organ¹⁹). Although there have been a few studies on the pulmonary effects of TiO₂ exposure, to the best of our knowledge, the current study is the first investigation in a rabbit model evaluating acute lung changes after P25 TiO₂ exposure. Generally, mice or rat models are widely used in research due to ease of handling and their being relatively cost-effective compared to other models. Despite these advantages, mice and humans have considerable differences in lung structure and function which limits their suitability to lung disease studies^{20, 21}). The rabbit is known to have a very similarity to human in terms of airway anatomy and responses to inflammatory mediators, although it has not been widely used probably due to limitation of cost and reagent availability. Differences in the pulmonary effects of TiO₂ NPs in mouse and rat species have previously been reported^{22, 26}). Considering rabbit is phylogenetically closer to human than rodents, our study may further provide important knowledge to understanding the acute lung impacts of TiO₂ exposure in human.

TiO₂ NPs was previously reported to induce pulmonary response¹¹), which has been mainly evaluated subacute and chronic change by histopathological analysis. There were limitations in that the sequential acute changes following TiO₂ exposure were not investigated. This is why image analysis was utilized in our study to evaluate acute lung inflammation following TiO₂ NPs intratracheal instillation. We observed ground glass opacities of acute pneumonitis at 1 h after single P25 TiO₂ NPs exposure. Furthermore we observed persistent pneumonitis in the P25 TiO₂ exposed lung, as well as newly developed pneumonitis in the P25 TiO₂ unexposed opposite lung at 24 h. These results indicate that single instillation of P25 TiO₂ can induce severe acute pulmonary inflammation. Moreover, previous studies reported that high dose TiO₂ NPs cause more severe lung inflammation compared with that of low dose of TiO₂, as well as induce persistent pulmonary inflammation^{23–25}). This information may have clinical implications regarding safety in handling of TiO₂ NPs.

To understand the pathogenic mechanism of this acute lung inflammation by P25 TiO₂ exposure, in the present study, BAL fluids and histopathology of lung sections were examined. We found that eosinophils were significantly increased during acute response (1 h) after P25 TiO₂ NPs exposure, which persisted at 24 h in BAL fluids. Furthermore, eosinophil increases showed a dose-dependent pattern. This finding is consistent with those from the study in rat¹⁶). Although we did not observe any significant changes of other inflammatory cells, some stud-

ies in mouse models have shown increase of neutrophils and macrophages in the lung, as well as epithelial change after challenge with TiO₂^{10, 22, 26–28}). However, these inflammatory changes were associated with different properties of TiO₂ NPs, like crystal structure, surface chemistry, and surface area^{12, 29–31}). Therefore, it may be difficult to compare results between NP studies. TiO₂ NP exposure in rats induced innate immune activation of eosinophils in the acute and long-lasting lymphocyte responses¹²). Recently, it has been reported that lung challenge with TiO₂ NP in mice cause inflammation by activation of T-helper 2 cells³²). It has also been shown that lung exposure to TiO₂ NP aggravates an asthmatic response and also promotes allergic sensitization and lung inflammation in a mouse model^{17, 33}). In addition, our histopathologic analysis showed severe eosinophil inflammation in the lung after P25 TiO₂ challenge, compared with those of the control lung that was not exposed to P25 TiO₂. Considering that eosinophils are the main effector cells in an allergic inflammation such as asthma^{34, 35}), we speculate that P25 TiO₂ induce allergic lung inflammation by eosinophil activation. Additional investigations are needed to elucidate how eosinophil activation happens following TiO₂ exposure.

In conclusion, this is the first study that investigated lung inflammation after P25 TiO₂ exposure in a rabbit model and found the particles to induce eosinophilic lung inflammation. Further research is necessary to investigate the mechanism and implications of this eosinophil activation induced by TiO₂ NP.

Acknowledgements

This study was supported by a grant from Kosin University College of Medicine, Busan, Republic of Korea. We thank Professor Toshihiko Myojo (Department of Environmental Health Engineering, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health, Japan) for his valuable comments.

References

- 1) Liu WT (2006) Nanoparticles and their biological and environmental applications. *J Biosci Bioeng* **102**, 1–7. [Medline] [CrossRef]
- 2) Hoet PH, Bröske-Hohlfeld I, Salata OV (2004) Nanoparticles—known and unknown health risks. *J Nanobiotechnology* **2**, 12. [Medline] [CrossRef]
- 3) Chen X, Mao SS (2007) Titanium dioxide nanomaterials:

- synthesis, properties, modifications, and applications. *Chem Rev* **107**, 2891–959. [[Medline](#)] [[CrossRef](#)]
- 4) Smijs TG, Bouwstra JA (2010) Focus on skin as a possible port of entry for solid nanoparticles and the toxicological impact. *J Biomed Nanotechnol* **6**, 469–84. [[Medline](#)] [[CrossRef](#)]
  - 5) Yah CS, Simate GS, Iyuke SE (2012) Nanoparticles toxicity and their routes of exposures. *Pak J Pharm Sci* **25**, 477–91. [[Medline](#)]
  - 6) Li JJ, Muralikrishnan S, Ng CT, Yung LY, Bay BH (2010) Nanoparticle-induced pulmonary toxicity. *Exp Biol Med* (Maywood) **235**, 1025–33. [[Medline](#)] [[CrossRef](#)]
  - 7) Bakand S, Hayes A, Dechsalukthorn F (2012) Nanoparticles: a review of particle toxicology following inhalation exposure. *Inhal Toxicol* **24**, 125–35. [[Medline](#)] [[CrossRef](#)]
  - 8) Garabrant DH, Fine LJ, Oliver C, Bernstein L, Peters JM (1987) Abnormalities of pulmonary function and pleural disease among titanium metal production workers. *Scand J Work Environ Health* **13**, 47–51. [[Medline](#)] [[CrossRef](#)]
  - 9) Chen JL, Fayerweather WE (1988) Epidemiologic study of workers exposed to titanium dioxide. *J Occup Med* **30**, 937–42. [[Medline](#)] [[CrossRef](#)]
  - 10) Chen HW, Su SF, Chien CT, Lin WH, Yu SL, Chou CC, Chen JJ, Yang PC (2006) Titanium dioxide nanoparticles induce emphysema-like lung injury in mice. *FASEB J* **20**, 2393–5. [[Medline](#)] [[CrossRef](#)]
  - 11) Iavicoli I, Leso V, Fontana L, Bergamaschi A (2011) Toxicological effects of titanium dioxide nanoparticles: a review of in vitro mammalian studies. *Eur Rev Med Pharmacol Sci* **15**, 481–508. [[Medline](#)]
  - 12) Li B, Ze Y, Sun Q, Zhang T, Sang X, Cui Y, Wang X, Gui S, Tan D, Zhu M, Zhao X, Sheng L, Wang L, Hong F, Tang M (2013) Molecular mechanisms of nanosized titanium dioxide-induced pulmonary injury in mice. *PLoS ONE* **8**, e55563. [[Medline](#)] [[CrossRef](#)]
  - 13) Gurr JR, Wang AS, Chen CH, Jan KY (2005) Ultrafine titanium dioxide particles in the absence of photoactivation can induce oxidative damage to human bronchial epithelial cells. *Toxicology* **213**, 66–73. [[Medline](#)] [[CrossRef](#)]
  - 14) Nemmar A, Melghit K, Ali BH (2008) The acute proinflammatory and prothrombotic effects of pulmonary exposure to rutile TiO₂ nanorods in rats. *Exp Biol Med* (Maywood) **233**, 610–9. [[Medline](#)] [[CrossRef](#)]
  - 15) Warheit DB, Webb TR, Sayes CM, Colvin VL, Reed KL (2006) Pulmonary instillation studies with nanoscale TiO₂ rods and dots in rats: toxicity is not dependent upon particle size and surface area. *Toxicol Sci* **91**, 227–36. [[Medline](#)] [[CrossRef](#)]
  - 16) Gustafsson Å, Lindstedt E, Elfsmark LS, Bucht A (2011) Lung exposure of titanium dioxide nanoparticles induces innate immune activation and long-lasting lymphocyte response in the Dark Agouti rat. *J Immunotoxicol* **8**, 111–21. [[Medline](#)] [[CrossRef](#)]
  - 17) Hussain S, Vanoirbeek JA, Luyts K, De Vooght V, Verbeken E, Thomassen LC, Martens JA, Dinsdale D, Boland S, Marano F, Nemery B, Hoet PH (2011) Lung exposure to nanoparticles modulates an asthmatic response in a mouse model. *Eur Respir J* **37**, 299–309. [[Medline](#)] [[CrossRef](#)]
  - 18) Chen J, Dong X, Zhao J, Tang G (2009) In vivo acute toxicity of titanium dioxide nanoparticles to mice after intraperitoneal injection. *J Appl Toxicol* **29**, 330–7. [[Medline](#)] [[CrossRef](#)]
  - 19) Oberdörster G, Oberdörster E, Oberdörster J (2005) Nanotoxicology: an emerging discipline evolving from studies of ultrafine particles. *Environ Health Perspect* **113**, 823–39. [[Medline](#)] [[CrossRef](#)]
  - 20) Kamaruzaman NA, Kardia E, Kamaldin N, Latahir AZ, Yahaya BH (2013) The rabbit as a model for studying lung disease and stem cell therapy. *Biomed Res Int* **2013**, 691830. [[Medline](#)] [[CrossRef](#)]
  - 21) Persson CG (2002) Con: mice are not a good model of human airway disease. *Am J Respir Crit Care Med* **166**, 6–7, discussion 8. [[Medline](#)] [[CrossRef](#)]
  - 22) Bermudez E, Mangum JB, Wong BA, Asgharian B, Hext PM, Warheit DB, Everitt JI (2004) Pulmonary responses of mice, rats, and hamsters to subchronic inhalation of ultrafine titanium dioxide particles. *Toxicol Sci* **77**, 347–57. [[Medline](#)] [[CrossRef](#)]
  - 23) Zhang Y, Tao J, He P, Tang Y, Wang Y (2009) [Bio-effects of nano-TiO₂ on lungs of mice]. *Sheng Wu Yi Xue Gong Cheng Xue Za Zhi* **26**, 803–6 (in Chinese). [[Medline](#)]
  - 24) Warheit DB, Yuen IS, Kelly DP, Snajdr S, Hartsky MA (1996) Subchronic inhalation of high concentrations of low toxicity, low solubility particulates produces sustained pulmonary inflammation and cellular proliferation. *Toxicol Lett* **88**, 249–53. [[Medline](#)] [[CrossRef](#)]
  - 25) Warheit DB, Hansen JF, Yuen IS, Kelly DP, Snajdr SI, Hartsky MA (1997) Inhalation of high concentrations of low toxicity dusts in rats results in impaired pulmonary clearance mechanisms and persistent inflammation. *Toxicol Appl Pharmacol* **145**, 10–22. [[Medline](#)] [[CrossRef](#)]
  - 26) Rossi EM, Pytkäinen L, Koivisto AJ, Vippola M, Jensen KA, Miettinen M, Sirola K, Nykäsenoja H, Karisola P, Stjernvall T, Vanhala E, Kiilunen M, Pasanen P, Mäkinen M, Hämeri K, Joutsensaari J, Tuomi T, Jokiniemi J, Wolff H, Savolainen K, Matikainen S, Alenius H (2010) Airway exposure to silica-coated TiO₂ nanoparticles induces pulmonary neutrophilia in mice. *Toxicol Sci* **113**, 422–33. [[Medline](#)] [[CrossRef](#)]
  - 27) Ma-Hock L, Burkhardt S, Strauss V, Gamer AO, Wiench K, van Ravenzwaay B, Landsiedel R (2009) Development of a short-term inhalation test in the rat using nano-titanium dioxide as a model substance. *Inhal Toxicol* **21**, 102–18. [[Medline](#)] [[CrossRef](#)]
  - 28) Sager TM, Kommineni C, Castranova V (2008) Pulmonary response to intratracheal instillation of ultrafine versus fine titanium dioxide: role of particle surface area. *Part Fibre Toxicol* **5**, 17. [[Medline](#)] [[CrossRef](#)]
  - 29) Singh S, Shi T, Duffin R, Albrecht C, van Berlo D, Höhr D,

- Fubini B, Martra G, Fenoglio I, Borm PJ, Schins RP (2007) Endocytosis, oxidative stress and IL-8 expression in human lung epithelial cells upon treatment with fine and ultrafine TiO₂: role of the specific surface area and of surface methylation of the particles. *Toxicol Appl Pharmacol* **222**, 141–51. [[Medline](#)] [[CrossRef](#)]
- 30) Warheit DB, Brock WJ, Lee KP, Webb TR, Reed KL (2005) Comparative pulmonary toxicity inhalation and instillation studies with different TiO₂ particle formulations: impact of surface treatments on particle toxicity. *Toxicol Sci* **88**, 514–24. [[Medline](#)] [[CrossRef](#)]
- 31) Kobayashi N, Naya M, Endoh S, Maru J, Yamamoto K, Nakanishi J (2009) Comparative pulmonary toxicity study of nano-TiO₂ particles of different sizes and agglomerations in rats: different short- and long-term post-instillation results. *Toxicology* **264**, 110–8. [[Medline](#)] [[CrossRef](#)]
- 32) Park EJ, Yoon J, Choi K, Yi J, Park K (2009) Induction of chronic inflammation in mice treated with titanium dioxide nanoparticles by intratracheal instillation. *Toxicology* **260**, 37–46. [[Medline](#)] [[CrossRef](#)]
- 33) Larsen ST, Roursgaard M, Jensen KA, Nielsen GD (2010) Nano titanium dioxide particles promote allergic sensitization and lung inflammation in mice. *Basic Clin Pharmacol Toxicol* **106**, 114–7. [[Medline](#)] [[CrossRef](#)]
- 34) Akuthota P, Xenakis JJ, Weller PF (2011) Eosinophils: offenders or general bystanders in allergic airway disease and pulmonary immunity? *J Innate Immun* **3**, 113–9. [[Medline](#)] [[CrossRef](#)]
- 35) Barnes PJ (2011) Pathophysiology of allergic inflammation. *Immunol Rev* **242**, 31–50. [[Medline](#)] [[CrossRef](#)]