Relations of biomarkers of manganese exposure and neuropsychological effects among welders and ferroalloy smelters

Hamid HASSANI^{1,2}, Farideh GOLBABAEI^{1*}, Hamid SHIRKHANLOO^{2,3} and Mehdi TEHRANI-DOUST⁴

¹Department of Occupational Health Engineering, School of Public Health, Tehran University of Medical Sciences, Iran

²Iranian Petroleum Industry Occupational and Environmental Health Research Center (IPIOEHRC), Iranian Petroleum Industry Health Research Institute (IPIHRI), Iran

³Research Institute of Petroleum Industry (RIPI), Iran

⁴Department of Psychiatry, School of Medicine, Tehran University of Medical Sciences, Iran

Received December 4, 2014 and accepted August 24, 2015

Published online in J-STAGE September 30, 2015

Abstract: The objective of present study was to assess relationship between biomarkers of Manganese (Mn) and neuropsychological effects. The study was carried out on 27 welders and 31 ferroalloy smelters as Mn-exposed groups and 30 office workers as unexposed controls. Air Mn concentrations were determined according to NIOSH method 7300. The biological samples were prepared using microwave assisted acid digestion and all samples were analyzed by graphite furnace- atomic absorption spectroscopy (GF-AAS) in order to determine manganese. Questionnaire 16 (Q16) and Cambridge Neuropsychological Test Automated Battery (CANTAB) were used to evaluate the neuropsychological effects. The mean concentrations of air Mn for the welder and ferroalloy smelter groups were $0.023 \pm 0.012 \text{ mg/m}^3$ and $0.008 \pm 0.005 \text{ mg/m}^3$, respectively. Manganese concentrations in blood, urine, and toenail samples of exposed workers ranged between 1.80–32.60 ($\mu\text{g/l}$), 1.00–42.50 ($\mu\text{g/l}$), and 0.10–6.08 ($\mu\text{g/g}$), respectively. Mean Mn concentrations in all biological samples of cases were significantly higher than unexposed controls ($p < 0.05$). A moderate relationship was observed between biomarkers of Mn exposure, air Mn, Q16 as well as some neurocognitive outcome measures. The present study shows that blood Mn, urine Mn and toenail Mn could be used to distinguish Mn-exposed workers from unexposed population at the group level.

Key words: Biomarkers of manganese exposure, Blood, Neuropsychological effects, Toenail, Urine, Welding and smelting processes

Introduction

Manganese (Mn) is an essential element for many

biological functions in human being^{1–3}). However, over-exposure to Mn can cause health problems especially neurotoxicity⁴). Manganese accumulates in brain regions and it can cause central nervous system (CNS) abnormalities as well as neuropsychiatric disorders⁵). The respiratory system is the most important way of Mn absorption for workers in the Mn-related industries. In this matter, entry

*To whom correspondence should be addressed.
E-mail: fgolbabaei@sina.tums.ac.ir

of Mn to brain via the olfactory nerve has been reported as the main pathway⁶⁾.

Occupational exposure to Mn occurs in various industrial processes such as welding, ferroalloy steel production and dry cell battery manufacturing^{7, 8)}. A significant number of studies have shown that welders are exposed to high concentrations of Mn-containing welding fumes which leads to increasing the prevalence of neurobehavioral symptoms among them^{3, 5, 7, 9–14)}. Several neurobehavioral symptoms have also been reported among the ferroalloy smelting workers exposed to Mn^{15–17)}. Ellingsen *et al.*, in a study on welders, reported that Mn exposures ranged between 0.003 to 4.26 mg/m³¹⁸⁾. Two neurological function alterations including Impaired Finger Tapping Speed and Grooved Pegboard Performance were considerable among former welders¹³⁾. Wastensson *et al.*, in a study on ship welders, showed that long-term exposure to Mn can have adverse effect on motor function³⁾. Occupational exposure to Mn can also affect other systems including respiratory, cardiac, liver and reproductive systems^{6, 7)}. A study on smelting workers by Cowan *et al.*⁸⁾ showed that airborne Mn concentration ranged between 0.01 to 0.374 mg/m³. Cowan *et al.*¹⁷⁾ have found a relationship between blood manganese-iron ratio and early onset neurobehavioral alterations. Bouchard *et al.*¹⁶⁾ have shown an exposure-effect relation for symptoms related to memory, concentration and balance in a group of Mn-alloy workers.

Biological monitoring (or biomonitoring) is a useful tool for occupational exposure assessment and potential related health risks¹⁹⁾. Biomonitoring provides valuable information that can be used to prevent adverse health effects. Having studied potential biomarkers of Mn including Mn in blood, urine and hair, Bader *et al.*²⁰⁾ reported that blood Mn is better than urine and/or hair Mn for assessment of Mn exposure. In their study, the mean concentration of blood Mn was 12.2 ± 4.8 µg/l (ranged: 3.2–25.8 µg/l). Similarly, Wongwit *et al.*²¹⁾ suggested the blood Mn as a recommended biomarker in occupational studies. However, Crossgrove and Zheng⁶⁾ noted that blood Mn indicates recent exposure, thus, it can't be used to assess long-term effects. The number of studies on biomarkers of Mn exposure has been increased during recent two decades, but suitable biomarkers are not well determined.

The objectives of this study were: 1) to assess the correlation between air Mn, blood Mn, urine Mn, and toenail Mn concentrations, 2) to evaluate the neurobehavioral symptoms, and 3) to examine the relations of biomarkers of Mn exposure with neurobehavioral and neurocognitive outcome measures among welders and ferroalloy smelters

as two occupationally Mn-exposed groups.

Subjects and Methods

Subjects

This cross-sectional study was carried out in an automotive parts manufacturing industry in Tehran, Iran, in 2013. Workers who had more than 1 yr of work history were recruited for the study. The exclusion criteria were similar to previous studies^{12, 13, 15)}, including: alcohol consumption, drug abuse, diabetes mellitus, damages of central nervous system (CNS) such as concussion, prior exposure to organic solvents and neurotoxic metals (e.g., lead and mercury), and serious kidney or liver diseases. Based on 95% confidence intervals and 80% power of statistical test, the number of calculated sample (sample size) was at least 27 for each group. Therefore, decision was made to recruit about 30 workers for each study group. 30 welders, 31 ferroalloy smelters and 30 office workers were invited to participate in the study. One welder refused and 2 others were excluded from this study due to kidney disease records. After exclusions, 27 welders and 31 ferroalloy smelters as exposed groups (cases) and 30 office workers as unexposed control group were involved in the study (participation rate: 96.70%). It should be noted that the unexposed controls were selected from matched people of the same sex (male), age, work history, and education level without any diseases affecting. In this study, the accuracy of participants' answers relating to alcohol consumption and drug abuse was based on two factors: firstly, all participants were volunteers who were informed about objectives of study, importance of their answers, and ethical issues. Informed consents were obtained from participants. The study was approved by ethics committee at Tehran University of Medical Sciences (TUMS). Secondly, there were no medical history of related items in medical records of the studied participants.

In the studied industry, workers were working 44 h per wk. The average time of welding was about 4 h per d during a work shift.

Most exposed workers (77.59%) reported that they usually uses respiratory protection masks (Type: FFP3/EN149:2001 series).

Work description

Welding type

The type of welding was Shielded Metal Arc Welding (SMAW) and two types of electrode were used (E 6013 and E 7018). The percentage of Mn in these electrodes

was about 0.50 to 0.90. Welders were exposed to Mn-containing welding fumes through inhalation during welding processes. Welding fumes contains metals especially iron, manganese (Mn), chromium, and nickel⁷⁾. Among these metals, Mn is a well-known neurotoxic metal.

Smelting process

Eight furnaces were operated for smelting process in the studied industry (producing grey iron and ductile iron automotive parts). The percentage of Mn in the molten materials was about 0.30 to 0.50.

Iron, manganese (Mn), nickel, graphite, and silicon were commonly used in the studied industry. Like welding fumes, smelting fumes also contains Mn. So, Smelters were exposed to Mn-containing smelting fumes through inhalation during smelting processes.

Collection of air samples and analysis

Personal air samples were collected using mixed cellulose ester membrane filters (0.8 mm pore size, 25 mm diameter; SKC Corp) and SKC pumps (224-PCMTX8, 224-PCMTX8, S/No. 09523488- SKC Ltd. DT11 8ST. UK). The flow rate of pumps was adjusted at 2.0 l/min for sampling. All pumps were calibrated before and after use and the mean of flow rate was considered in calculations. Analysis was done using graphite furnace-atomic absorption spectroscopy (GF-AAS) according to NIOSH analytical method 7300²²⁾.

Biological monitoring

Biological samples including blood, urine and toenail samples were collected at the end of work shift. The time lag between collecting air samples and biological samples was about 7 d. It should be noted that all of the biological samples were collected at the same working day.

Five milliliter (ml) of venous blood was collected from each worker. Blood samples were divided into two tubes (4 ml in a heparinized tube and 1 ml in a non-heparinized tube). Urine samples were collected in polyethylene tubes and for each sample 1ml nitric acid was added per 10 ml urine sample. Toenails were from all toes of participants. Superficial contamination was removed by sonicating toenail samples in 10 ml 1% Tritone X-100 solution for 20 min and then washed with deionized water and they were dried before weighting.

Digestion method for biological samples

The biological samples were prepared using microwave assisted acid digestion (Microwave digestion- EPA

Method 3052 on the Multiwave 3000). In this clean and reproducible closed digestion method, the quality assurance/quality control (QA/QC) has been improved. Another benefit of microwave assisted acid digestion method is reduced preparation time²³⁾. Nitric acid, hydrochloric acid, and hydrogen peroxide were used as reagents. After adding reagents in the reaction vessels with the biological samples, vessels were placed in the microwave.

Analysis of biological samples

The graphite furnace-atomic absorption spectroscopy (GF-AAS) was used for analysis of biological samples. The validity of analytical method was examined by using Standard Reference Material (SRM 2670a, NIST). Data showed that the recovery percentage was 96.2% (SRM: $2.6 \pm 0.7 \mu\text{g/l}$, value obtained: $2.5 \pm 0.4 \mu\text{g/l}$). The Limit of Quantitation (LOQ), Limit of Detection (LOD) and linear range for Mn analysis were $1.0 \mu\text{g/l}$, $0.25 \mu\text{g/l}$, and $1\text{--}35 \mu\text{g/l}$, respectively.

Neurobehavioral symptoms

In this study, the questionnaire 16 (Q16)²⁴⁾ was used for assessing neurobehavioral symptoms. This questionnaire contains 16 Yes/No questions about abnormal tiredly, painful tingling in body, palpitation, irritation and depression without particular reason, problem with concentrating, having a short memory, perspire without particular reason, problems with buttoning and unbuttoning, difficulties to get the meaning from reading newspapers or books, oppression of chest, memory problems, headache (at least once a week), and being less interested in sex. The frequency of neurobehavioral symptoms of welders and ferroalloy smelters was compared to unexposed office workers.

Neurocognitive tests

Neurocognitive tests including choice reaction time (CRT) as an attention test, spatial working memory (SWM) as an executive function test, paired associates learning (PAL) as a visual memory test, and stocking of Cambridge (SOC) as a planning test were performed using Cambridge Neuropsychological Test Automated Battery (CANTAB)²⁵⁾. This computerized cognitive assessment tool would improve the quality of study data, providing greater sensitivity with maximization of standardization of administration and scoring as well as the precision of measurement compared to manually-administered (paper and pencil) test batteries used in previous studies.

Table 1. The demographic Information of the study groups

Groups	n	Age (yr) (Mean $\pm$ SD)	Work history (yr) (Mean $\pm$ SD)	BMI (kg/m ²) (Mean $\pm$ SD)	Number of smokers (%)	Number of married people (%)
Welder	27	31.18 $\pm$ 4.06	7.04 $\pm$ 4.64	24.47 $\pm$ 2.92	7 (25.92)	24 (88.88)
Control	30	34.45 $\pm$ 7.11	8.61 $\pm$ 4.70	26.11 $\pm$ 4.72	4 (13.33)	26 (86.66)
<i>p</i> value	-	0.053	0.228	0.149	0.248	0.597
Ferroalloy smelter	31	34.88 $\pm$ 5.63	9.15 $\pm$ 4.38	27.06 $\pm$ 4.15	7 (22.58)	28 (90.32)
Control	30	34.45 $\pm$ 7.11	8.61 $\pm$ 4.70	26.11 $\pm$ 4.72	4 (13.33)	26 (86.66)
<i>p</i> value	-	0.789	0.638	0.404	0.384	0.155

*Differences between case and control groups, Student's *t*-test, *p*<0.05; statistically significant

Table 2. Mn concentrations in air, blood, urine, and toenail samples

Groups	Sample (unit)	n	Mean $\pm$ SD	Range	<i>p</i> value
Welder	Air (mg/m ³)	27	0.023 $\pm$ 0.012	0.002–0.062	-
	Blood (μ g/l)	27	15.85 $\pm$ 7.11	1.80–32.60	0.007
	Urine (μ g/l)	27	13.00 $\pm$ 7.98	1.00–24.10	0.026
	Toenail (μ g/g)	24	3.08 $\pm$ 1.44	1.15–6.08	<0.001
Ferroalloy smelter	Air (mg/m ³)	31	0.008 $\pm$ 0.005	0.002–0.029	-
	Blood (μ g/l)	31	17.33 $\pm$ 8.66	4.00–32.06	0.001
	Urine (μ g/l)	31	12.56 $\pm$ 10.00	1.00–42.50	0.047
	Toenail (μ g/g)	31	1.52 $\pm$ 0.95	0.10–3.70	0.048
Case (Welder and Ferroalloy smelter)	Air (mg/m ³)	58	0.014 $\pm$ 0.011	0.002–0.062	-
	Blood (μ g/l)	58	16.74 $\pm$ 8.04	1.80–32.60	<0.001
	Urine (μ g/l)	58	12.73 $\pm$ 9.17	1.00–42.50	0.018
	Toenail (μ g/g)	55	2.14 $\pm$ 1.39	0.10–6.08	0.001
Control	Blood (μ g/l)	30	9.37 $\pm$ 8.70	1.00–33.40	-
	Urine (μ g/l)	30	7.86 $\pm$ 7.82	1.00–27.80	-
	Toenail (μ g/g)	28	1.08 $\pm$ 1.02	0.10–4.47	-

*Comparison with control, Student's *t*-test, *p*<0.05; statistically significant

Statistical analyses

Data were analyzed using descriptive statistics, compare means, Mann-Whitney test, and partial correlation tests. Student's *t*-test was used to determine differences among means for some characteristics as well as Mn concentrations in air, blood, urine, and toenail samples of case and control groups. Comparisons between medians of Q16 (Number of positive answers) of studied groups were performed using Mann-Whitney test. Partial correlation tests were carried out to determine the correlation among air, blood, urine and toenail samples, neurobehavioral symptoms (Q16 score) as well as neurocognitive outcome measures in the exposed groups. We used the statistical package for the social sciences software (SPSS version 17) for all statistical analyses. Results were considered statistically significant at *p*<0.05.

Results

The demographic information of the case and control groups is shown in Table 1. There were no significant differences between the case and control groups in terms of age, work history, body mass index (BMI), smoking, and marriage status.

The Mn concentrations in air, blood, urine, and toenail samples of our study population are shown in Table 2. The mean concentrations of air Mn for the welder and ferroalloy smelter groups were 0.023 $\pm$ 0.012 mg/m³ (range: 0.002–0.062 mg/m³) and 0.008 $\pm$ 0.005 mg/m³ (range: 0.002–0.029 mg/m³), respectively.

Manganese concentrations in blood, urine, and toenail samples of exposed workers ranged between 1.80–32.60 (μ g/l), 1.00–42.50 (μ g/l), and 0.10–6.08 (μ g/g), respectively. As it is shown in Table 2, the mean Mn concentrations in all biological samples were significantly

higher in the welders and ferroalloy smelters than in the unexposed controls ($p<0.05$).

The number of self-reported neurobehavioral symptoms was significantly more prevalent among welders and ferroalloy smelters than among unexposed controls ($p<0.001$), (Table 3).

The results of partial correlation showed that there were significant relationships between blood Mn and toenail Mn with Q16 score, and urine Mn with toenail Mn in the welder group. Correlations between blood Mn with urine Mn, toenail Mn, and air Mn, as well as urine Mn with toenail Mn and air Mn were statistically significant in the ferroalloy smelter group (Table 4).

Table 5 shows significant inverse relationships between neurocognitive outcome measures and biomarkers of Mn exposure. The correlation coefficients ranged from 0.299 to 0.566, ($p<0.05$).

Discussion

The results of this study showed that the concentrations of Mn in biological samples (blood, urine, and toenail) of both welder and ferroalloy smelter groups were significantly higher than the unexposed-control group. Welders and ferroalloy smelters reported more neurobehavioral symptoms than the controls. Significant relationships were also observed between blood Mn and toenail Mn with the number of neurobehavioral symptoms in the welder group. We found significant inverse relationships between some neurocognitive outcome measures and biomarkers of Mn exposure.

The results of previous studies on biomarkers of Mn exposure are contradictory, thus sufficient data, well quantitative and qualitative studies, and review of literature are needed to determine reliable biological indicators. Some previous studies showed that the blood Mn and urine Mn levels of exposed workers were significantly higher than unexposed controls^{2, 4, 21, 26}. Consistently, our results showed that blood Mn and urine Mn can be utilized for distinguishing exposed from unexposed workers at the group based level. Bader *et al.*²⁰ examined the relationship between the Mn dioxide (MnO_2) exposure and three potential biomarkers including blood, urine and axillary hair in a study on dry cell battery manufacturers. They concluded that blood Mn is better indicator than urine and hair Mn for the biomonitoring of MnO_2 exposure. Similarly, blood Mn is recommended by Wongwit *et al.*²¹ for the biomonitoring of Mn exposure among occupationally-exposed workers. In Laohaudomchok *et al.*²⁷ study,

Table 3. Neurobehavioral symptoms in the exposed and unexposed workers

Groups	n	Q16, No of positive answers		p value
		Median	Range	
Welder	27	3	1–10	<0/001
Ferroalloy smelter	31	4	1–11	<0/001
Control	30	1	0–6	-

*Comparison with control, Mann-Whitney test, $p<0.05$; statistically significant

toenail Mn was significantly correlated with cumulative Mn exposure indices; however, there were no significant correlation between blood Mn and urine Mn with cumulative Mn exposure indices. According to this finding, toenail Mn could be used as a suitable biomarker for long-term exposure assessment. This shows that the toenail Mn might be considered as a biomarker of exposure and effect. Similar to Laohaudomchok *et al.*²⁷ study, our results showed that toenail Mn seems to be a useful biomarker of exposure and effect. In general, blood Mn and urine Mn could be used as two practical biomarkers in the group based studies. However, toenail Mn is the preferred biomarker for long-term exposure assessment.

Despite the difference in air Mn levels between the two exposed groups, the blood Mn and urine Mn levels were similar among the welders and ferroalloy smelters. One reason of this observation was that the individual air Mn measurement for just one day may not represent actual individual exposure. Also, it is worth noting that half-life of Mn varies in blood, urine, and toenail (hours or several days in blood and urine, and several months in toenail). So, we suggest to consider the cumulative exposure index (CEI) for future studies.

Although mean air Mn concentrations for both of our studied groups were below the Occupational Exposure Limit -Time Weighted Average (OEL-TWA; 0.2 mg/m^3) recommended by Environmental and Occupational Health Center of the Ministry of Health and Medical Education of Iran²⁸, the number of self-reported neurobehavioral symptoms was significantly higher in the Mn-exposed workers. Thus, there is a need to modify the current OEL-TWA for Mn to prevent the occurrence of related adverse effects especially neurotoxicity. In other words, a revised OEL-TWA for Mn (lower than 0.2 mg/m^3) can protect nearly all industrial workers.

“The questionnaire 16 (Q16) is commonly used to study prevalence of neurotoxic symptoms among workers exposed to organic solvents.”²⁴. However, a few studies

Table 4. Partial correlation* of Mn concentrations among air, blood, urine, and toenail samples and associations with neurobehavioral symptoms (Q16 score) in the exposed groups

Groups	Variable	Blood		Urine		Toenail		Air	
		r	p value	r	p value	r	p value	r	p value
Welder	Urine	0.507	0.016	-	-	-	-	-	-
	Toenail	0.209	0.376	0.541	0.014	-	-	-	-
	Air	0.332	0.131	0.218	0.329	0.397	0.083	-	-
	Q16	0.437	0.042	0.335	0.128	0.646	0.002	0.224	0.304
Ferroalloy smelter	Urine	0.679	<0.001	-	-	-	-	-	-
	Toenail	0.646	<0.001	0.649	<0.001	-	-	-	-
	Air	0.377	0.030	0.407	0.019	0.304	0.097	-	-
	Q16	0.087	0.643	0.046	0.801	0.200	0.280	0.302	0.083

*Adjusted for age, work history and smoking. $p < 0.05$; statistically significant

Table 5. Significant correlations* between neurocognitive outcome measures and biomarkers of manganese exposure in the exposed groups

Groups	Variables	r	p value
Welder	PAL Stages completed on first trial & Mn- Urine	-0.546	0.019
	PAL Total error 6 shape & Mn- Toenail	-0.470	0.049
Ferroalloy smelter	SWM Double error & Mn- Blood	-0.426	0.038
	SWM Double error & Mn- Urine	-0.566	0.004
	SWM Double error & Mn- Toenail	-0.503	0.012
	SWM Double error (8 boxes) & Mn- Urine	-0.422	0.040
	SWM Within errors & Mn- Blood	-0.425	0.038
	SWM Within errors & Mn- Urine	-0.512	0.010
Case (Welder and Ferroalloy smelter)	PAL Stages completed on first trial & Mn- Urine	-0.343	0.023
	SOC Mean subsequent thinking time (4 moves) & Mn- Blood	-0.382	0.011
	SOC Mean subsequent thinking time (4 moves) & Mn- Urine	-0.299	0.048
	SWM Double error & Mn- Blood	-0.407	0.006
	SWM Double error & Mn- Urine	-0.376	0.012
	SWM Double error (8 boxes) & Mn- Blood	-0.300	0.048
	SWM Double error (8 boxes) & Mn- Urine	-0.318	0.035
	SWM Within errors & Mn- Blood	-0.445	0.002
	SWM Within errors & Mn- Urine	-0.435	0.003
	SWM Within error (6 boxes) & Mn- Blood	-0.299	0.049

*Adjusted for age, work history and smoking. $p < 0.05$; statistically significant

used the Q16 for investigating neurobehavioral symptoms among welders^{13, 29, 30}). In a study on welders by Ellingsen *et al.*¹³), the arithmetic mean of Q16 positive answers was 3.6 (range: 1–13). In our study, the median of positive answers for the welder and ferroalloy smelter groups were 3 (range: 1–10) and 4 (range: 1–11), respectively. Generally, Mn-exposed groups reported more neurobehavioral symptoms than unexposed controls.

The correlation test showed statistically significant associations between biomarkers of Mn exposure, air Mn and Q16. Correlation coefficients ranged between 0.377–0.679 (moderate relationship) for observed significant associa-

tions. As a result, a direct relationship between the level of Mn exposure and the number of neurotoxic symptoms was found.

Laohaudomchok *et al.*³¹) suggested that even a low-level Mn exposure may have neuropsychological effects on welders. In their study, welders' cumulative Mn exposure index (Mn-CEI) was significantly associated with slower reaction time as well as worse mood. Similarly, Wastensson *et al.*³) showed that the poorer performance in the grooved pegboard test, as a neuromotor function test, was associated with Mn-CEI among welders. Using the computerized cognitive assessment tool "CANTAB"

for neurocognitive tests in the current study, the accuracy and precision of our neurocognitive results seems to be more reliable compared to those previous studies that have used manually-administered test batteries. Our results showed that biomarkers of Mn exposure were significantly correlated with some neurocognitive outcome measures. An increase in level of Mn in biological samples results in a decrease in neurocognitive performances (negative relationship). However, similar to many field studies, the major limitation of this cross-sectional study was small sample size. Therefore, we suggest a longitudinal study on large samples for further research. A potential limitation of this study is the lack of information related to dietary patterns among workers. In a similar study, Laohaudomchok *et al.*²⁷⁾ have noted that “Mn from food intake may not make a significant contribution to internal dose”. However, we collected and analyzed ten drinking water samples to control Mn levels in drinking water as a potential confounding factor. The results showed that the mean concentration of Mn in drinking water samples was $22.1 \pm 0.9 \mu\text{g/l}$ which was lower than acceptable concentration for Mn in drinking water ($50 \mu\text{g/l}$). Therefore, it couldn't have significant effect on the obtained results. Another potential limitation of the present study seems to be the lack of data related to ventilation system efficiency. In general, one advantage of this case-control study is comparing the different biomarkers of Mn exposure among two Mn-exposed groups. Significant relations of biomarkers of Mn exposure with neuropsychological effects indicates adverse effects among welder and smelter groups.

The current Occupational Exposure Limit-Time Weighted Average of Mn (OEL-TWA; 0.2 mg/m^3) seems not providing sufficient protection against the neuropsychological effects. So, a change is needed to modify the current OEL-TWA of Mn for protecting exposed workers. The present study showed that blood Mn, urine Mn and toenail Mn could be used to distinguish Mn-exposed workers from unexposed population at the group level. Further studies are needed to identify reliable biomarkers for assessing Mn exposure at the individual level as well as to determine suitable biomarkers and biological exposure indices (BEIs) for Mn with concerning neuropsychological effects.

Acknowledgements

We would like to thanks Mr. Majid Minaei, Mr. Ahmad Mirzaei, Ms. Azar Mohammadzadeh, Mrs. Somayeh Farhang Dehghan and Mrs. Sarvar Charehdar for their kind assistance during this research. This study was supported

by Tehran University of Medical Sciences.

References

- 1) Zheng W, Fu SX, Dydak U, Cowan DM (2011) Biomarkers of manganese intoxication. *Neurotoxicology* **32**, 1–8. [[Medline](#)] [[CrossRef](#)]
- 2) Järvisalo J, Olkinuora M, Kiilunen M, Kivistö H, Ristola P, Tossavainen A, Aitio A (1992) Urinary and blood manganese in occupationally nonexposed populations and in manual metal arc welders of mild steel. *Int Arch Occup Environ Health* **63**, 495–501. [[Medline](#)] [[CrossRef](#)]
- 3) Wastensson G, Sallsten G, Bast-Pettersen R, Barregard L (2012) Neuromotor function in ship welders after cessation of manganese exposure. *Int Arch Occup Environ Health* **85**, 703–13. [[Medline](#)] [[CrossRef](#)]
- 4) Ellingsen DG, Hetland SM, Thomassen Y (2003) Manganese air exposure assessment and biological monitoring in the manganese alloy production industry. *J Environ Monit* **5**, 84–90. [[Medline](#)] [[CrossRef](#)]
- 5) Antonini JM, Santamaria AB, Jenkins NT, Albin E, Lucchini R (2006) Fate of manganese associated with the inhalation of welding fumes: potential neurological effects. *Neurotoxicology* **27**, 304–10. [[Medline](#)] [[CrossRef](#)]
- 6) Crossgrove J, Zheng W (2004) Manganese toxicity upon overexposure. *NMR Biomed* **17**, 544–53. [[Medline](#)] [[CrossRef](#)]
- 7) Hassani H, Golbabaie F, Ghahri A, Hosseini M, Shirkhanloo H, Dinari B, Eskandari D, Fallahi M (2012) Occupational exposure to manganese-containing welding fumes and pulmonary function indices among natural gas transmission pipeline welders. *J Occup Health* **54**, 316–22. [[Medline](#)] [[CrossRef](#)]
- 8) Cowan DM, Fan Q, Zou Y, Shi X, Chen J, Aschner M, Rosenthal FS, Zheng W (2009) Manganese exposure among smelting workers: blood manganese-iron ratio as a novel tool for manganese exposure assessment. *Biomarkers* **14**, 3–16. [[Medline](#)] [[CrossRef](#)]
- 9) Hobson A, Seixas N, Sterling D, Racette BA (2011) Estimation of particulate mass and manganese exposure levels among welders. *Ann Occup Hyg* **55**, 113–25. [[Medline](#)] [[CrossRef](#)]
- 10) Laohaudomchok W, Cavallari JM, Fang SC, Lin X, Herrick RF, Christiani DC, Weisskopf MG (2010) Assessment of occupational exposure to manganese and other metals in welding fumes by portable X-ray fluorescence spectrometer. *J Occup Environ Hyg* **7**, 456–65. [[Medline](#)] [[CrossRef](#)]
- 11) Richman JD, Livi KJ, Geyh AS (2011) A scanning transmission electron microscopy method for determining manganese composition in welding fume as a function of primary particle size. *J Aerosol Sci* **42**, 408–18. [[Medline](#)] [[CrossRef](#)]
- 12) Wang X, Yang Y, Wang X, Xu S (2006) The effect of occupational exposure to metals on the nervous system

- function in welders. *J Occup Health* **48**, 100–6. [[Medline](#)] [[CrossRef](#)]
- 13) Ellingsen DG, Konstantinov R, Bast-Pettersen R, Merkurjeva L, Chashchin M, Thomassen Y, Chashchin V (2008) A neurobehavioral study of current and former welders exposed to manganese. *Neurotoxicology* **29**, 48–59. [[Medline](#)] [[CrossRef](#)]
 - 14) Sriram K, Lin GX, Jefferson AM, Roberts JR, Chapman RS, Chen BT, Soukup JM, Ghio AJ, Antonini JM (2010) Dopaminergic neurotoxicity following pulmonary exposure to manganese-containing welding fumes. *Arch Toxicol* **84**, 521–40. [[Medline](#)] [[CrossRef](#)]
 - 15) Bouchard M, Mergler D, Baldwin M, Panisset M, Roels HA (2007) Neuropsychiatric symptoms and past manganese exposure in a ferro-alloy plant. *Neurotoxicology* **28**, 290–7. [[Medline](#)] [[CrossRef](#)]
 - 16) Bouchard M, Mergler D, Baldwin ME, Panisset M (2008) Manganese cumulative exposure and symptoms: a follow-up study of alloy workers. *Neurotoxicology* **29**, 577–83. [[Medline](#)] [[CrossRef](#)]
 - 17) Cowan DM, Zheng W, Zou Y, Shi X, Chen J, Rosenthal FS, Fan Q (2009) Manganese exposure among smelting workers: relationship between blood manganese-iron ratio and early onset neurobehavioral alterations. *Neurotoxicology* **30**, 1214–22. [[Medline](#)] [[CrossRef](#)]
 - 18) Flynn MR, Susi P (2009) Neurological risks associated with manganese exposure from welding operations—a literature review. *Int J Hyg Environ Health* **212**, 459–69. [[Medline](#)] [[CrossRef](#)]
 - 19) Ong CN (1999) Reference values and action levels of biological monitoring in occupational exposure. *Toxicol Lett* **108**, 127–35. [[Medline](#)] [[CrossRef](#)]
 - 20) Bader M, Dietz MC, Ihrig A, Triebig G (1999) Biomonitoring of manganese in blood, urine and axillary hair following low-dose exposure during the manufacture of dry cell batteries. *Int Arch Occup Environ Health* **72**, 521–7. [[Medline](#)] [[CrossRef](#)]
 - 21) Wongwit W, Kaewkungwal J, Chantachum Y, Visemanee V (2004) Comparison of biological specimens for manganese determination among highly exposed welders. *Southeast Asian J Trop Med Public Health* **35**, 764–9. [[Medline](#)]
 - 22) National Institute for Occupational Safety and Health (NIOSH) (2003) Manual of analytical methods, method 7300, 4th Ed. NIOSH, Cincinnati.
 - 23) Mangum SJ (2009) Microwave digestion- EPA Method 3052 on the Multiwave 3000. Field application report. PerkinElmer Inc., Waltham.
 - 24) Lundberg I, Högberg M, Michélsen H, Nise G, Hogstedt C (1997) Evaluation of the Q16 questionnaire on neurotoxic symptoms and a review of its use. *Occup Environ Med* **54**, 343–50. [[Medline](#)] [[CrossRef](#)]
 - 25) Cambridge Neuropsychological Test Automated Battery (CANTAB): <http://www.cambridgecognition.com/clinicaltrials/cantabsolutions/tests>.
 - 26) Apostoli P, Lucchini R, Alessio L (2000) Are current biomarkers suitable for the assessment of manganese exposure in individual workers? *Am J Ind Med* **37**, 283–90. [[Medline](#)] [[CrossRef](#)]
 - 27) Laohaudomchok W, Lin X, Herrick RF, Fang SC, Cavallari JM, Christiani DC, Weisskopf MG (2011) Toenail, blood, and urine as biomarkers of manganese exposure. *J Occup Environ Med* **53**, 506–10. [[Medline](#)] [[CrossRef](#)]
 - 28) Environmental and Occupational Health Center Ministry of Health and Medical Education, Iran (2012) Occupational Exposure Limits (OEL), last revision.
 - 29) Sjögren B, Gustavsson P, Hogstedt C (1990) Neuropsychiatric symptoms among welders exposed to neurotoxic metals. *Br J Ind Med* **47**, 704–7. [[Medline](#)]
 - 30) Zoni S, Albini E, Lucchini R (2007) Neuropsychological testing for the assessment of manganese neurotoxicity: a review and a proposal. *Am J Ind Med* **50**, 812–30. [[Medline](#)] [[CrossRef](#)]
 - 31) Laohaudomchok W, Lin X, Herrick RF, Fang SC, Cavallari JM, Shrairman R, Landau A, Christiani DC, Weisskopf MG (2011) Neuropsychological effects of low-level manganese exposure in welders. *Neurotoxicology* **32**, 171–9. [[Medline](#)] [[CrossRef](#)]