Advance Publication

INDUSTRIAL HEALTH

Received: October 13, 2020

Accepted: February 3, 2022

J-STAGE Advance Published Date: February 15, 2022

TITLE PAGE

Original Article

Title: Do holidays change subjective sleep length or sleep debt in shift work disorder?

Running Title: SLEEP LENGTH AND SLEEP DEBT AFTER HOLIDAYS IN SWD

Authors: Päivi VANTTOLA^{1*}, Mikko HÄRMÄ¹, Christer HUBLIN¹, Katriina VIITASALO², Mikael SALLINEN^{1, 3}, Jussi VIRKKALA^{1, 4} and Sampsa PUTTONEN¹

Affiliations: ¹Finnish Institute of Occupational Health, Finland

²Finnair Health Services, Finland

³Department of Psychology, University of Jyväskylä, Finland

⁴Department of Clinical Neurophysiology, University of Helsinki and Helsinki University Hospital, Finland

Correspondence: Päivi Vanttola; Finnish Institute of Occupational Health, PO Box 40, FI-00032 TYÖTERVEYSLAITOS, Finland; Tel. +358 50 570 8639; Fax: N/A; E-mail: paivi.vanttola@ttl.fi

Abstract:

In shift work disorder (SWD), disturbed sleep acutely impairs employees' recovery, but little attention has been paid to sleep during longer recovery periods. We examined how holidays affect self-estimated sleep length, sleep debt, and recovery in cases of SWD. Twenty-one shift workers with questionnaire-based SWD and nine reference cases without SWD symptoms completed a questionnaire on recovery and sleep need. They also reported sleep length on two separate occasions: during a work period and after ≥ 2 weeks of holidays. Sleep debt was calculated by subtracting sleep length from sleep need. We used parametric tests to compare the groups and the periods. The groups reported shorter sleep on workdays than during holidays (median difference: SWD group 1.7 h, $p < 0.001$; reference group 1.5 h; $p < 0.05$). The SWD group's self-estimated sleep during holidays increased less above the sleep need (median 0.0 h) than the reference group's sleep (1.0 h, $p < 0.05$). In addition, the SWD group reported good recovery from irregular working hours less often (14%) than the reference group (100%, $p < 0.001$). Although holidays were generally associated with longer sleep estimates than workdays, employees with SWD experienced consistently less efficient recovery than those without SWD.

Keywords: Circadian rhythm sleep-wake disorder, Day off, Excessive sleepiness, Free day, Insomnia, Non-workday, Shift work sleep disorder, Vacation

Introduction

Work schedules restrict shift workers' sleep on workdays. During weekly days off, individuals can time their sleep more freely and increase their sleep length to recover from work-related strain. The epidemiological literature suggests that habitual sleep of seven to eight hours is beneficial for health and that outside this range, the risk of mortality, for example, increases¹⁻³). In laboratory conditions, the length of asymptotic sleep has been 8.2—8.9 h among young adults and 7.4 h among the elderly^{4, 5}). During a longer holiday, the prior work schedule should not considerably affect the amount of sleep. De Bloom *et al.*⁶) found that employees report approximately 40-min longer night sleep during (7.4 h) than before (6.7 h) a holiday period of at least two weeks. During days off, sleep length increases by approximately 20–45 min in comparison to workdays⁷⁻⁹), and sleep quality improves⁹).

In shift work, sleep length decreases by up to four hours prior to morning shifts and between night shifts, and many shift workers tackle sleep loss with prophylactic and restorative napping¹⁰). In addition to morning and night shifts, short shift intervals typically restrict sleep length and cause sleepiness^{11, 12}), increasing the risk of shift work disorder (SWD)¹³). SWD is characterised by chronic shift work schedule-related insomnia and/or excessive sleepiness, based on the second¹⁴) and third¹⁵) editions of the International Classification of Sleep Disorders (ICSD-2 and ICSD-3, respectively). We recently concluded that approximately 6% of hospital night shift workers are affected by ICSD-3-based SWD¹⁶). The ICSD-3 requires a reduced total sleep time in addition to the primary symptoms of SWD. Night

work and morning shifts – but not days off – have been associated with reduced sleep in SWD^{17, 18)}. Even so, Kalmbach *et al.*¹⁹⁾ have found that new rotating shift workers who develop SWD report shorter sleep on weekend nights than those who do not develop SWD.

Shift work increases the subjective need for recovery from work²⁰⁾. Consequently, shift workers experience more fatigue on days off than dayworkers, and report extended sleep length, probably due to their increased need for restorative sleep²¹⁾. Field studies have shown that employees' sleep may be at its shortest directly after the last night shift²²⁾ or on the first day off after a night shift²³⁾, after which sleep duration increases. Garde *et al.*²²⁾ indicated that after the last night shift, police officers' sleep improved with more consecutive recovery days. In addition, Haluza *et al.*²⁴⁾ suggest that having three consecutive days off could improve recovery from shift work. Subjective sleepiness typically reverses after at least two nights of recovery sleep following a night shift²⁵⁾. We have previously found that in addition to inefficient sleep, SWD is associated with excessive sleepiness²⁶⁾ and slower recovery¹⁷⁾ on weekly days off. Yet, although they could alleviate the symptoms of SWD, longer recovery periods are rarely studied among shift workers.

To examine recovery during holidays in cases of SWD, we compared shift workers with and without questionnaire-based SWD during work and holiday periods. In addition to their general questionnaire responses, we utilised their responses to questions on sleep length at two separate points of time: during a work period and after at least two weeks of being on holiday. We had two main hypotheses. First, we hypothesised that shift workers both with and

without SWD would report longer sleep during holidays than on workdays. Secondly, we hypothesised that shift workers with SWD would experience longer sleep but similar sleep debt during holidays to those without SWD.

Subjects and methods

Study design and setting

This observational study consisted of three questionnaires and a three-week sleep diary and actigraphy registration (Fig. 1A. general questionnaire, B. Sleep diary and actigraphy monitoring, C. questionnaire on shift work period, and D. questionnaire on holiday period) at different points of time between 2012 and 2013. The shift workers completed the questionnaire on the shift work period (Fig. 1C) either before or during the three-week sleep diary and actigraphy monitoring period (Fig. 1B). We used the sleep diaries and actigraphy to observe and verify that the shift workers' symptoms and shift schedules (Fig. 1B) were in accordance with the ICSD-3 criteria of SWD¹⁷⁾ between May 2012 and July 2013. The interval between the questionnaires on shift work (Fig. 1C) and holidays (Fig. 1D) varied from one to six months. The study protocol was approved by the Coordinating Ethics Committee of the Hospital District of Helsinki and Uusimaa (35/13/03/00/12). The study followed the requirements of the Helsinki Declaration. All the participants provided written informed consent.

[Insert
Fig. 1
here]

Participants

The study included 54 full-time ground staff members of Helsinki Airport (Finland), whose shift schedules included early morning shifts (starting from 03:01 to 06:00) and/or night shifts (at least 3 h of work between 23:00 and 06:00 h), and evening shifts¹⁷⁾. Eleven shift workers discontinued due to health, personal, or unknown reasons. To form and compare clear SWD and clear non-SWD (i.e. reference) case groups, and to simplify the interpretation of the results,

we excluded a further 13 shift workers with milder SWD symptoms (only occurring ‘rather often’) who could not be distinctly classified as either SWD or reference cases (see criteria for the groups in ‘Shift work disorder’ section below)¹⁷⁾. The final sample was composed of 21 SWD cases and nine reference cases.

General questionnaire

The general questionnaire included questions on background characteristics, SWD, recovery, and sleep (Fig. 1A). It was used to define SWD and to explore aspects of recovery on a general level.

Shift work disorder

We formed the study groups on the basis of the answers to the shift-specific questions on insomnia and sleepiness (SS-Q)¹⁷⁾: ‘How often do you 1) experience difficulties initiating sleep, 2) awaken during a sleep period, 3) experience difficulties falling asleep after awakening, 4) experience non-restorative sleep, 5) experience daytime sleepiness, and 6) experience difficulties staying awake at work, in relation to i) morning shifts, ii) evening shifts, iii) night shifts, and iv) two weeks of being on holiday (excluding Question 6)’? The participants used a four-point scale (‘never/rarely’, ‘rather rarely’, ‘rather often’, or ‘often/continuously’) to answer the sub-questions (i–iv) of all the questions (1–6). ‘Not applicable’ was also a response option.

To fulfil the criteria of SS-Q-based SWD case, the shift workers had to report symptoms of insomnia or sleepiness ‘never/rarely’ or ‘rather rarely’ after two weeks of being on holiday,

and report at least one of the ‘never/rarely’ or ‘rather rarely’ occurring symptoms as ‘often/continuously’ in connection with morning, evening, and/or night shift. Nevertheless, one of the six symptoms could occur during the holiday period if at least one other symptom only occurred in connection with work shifts¹⁷⁾. To fulfil the criteria of SS-Q-based reference case, the shift workers had to report symptoms of insomnia and sleepiness as ‘never/rarely’ or ‘rather rarely’ in relation to both shifts and holidays. However, like the SWD cases, no more than one holiday-related symptom could occur ‘rather often’ or ‘often/continuously’¹⁷⁾.

To meet the ICSD-3 criteria of SWD, we used a three-week actigraphy and sleep diary monitoring (Fig. 1B and Appendix) to verify disturbed sleep-wake patterns and reduced total sleep time¹⁷⁾ and confirmed that the frequency of work shifts and the SWD symptoms were adequate^{16, 17)} (see Appendix T1–3). Our definition of SWD also complies with the circadian rhythm sleep-wake disorder – shift work type – in the 11th revision of the International Classification of Diseases²⁷⁾.

Other general questionnaire variables

The participants responded to items on age (yrs), sex, chronotype (four-point scale from ‘definitely a morning-type’ to ‘definitely an evening-type’ dichotomised to morning/evening-type)²⁸⁾, shift work experience (yrs), average length of shifts (h), obstructive sleep apnoea, sleep need (h)¹⁷⁾, and recovery from irregular working hours (four-point scale from ‘poor’ to ‘good’) (Fig. 1A).

Three-week actigraphy and sleep diary monitoring

The participants wore an actigraph Actiwatch AW7 (Cambridge Neurotechnology Ltd, Cambs, UK) on their non-dominant wrist for a three-week shift work period. We used Actiwatch Activity and Sleep Analysis 7 software (Cambridge Neurotechnology Ltd, Cambs, UK) to build a variable on average 24-hour total sleep time including workdays and days off (Fig. 1B) from one-minute epochs¹⁷⁾. In addition, the participants filled in a sleep diary in which they reported shift start and end times at bedtime (Fig. 1B)¹⁷⁾.

Questionnaires on work and holiday periods

The participants completed questionnaires during a visit to a medical centre once during a shift work period (Fig. 1C) and once after at least two weeks of being on holiday (Fig. 1D). The participants answered the questions ‘How long during the last two weeks have you slept on average per 24 h, including naps 1) on workdays (sleep length on workdays, h and min) and 2) on days off/holidays (answering during a work period: sleep length on days off; answering after holidays: sleep length during holidays; h and min)?’. We calculated sleep debt for workdays, days off, and holidays separately by subtracting self-reported sleep length on these days from self-reported sleep need^{17, 29)}.

To enable us to evaluate the presence of SWD symptoms in relation to the questionnaire responses on sleep length, the participants also estimated the occurrence of symptoms of insomnia and sleepiness in the last two weeks (Fig. 1C and D) as part of the questionnaires on work and holiday periods (see Appendix T4).

The comparison of shift work and holiday periods verified that the SWD symptoms decreased during holidays. Appendix T5 presents the results of the comparisons of the SWD and reference groups.

Statistical analyses

We used IBM SPSS Statistics 25.0 (IBM Corp., Armonk, NY, USA) and SAS 9.4 (SAS Institute Inc., Cary, North Carolina, USA) for the statistical analyses. We used the paired samples *t-test* to verify whether the length and timing of sleep differed during work shifts and on days off in the SWD group (Appendix T2–3). We used the Wilcoxon signed-ranks test to explore whether the occurrence of SWD symptoms during the shift work and holiday periods differed in the SWD and reference groups (Appendix T5). To compare the groups, we applied the independent samples *t-test*, the Mann–Whitney *U-test*, or Fisher’s exact test, depending on the scale and distribution of each variable. We applied the Wilcoxon signed-ranks test to compare the sleep length and sleep debt on workdays and days off, workdays and holidays, and days off and holidays of the SWD and reference groups. Exact and repeated logistic regression models were used to adjust the analyses for age or chronotype. However, the results of the chronotype-adjusted analyses are not shown, because they were similar to the results of the unadjusted and age-adjusted analyses.

Results

Sample characteristics

Table 1 presents the characteristics of the SWD ($n=21$, 76% men) and reference ($n=9$, 78% men) groups. The SWD group was on average seven yrs (95% CI -14, 0) younger than the reference group ($p=0.043$). The SWD group had a longer subjective sleep need than the reference group ($U=48.5$, $p=0.030$). The groups were comparable in terms of sex, chronotype, shift work experience, objective 24-hour total sleep time, work shifts, and having treated obstructive sleep apnoea.

[Insert
Table 1
here]

Self-estimated sleep length

Table 2 presents the sleep length of the study groups. Compared to that of the reference group, those with SWD reported significantly shorter sleep on workdays ($U=32.5$, $p=0.003$), but not on days off or during holidays, as indicated by the Mann–Whitney *U*-tests. The age-adjusted exact logistic regression analyses indicated similar results (workdays: OR 0.12, 95% CI 0.01–0.64, $p=0.007$).

[Insert
Table 2
here]

The Wilcoxon signed-ranks tests indicated significantly shorter sleep on workdays than on days off among those with SWD ($Z=-3.97$, $p<0.001$), but not among those in the reference group ($Z=-1.76$, $p=0.078$). Sleep was also significantly shorter on workdays than during holidays among those with SWD ($Z=-4.05$, $p<0.001$) and among those in the reference group ($Z=-2.39$, $p=0.017$). Sleep length did not significantly differ on days off and during holidays among those with SWD ($Z=-1.17$, $p=0.242$) or among those in the reference group ($Z=-1.89$, $p=0.058$). Apart from an analysis of workdays and

days off which was inconclusive, age-adjusted repeated logistic regression analyses indicated similar results (workdays and holidays: OR -0.00, 95% CI -0.00--0.00, $p=0.007$; days off and holidays: $p=0.568$) to the Wilcoxon signed-ranks tests.

Sleep debt

Fig. 2 presents the sleep debt of the groups. Those with SWD had a significantly greater sleep debt on workdays ($U=15.5$, $p<0.001$), days off ($U=50.5$, $p=0.040$), and during holidays ($U=42.0$, $p=0.016$) than those in the reference group, as indicated by the Mann–Whitney *U*-tests. Age-adjusted exact logistic regression analyses indicated that SWD was positively associated with sleep debt on workdays (OR 9.29, 95% CI 1.83–202.20, $p<0.001$) and during holidays (OR 2.41, 95% CI 1.14–6.58, $p=0.018$), but not on days off.

[Insert
Fig. 2
here]

Recovery

Those with SWD experienced good recovery from irregular working hours less often (14%, $n=3/21$) than those in the reference group (100%, $n=9/9$; $p<0.001$, Fig. 3). This was supported by an age-adjusted exact logistic regression analysis (OR 0.02, 95% CI 0.00–0.11, $p<0.0001$).

[Insert
Fig. 3
here]

DISCUSSION

We studied recovery during holidays in SWD utilising questionnaire responses on sleep length during work and holiday periods. This is the first study to investigate the amount of sleep or sleep debt in SWD cases during a naturalistic shift work washout period. Holidays were associated with longer sleep estimates than workdays among shift workers both with and without SWD. Whereas the employees without SWD reported longer sleep than sleep need during holidays, the employees with SWD estimated having similar sleep length and sleep need. This, together with sleep debt and suboptimal recovery from irregular working hours among the shift workers with SWD, suggests that the recovery of those with SWD is compromised.

As we expected, the employees both with and without SWD estimated that they slept longer during holidays than on workdays. Although longer sleep need has been associated with SWD¹⁷⁾ and dissatisfaction with shift work³⁰⁾, those with SWD did not report longer sleep during holidays than those without SWD. Contrary to our hypothesis, the employees with SWD estimated that they slept approximately the same amount during holidays as their perceived sleep need (i.e., they had zero hours of sleep debt), whereas those without SWD estimated that they slept more. Future studies could explore whether leaving shift work would diminish the group difference if, for example, the perceived sleep need in the different groups changed in different ways.

All the shift workers without SWD recovered well from irregular working hours, while the employees with SWD reported mainly suboptimal recovery. This is in line with studies that have associated the manifestation of SWD on workdays with shorter subjective sleep^{17–19)} and insufficient sleep³¹⁾. Compared to workdays, the employees with SWD estimated their sleep length as longer on days that were not workdays, but they still showed sleep debt on days off. Those without SWD estimated that they slept longer on days off than their subjective sleep need. Thus, in addition to recovery sleep, the reference group members' sleep might also be prophylactic, counteracting the negative effects of shift work on workdays following days off³²⁾. Those with SWD might lack this kind of prophylactic sleep. Indeed, objective 24-hour sleep was close to the reference cases' subjective sleep need, which is in line with a recent study of truck drivers³³⁾, whereas those with SWD slept over an hour less than their sleep need when not on holiday.

This is the first study of SWD to evaluate sleep length in relation to holidays and to compare responses after a real-life shift work washout period with responses during a work period. As SWD symptoms should be associated with a shift work schedule¹⁵⁾, and not with a long recovery period, we considered severe symptoms during holidays as being unrelated to SWD. This probably prevented us from confusing SWD with conditions of continuing insomnia and/or sleepiness. However, as SWD and other disorders can occur simultaneously, the participants of the current study could have one holiday-related insomnia/sleepiness symptom, provided that the separate symptom that defined a person as an SWD case did not occur

during holidays in the SS-Q. The work- and holiday-related questionnaire responses of those with SWD verified that the subjective frequency of the symptoms of insomnia and sleepiness did indeed decrease in relation to a real holiday. We also verified disturbed sleep-wake patterns and reduced total sleep time among those with questionnaire-based SWD using sleep diaries and actigraphy, as required by the ICSD-3 criteria.

The current study has limitations, such as its usage of subjective measures and variables averaged over two weeks. Thus, probable daily variation of sleep need and sleep debt in longer recovery from SWD remains to be studied. We used a subjective measure of sleep need, which can provide inaccurate or overly short estimations of actual sleep need. On the other hand, objective assessment of sleep need may not be feasible in an observative field setting if full recovery takes longer than the recovery periods available in a shift work schedule. In addition, in real life, habitual sleep (that has been used as an estimate for sleep need) can be shorter than actual sleep need, even if individuals are not shift workers^{34, 35}). Nevertheless, studies that use objective measures of sleep need, such as sleep restriction or sleep extension protocols, are required to confirm our results concerning sleep need. Furthermore, although the ICSD-3 states that SWD symptoms should manifest for at least three months, the SS-Q did not specify the manifestation period; it focused on generic symptoms in connection with different shift types. In addition, to simplify the interpretation of the results, we excluded shift workers with milder SWD symptoms whom we could not classify as either clear SWD cases or clear non-SWD cases. This added a contrast

between the groups. Statistical power was limited because of small the sample size, but it was adequate for observing significant differences between the groups. Further, younger age has been related to longer and better sleep³⁶⁾ and lesser need for recovery³⁷⁾ than older age, and in our study, the shift workers with SWD were younger than those without SWD, which may have diminished the differences between the groups. The younger age of the employees with SWD may have partially resulted from the healthy worker effect³⁸⁾, since those with SWD are possibly at a higher risk of leaving shift work.

To conclude, holidays increased the amount of self-estimated sleep, regardless of SWD status. However, the employees with SWD appeared to have a consistently longer sleep debt than those without SWD, which partially resulted from the greater sleep need among those with SWD. Moreover, the shift workers with SWD reported primarily suboptimal recovery from irregular working hours, unlike those without SWD, who all recovered well. All the above support the notion of less efficient recovery among shift workers with SWD. Future studies could examine whether shift workers with SWD would benefit from extended recovery periods in their roster or from individualised rota planning.

Conflict of interest

CH reports outside the present work a grant from Finska Läkaresällskapet and from the Finnish Work Environment Fund, and personal fees from the National Institute of Health and Welfare and the Orion Corporation. The other authors have no conflict of interest.

Acknowledgements

We thank Paula Niemelä for recruiting the participants and for making the practical arrangements, and Alice Lehtinen (Altexta) for the language editing of this article. The study was funded by the Finnish Work Environment Fund (111102) and NordForsk (74809).

References

- 1) Hublin C, Partinen M, Koskenvuo M, Kaprio J (2007) Sleep and mortality: a population-based 22-year follow-up study. *Sleep* **30**, 1245–53.
- 2) Kwok CS, Kontopantelis E, Kuligowski G, Gray M, Muhyaldeh A, Gale CP, Peat GM, Cleator J, Chew-Graham C, Loke YK, Mamas MA (2018) Self-reported sleep duration and quality and cardiovascular disease and mortality: a dose-response meta-analysis. *J Am Heart Assoc* **7**, e008552.
- 3) Yin J, Jin X, Shan Z, Li S, Huang H, Li P, Peng X, Peng Z, Yu K, Bao W, Yang W, Chen X, Liu L (2017) Relationship of sleep duration with all-cause mortality and cardiovascular events: a systematic review and dose-response meta-analysis of prospective cohort studies. *J Am Heart Assoc* **6**, e005947.
- 4) Klerman EB, Dijk DJ (2008) Age-related reduction in the maximal capacity for sleep--implications for insomnia. *Curr Biol* **18**, 1118–23.
- 5) Van Dongen HP, Maislin G, Mullington JM, Dinges DF (2003) The cumulative cost of additional wakefulness: dose-response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. *Sleep* **26**, 117–26.
- 6) De Bloom J, Geurts SAE, Kompier MAJ (2013) Vacation (after-) effects on employee health and well-being, and the role of vacation activities, experiences and sleep. *J Happiness Stud* **14**, 613–33.

- 7) Petersen H, Kecklund G, D'Onofrio P, Axelsson J, Åkerstedt T (2017) Thank god it's Friday - sleep improved. *J Sleep Res* **26**, 567–71.
- 8) Swanson LM, Arnedt JT, Rosekind MR, Belenky G, Balkin TJ, Drake C (2011) Sleep disorders and work performance: findings from the 2008 National Sleep Foundation Sleep in America poll. *J Sleep Res* **20**, 487–94.
- 9) Groeger JA, Zijlstra FR, Dijk DJ (2004) Sleep quantity, sleep difficulties and their perceived consequences in a representative sample of some 2000 British adults. *J Sleep Res* **13**, 359–71.
- 10) Åkerstedt T (2003) Shift work and disturbed sleep/wakefulness. *Occup Med (Lond)* **53**, 89–94.
- 11) Sallinen M, Kecklund G (2010) Shift work, sleep, and sleepiness - differences between shift schedules and systems. *Scand J Work Environ Health* **36**, 121–33.
- 12) Vedaa Ø, Harris A, Bjorvatn B, Waage S, Sivertsen B, Tucker P, Pallesen S (2016) Systematic review of the relationship between quick returns in rotating shift work and health-related outcomes. *Ergonomics* **59**, 1–14.
- 13) Flo E, Pallesen S, Moen BE, Waage S, Bjorvatn B (2014) Short rest periods between work shifts predict sleep and health problems in nurses at 1-year follow-up. *Occup Environ Med* **71**, 555–61.
- 14) American Academy of Sleep Medicine (2005) International classification of sleep disorders: Diagnostic and coding manual (2nd Ed.) American Academy of Sleep Medicine, Westchester, IL.

- 15) American Academy of Sleep Medicine (2014) International classification of sleep disorders (3rd Ed.) American Academy of Sleep Medicine, Darien, IL.
- 16) Vanttola P, Puttonen S, Karhula K, Oksanen T, Härmä M (2019) Prevalence of shift work disorder among hospital personnel: a cross-sectional study using objective working hour data. *J Sleep Res* **29**, e12906.
- 17) Vanttola P, Härmä M, Viitasalo K, Hublin C, Virkkala J, Sallinen M, Karhula K, Puttonen S (2019) Sleep and alertness in shift work disorder: findings of a field study. *Int Arch Occup Environ Health* **92**, 523–33.
- 18) Gumenyuk V, Howard R, Roth T, Korzyukov O, Drake CL (2014) Sleep loss, circadian mismatch, and abnormalities in reorienting of attention in night workers with shift work disorder. *Sleep* **37**, 545–56.
- 19) Kalmbach DA, Pillai V, Cheng P, Arnedt JT, Drake CL (2015) Shift work disorder, depression, and anxiety in the transition to rotating shifts: the role of sleep reactivity. *Sleep Med* **16**, 1532–8.
- 20) Jansen N, Kant I, van Amelsvoort L, Nijhuis F, van den Brandt P (2003) Need for recovery from work: evaluating short-term effects of working hours, patterns and schedules. *Ergonomics* **46**, 664–80.
- 21) Härmä M, Karhula K, Puttonen S, Ropponen A, Koskinen A, Ojajärvi A, Kivimäki M (2019) Shift work with and without night work as a risk factor for fatigue and changes in sleep length: a cohort study with linkage to records on daily working hours. *J Sleep Res* **28**, e12658.

- 22) Garde AH, Nabe-Nielsen K, Jensen MA, Kristiansen J, Sorensen JK, Hansen AM (2020) The effects of the number of consecutive night shifts on sleep duration and quality. *Scand J Work Environ Health* **46**, 446–53.
- 23) Totterdell P, Spelten E, Smith L, Barton J, Folkard S (1995) Recovery from work shifts: how long does it take? *J Appl Psychol* **80**, 43–57.
- 24) Haluza D, Schmidt VM, Blasche G (2019) Time course of recovery after two successive night shifts: a diary study among Austrian nurses. *J Nurs Manag* **27**, 190–6.
- 25) Åkerstedt T, Kecklund G, Gillberg M, Lowden A, Axelsson J (2000) Sleepiness and days of recovery. *Transportation Research Part F* **3**, 251–61.
- 26) Vanttola P, Puttonen S, Karhula K, Oksanen T, Härmä M (2020) Employees with shift work disorder experience excessive sleepiness also on non-work days: a cross-sectional survey linked to working hours register in Finnish hospitals. *Ind Health* **58**, 366–74.
- 27) World Health Organization. International Classification of Diseases 11th Revision for Mortality and Morbidity Statistics (Version: 04/2019). <https://icd.who.int/browse11/l-m/en#/http%3a%2f%2fid.who.int%2fid%2fentity%2f756278911>. Accessed August 15, 2019.
- 28) Horne JA, Östberg O (1976) A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int J Chronobiol* **4**, 97–110.

- 29) Hublin C, Kaprio J, Partinen M, Koskenvuo M (2001) Insufficient sleep--a population-based study in adults. *Sleep* **24**, 392–400.
- 30) Axelsson J, Åkerstedt T, Kecklund G, Lowden A (2004) Tolerance to shift work-how does it relate to sleep and wakefulness? *Int Arch Occup Environ Health* **77**, 121–9.
- 31) Waage S, Moen BE, Pallesen S, Eriksen HR, Ursin H, Åkerstedt T, Bjorvatn B (2009) Shift work disorder among oil rig workers in the North Sea. *Sleep* **32**, 558–65.
- 32) Axelsson J, Vyazovskiy VV (2015) Banking sleep and biological sleep need. *Sleep* **38**, 1843–5.
- 33) Onninen J, Pylkkönen M, Tolvanen A, Sallinen M (2021) Accumulation of sleep loss among shift-working truck drivers. *Chronobiol Int* **38**, 1344–53.
- 34) Kitamura S, Katayose Y, Nakazaki K, Motomura Y, Oba K, Katsunuma R, Terasawa Y, Enomoto M, Moriguchi Y, Hida A, Mishima K (2016) Estimating individual optimal sleep duration and potential sleep debt. *Sci Rep* **6**, 35812.
- 35) Klerman EB, Dijk DJ (2005) Interindividual variation in sleep duration and its association with sleep debt in young adults. *Sleep* **28**, 1253–9.
- 36) Costa G, Sartori S (2007) Ageing, working hours and work ability. *Ergonomics* **50**, 1914–30.
- 37) Kiss P, De Meester M, Braeckman L (2008) Differences between younger and older workers in the need for recovery after work. *Int Arch Occup Environ Health* **81**, 311–20.

- 38) Knutsson A, Åkerstedt T (1992) The healthy-worker effect: self-selection among Swedish shift workers. *Work & Stress* **6**, 163–7.

ALL FIGURE TITLES AND LEGENDS

Fig. 1. Study design and variables/measures

Fig. 2. Box-and-whisker plot of sleep debt. ○ = outlier. SWD (black), Reference (grey). Mann–Whitney *U*-test: * = $p < 0.05$, *** = $p < 0.001$

Fig. 3. Percentage of shift workers with good recovery from irregular working hours. Fisher's exact test: *** = $p < 0.001$

Appendix

Do holidays change subjective sleep length or sleep debt in shift work disorder?

Industrial Health

Päivi Vanttola¹, Mikko Härmä¹, Christer Hublin¹, Katriina Viitasalo², Mikael Sallinen^{1,3}, Jussi Virkkala^{1,4},
Sampsa Puttonen²

¹ Finnish Institute of Occupational Health, Finland;

² Finnair Health Services, Finland; ³ Department of Psychology, University of Jyväskylä, Finland;

⁴ Department of Clinical Neurophysiology, University of Helsinki and Helsinki University Hospital, Finland

E-mail: paivi.vanttola@ttl.fi

Actigraphy and sleep diary monitoring to verify that each individual with SS-Q-based SWD met the ICSD-3-based criteria of SWD

The participants filled in a sleep diary twice a day during a three-week shift work period. At bedtime, they reported shift start and end times, the greatest Karolinska Sleepiness Scale sleepiness¹⁾ of each day, and bedtime. At awakening, they evaluated wake-up time, sleep latency, the number of awakenings, and the quality of sleep (five-point scale from 1 = ‘good’ to 5 = ‘poor’). We calculated the mid-point of sleep from these variables. In addition, the participants wore an actigraph Actiwatch AW7 (Cambridge Neurotechnology Ltd, Cambs, UK) on their non-dominant wrist. We used Actiwatch Activity and Sleep Analysis 7 software (Cambridge Neurotechnology Ltd, Cambs, UK) to build the variables: sleep latency, wake after sleep onset, sleep efficiency, and total sleep time from one-minute epochs²⁾.

Frequency of work shifts and SWD symptoms. Those with SWD (n = 21) had from 7 to 15 work shifts (morning, evening and/or night shifts) during the three-week monitoring period. In connection with these work shifts they had individual numbers of symptoms, presented at group level in Appendix Table 1.

Appendix Table 1. Number of days with SWD symptoms among SWD cases during three-week monitoring period

Number of days with symptoms	Sleep diary					Actigraphy				
	Mean	MD	SD	(Q ₁ , Q ₃)	Range	Mean	MD	SD	(Q ₁ , Q ₃)	Range
Sleep latency > 30 min	2.5	1	2.9	(1, 4)	(0, 11)	0.5	0	0.8	(0, 1)	(0, 2)
Wake after sleep onset > 30 min						7.7	9	3.9	(5, 11)	(0, 12)
≥ 4 awakenings during sleep period	2.1	1	2.3	(0, 4)	(0, 7)					
Sleep efficiency ≤ 85%						2.6	1	3.6	(0, 5)	(0, 11)
Sleep quality poor or rather poor	2.7	2	1.7	(2, 4)	(0, 6)					
Total sleep time < 6 h	5.2	5	2.8	(3, 8)	(1, 12)	5.8	6	2.7	(4, 8)	(1, 11)
Karolinska Sleepiness Scale ≥ 7	7.4	8	3.1	(5, 11)	(1, 12)					

MD, median; SD, standard deviation; Q₁ 1st quartile; Q₃, 3rd quartile

Reduced total sleep time. Each SWD case’s total sleep time decreased by at least one hour in relation to work shifts. Appendix Table 2 presents the group-level difference between total sleep time of the shifts and days off.

Appendix Table 2. Difference between total sleep time in connection with shifts and total sleep time in connection with days off among SWD cases during three-week monitoring period

Reduction of total sleep time	Sleep diary				Actigraphy			
	Mean Δ	(95% CI)	t (df)	p ^a	Mean Δ	(95% CI)	t (df)	p ^a
Morning shifts (h)	2.1	(1.6, 2.6)	9.023 (20)	<0.0001	1.7	(1.3, 2.1)	8.584 (20)	<0.0001
Evening shifts (h)	1.0	(0.6, 1.4)	5.167 (20)	<0.0001	1.0	(0.7, 1.3)	6.752 (20)	<0.0001
Night shifts (h)	2.7	(2.1, 3.2)	10.649 (16)	<0.0001	2.4	(1.9, 2.9)	10.258 (16)	<0.0001

Δ, difference; CI, Confidence interval; ^a Paired samples *t*-test

- 1) Ingre M, Kecklund G, Åkerstedt T, Kecklund L (2004) Variation in sleepiness during early morning shifts: a mixed model approach to an experimental field study of train drivers. *Chronobiol Int* **21**(6), 973-90.
- 2) Vanttola P, Härmä M, Viitasalo K, Hublin C, Virkkala J, Sallinen M, Karhula K, Puttonen S (2019) Sleep and alertness in shift work disorder: findings of a field study. *Int Arch Occup Environ Health* **92**, 523-533.

Disturbed sleep-wake pattern. The SWD cases' timing of sleep varied considerably. Appendix Table 3 presents the group-level difference between the mid-points of sleep for shifts and days off.

Appendix Table 3. Difference between mid-sleep in connection with shifts and in connection with days off among SWD cases during three-week monitoring period

Delay in mid-sleep	Sleep diary		t (df)	p^a	Actigraphy		t (df)	p^a
	Mean Δ	(95% CI)			Mean Δ	(95% CI)		
Morning shifts (h)	-0.8	(-0.3, -1.3)	-3.151 (20)	<0.01	-0.7	(-0.2, -1.2)	-2.857 (20)	<0.01
Evening shifts (h)	1.4	(0.9, 1.8)	6.427 (20)	<0.0001	1.6	(1.1, 2.0)	7.125 (20)	<0.0001
Night shifts (h)	6.3	(5.6, 7.0)	18.853 (16)	<0.0001	6.4	(5.8, 6.9)	25.269 (16)	<0.0001

Δ , difference; CI, Confidence interval; ^a Paired samples *t*-test

Questions to evaluate the presence of SWD symptoms in relation to sleep length estimations

The participants completed questions on SWD symptoms (Appendix Table 4) as part of the questionnaires on sleep length during a visit to a medical centre once during a shift work period and once after at least two weeks of being on holiday (Figure 1C–D). The questions were similar to the Shift-specific questions on insomnia and sleepiness (SS-Q, Figure 1A) used to define SWD in the current study.

Appendix Table 4. Questions on occurrence of symptoms of insomnia and sleepiness

Questions/symptoms	Less than once a week	On 1–2 days per week	On 3–5 days per week	Often/continuously
1. How often have you experienced difficulties initiating sleep in the last two weeks?	1	2	3	4
2. How often have you awakened during a sleep period in the last two weeks?	1	2	3	4
3. How often have you experienced difficulties falling asleep after awakening in the last two weeks?	1	2	3	4
4. How often have you experienced non-restorative sleep in the last two weeks?	1	2	3	4
5. How often have you experienced daytime sleepiness in the last two weeks?	1	2	3	4
6. How often have you experienced difficulties staying awake in the last two weeks?	1	2	3	4

Appendix Table 5. Occurrence of SWD symptoms during shift work and holiday periods among those with SWD ($n=21$) and those in the reference group ($n=9$), based on questions in Appendix Table 4

Symptoms of insomnia/sleepiness ^a	SWD					Reference				
	Work period		Holiday period		<i>p</i> ^b	Work period		Holiday period		<i>p</i> ^b
	MD	(Q ₁ , Q ₃)	MD	(Q ₁ , Q ₃)		MD	(Q ₁ , Q ₃)	MD	(Q ₁ , Q ₃)	
Difficulties initiating sleep	2	(1, 2)	1	(1, 2)	0.161	1	(1, 1)	1	(1, 1)	1.000
Awakenings during a sleep period	3	(3, 4)	2	(2, 4)	0.013	2	(2, 3)	2	(1, 2)	0.011
Difficulties falling asleep after awakening	1	(1, 1)	1	(1, 1)	0.414	1	(1, 1)	1	(1, 1)	1.000
Non-restorative sleep	2	(2, 2)	2	(1, 2)	0.035	1	(1, 1)	1	(1, 1)	0.317
Daytime sleepiness	3	(2, 3)	2	(1, 2)	0.000	2	(1, 2)	2	(1, 2)	0.414
Difficulties staying awake	2	(1, 2)	1	(1, 1)	0.003	1	(1, 1)	1	(1, 1)	1.000

MD, median; Q₁ 1st quartile; Q₃, 3rd quartile; ^a In last two weeks: Scale: 1 = less than once a week, 2 = on 1–2 days per week, 3 = on 3–5 days per week, and 4 = often/continuously; ^b Wilcoxon signed-ranks test

The shift workers with SWD reported awakenings during a sleep period ($Z=-2.5$, $p=0.013$), non-restorative sleep ($Z=-2.1$, $p=0.035$), daytime sleepiness ($Z=-3.5$, $p<0.001$), and difficulties staying awake ($Z=-3.0$, $p<0.010$) significantly more seldom during a holiday period than during a work period. The frequency of difficulties initiating sleep and the frequency of difficulties falling asleep after awakening did not significantly differ during work and holiday periods among those with SWD. Those in the reference group reported awakenings during a sleep period ($Z=-2.5$, $p=0.011$) during a holiday period significantly more seldom than during a work period. As expected, most of the SWD symptoms appeared to decrease in the SWD group during holidays, whereas only one symptom decreased in the reference group. The participants of the current study did not seem to frequently have difficulties initiating sleep or falling asleep after awakening.

Table 1. Sample characteristics

Variable	SWD						Reference						<i>p</i>		
	n (group)	Mean	(SD)	Median	(IQR)	%	(n)	n (group)	Mean	(SD)	Median	(IQR)		%	(n)
Sex (men)	21					76	(16)	9					78	(7)	1.000 ^c
Age (yr)	21	41	(8)					9	48	(7)					0.043 ^a
Morning chronotype	21					38	(8)	9					56	(5)	0.443 ^c
Sleep need (h)	21			8.0	(2.0)			9			7.0	(1.5)			0.030 ^b
24-hour TST (h) – actigraphy	21	6.7	(0.7)					9	7.1	(0.4)					0.066 ^a
Treated obstructive sleep apnoea	21					5	(1)	9					11	(1)	0.517 ^c
Shift work experience (yr)	21	17	(8)					9	23	(13)					0.183 ^a
Length of shifts (h)	21			9.0	(1.2)			9			9.7	(1.0)			0.156 ^b
Early morning shifts per month	10			5.5	(4.8)			5			5.3	(3.5)			0.499 ^b
Any morning shift per month	21			6.8	(3.6)			8			5.6	(3.5)			0.250 ^b
Evening shifts per month	21			4.3	(1.7)			9			4.1	(1.7)			0.666 ^b
Night shifts per month	17			5.0	(1.6)			8			4.6	(3.8)			0.884 ^b

IQR, interquartile range

TST, total sleep time

^a Independent samples *t*-test

^b Mann–Whitney *U*-test

^c Fisher’s Exact Test

^d ‘Absolutely morning type’ or ‘more morning than evening type’

Table 2. Self-estimated sleep length among those with SWD (n=21) and those in reference group (n=9)

Sleep length during last two weeks	SWD		Reference		<i>p</i> ^a
	Median	(Q ₁ , Q ₃)	Median	(Q ₁ , Q ₃)	
Workdays (h) ^b	6.3	(6.0, 7.0)	7.0	(7.0, 7.8)	0.003
Days off (h) ^b	8.0	(7.0, 8.5)	7.5	(7.0, 8.5)	0.744
Holidays (h) ^c	8.0	(7.4, 8.5)	8.5	(7.8, 9.0)	0.238

Q₁ 1st quartile; Q₃, 3rd quartile

^a Mann–Whitney U test

^b Response during work period

^c Response after at least two weeks of being on holiday

**A. General
questionnaire**

- Characteristics
- Shift work disorder (SS-Q)
- Sleep need
- Recovery

**B. Three-week
actigraphy and sleep
diary monitoring**

- Verification of SWD symptoms, reduced total sleep time, and disturbed sleep-wake pattern among participants with shift work disorder (Appendix)
- 24-hour total sleep time
- Frequency of different work shifts

**C. Questionnaire
during shift work
period**

- Sleep length on workdays
- Sleep length on days off
- Sleep debt on workdays
- Sleep debt on days off
- Symptoms of insomnia and sleepiness (Appendix)

**D. Questionnaire
after ≥ 2 weeks of
being on holiday**

- Sleep length during holidays
- Sleep debt during holidays
- Symptoms of insomnia and sleepiness (Appendix)