LETTERS TO THE EDITOR

METABOLIC AND RESPIRATORY CHANGES IN ACUTE CARBON MONOXIDE POISONING TREATED BY HYPERBARIC OXYGENATION

There has as yet been no report which has concerned itself with the effect of the hyperbaric oxygenation (OHP) on metabolic and respiratory changes in acute carbon monoxide (CO) poisoning, though OHP has been widely applied with satisfactory results. To elucidate this problem, blood gases and acid-base balance of 109 patients suffering from severe acute CO poisoning were investigated. All patients were comatose or semicomatose on admission to the respiratory care unit for OHP. They are divided into three groups based upon their outcomes. The first is the improved group of patients, whose consciousness was fully recovered after the treatments (89 cases). The second is the group of unchanged patients who were survived in vegetative state (14 cases). The last group is of the non-survived patients, who had fatal outcomes within 10 days of CO poisoning (6 cases). The patients who had sustained cardiac arrest before the first OHP therapy were excluded from this study to clarify the changes caused by CO alone.

Blood samples were taken from the femoral artery at the time of admission and shortly after the first OHP treatment under room air breathing. They were analyzed for respiratory gases and acid-base status with an IL Meter Model 113. Base excess (BE) was obtained by using Siggaard-Andersen nomogram from PaCO_2 , pH and Hb.

Immediately after the sampling of the blood and taking chest radiogram, patients were treated by OHP at a pressure of two atmospheres absolute (ATA) of pure O_2 for 1 hr. All patients received the infusion of lactate Ringer's solution throughout the initial treatment, and the correction of acidosis by sodium bicarbonate was not done before and during the first OHP treatment.

Table 1 shows mean values and standard errors of blood gases and acid-base status in three groups before and shortly after the first OHP therapy. At the time of admission arterial pH of the died group was remarkably reduced compared with the improved and unchanged groups. BE was distinctly negative in three groups of patients, especially that of died group was enormously low. This metabolic acidosis seemed to be compensated by respiratory alkalosis in improved and unchanged groups, since PaCO_2 was low and arterial pH was within normal at that time. On the contrary, marked hypocarbia was observed under remarkably low pH of the arterial blood in died group. It could be understood that respiratory compensation for metabolic acidosis was insufficient in the latter group of patients.

However, hypocarbia remained in three groups of the patients even after the first OHP treatment. In these determinations BE of the improved and unchanged

Table 1. Blood gases and acid-base status before and shortly after the first OHP therapy. (Mean $\pm$ S.E.)

		Improved	Unchanged	Died
Admission	pH	7.38 $\pm$ 0.01	7.39 $\pm$ 0.03	7.20 $\pm$ 0.05
	pO ₂	87.4 $\pm$ 2.6	85.0 $\pm$ 5.4	84.8 $\pm$ 4.8
	pCO ₂	28.4 $\pm$ 0.7	26.5 $\pm$ 1.4	22.6 $\pm$ 3.1
	BE	-6.2 $\pm$ 0.6	-7.3 $\pm$ 2.0	-16.6 $\pm$ 3.1
After first OHP	pH	7.46 $\pm$ 0.01†	7.49 $\pm$ 0.02†	7.41 $\pm$ 0.02†
	pO ₂	85.6 $\pm$ 2.7	79.0 $\pm$ 4.6	54.0 $\pm$ 7.9†
	pCO ₂	31.3 $\pm$ 0.6	29.8 $\pm$ 2.1	29.9 $\pm$ 1.5
	BE	-0.6 $\pm$ 0.5†	-0.8 $\pm$ 1.3†	-2.8 $\pm$ 1.3†

†: Statistically significant difference from the data on admission ($p < 0.01$)

patients were within normal, and of the died was nearly normal. These changes of PaCO₂ and BE resulted in alkalotic appearance of arterial pH in the improved and unchanged group, or normal pH in the died group shortly after the first OHP therapy.

The acid-base alteration following the first OHP above mentioned is more clearly depicted in Fig. 1. The broad range of values of pH, PaCO₂ and BE on admission is illustrated in Fig. 1a. PaCO₂ in almost all the patients were below 35 mmHg, and BE were also negative values below -2.3 mEq/l. Thus respiratory

Fig. 1. Metabolic shift from admission (a) to after the first OHP therapy (b).

Note respiratory alkalosis remained even after metabolic acidosis was corrected.

METABOLIC AND RESPIRATORY CHANGES IN CO POISONING

alkalosis and metabolic acidosis were common findings at the time of admission. The change by the first OHP therapy is demonstrated in Fig. 1b. Compared with Fig. 1a, the widespread shift of patients to the normal metabolic zone is evident. Almost all values were plotted between two oblique lines representing $BE \pm 2.3$ mEq/l showing metabolic acidosis was corrected by the initial OHP. On the other hand, $PaCO_2$ was not changed and respiratory alkalosis still remained after the metabolic acidosis was corrected. The persistent hypocarbia in the normal or nearly normal metabolic state indicates that there should be another mechanism to continue hypocarbia than compensation for metabolic acidosis alone.

In our previous study, we have revealed that blood volume was reduced remarkably following experimental CO poisoning by increased vascular permeability¹⁾. In this state perivascular edema and membrane permeability change in tissue cells as seen in shock state would make tissue hypoxia persistent even after CO-Hb has diminished by OHP. It would be also taken into consideration that brain edema caused by the mechanism above mentioned contributes to develop respiratory alkalosis through the change in bicarbonate ions in cerebrospinal fluid²⁾.

We had already reported that the progressive widening of A-aDO₂ in fatal patients suffering from CO poisoning³⁾. In our present study PaO_2 of died group fell significantly after OHP treatment, and was impossible to maintain breathing under room air. The possible mechanisms to develop progressive hypoxemia were mentioned previously³⁾. In addition to them, persistent alveolar-capillary block caused by perivascular edema could be considered as another mechanism. Pulmonary edema was frequently observed in acute CO poisoning as reported elsewhere⁴⁾, and it would also be concerned with this problem.

REFERENCES

- 1) Ogawa, M., Shimazaki, S., Tanaka, N., Ukai, T. and Sugimoto, T. (1973). *Igaku no Ayumi*, **84**, 201. (in Japanese)
- 2) Katsurada, K., Sugimoto, T. and Onji, Y. (1969). *J. Trauma*, **9**, 799.
- 3) Ogawa, M., Tamura, H., Katsurada, K. and Sugimoto, T. (1972). *Med. J. Osaka Univ.*, **22**, 251.
- 4) Ogawa, M., Katsurada, K., Sugimoto, T. and Sone, S. (1973). *Jap. J. Chest Dis.*, **32**, 268. (in Japanese)

Respiratory Care Unit
Department of Traumatology
Osaka University Hospital
Dojima Hamadori
Fukushima-ku, Osaka

Michio OGAWA
Shuji SHIMAZAKI
Kikushi KATSURADA
Tsuyoshi SUGIMOTO

(Received June 6, 1973)