Mini Review

Recognition of bacterial compounds by aortic endothelial cells activates Weibel-Palade body exocytosis

Takeshi Into* and Kenji Matsushita

Department of Oral Disease Research, National Institute for Longevity Sciences, National Center for Geriatrics and Gerontology, Obu, Aichi, Japan

The endothelial cell-specific granule Weibel-Palade body releases vasoactive substances capable of modulating vascular inflammation, thrombosis and atherogenesis. Pathogen recognition by endothelial Toll-like receptors (TLRs) is thought to play a crucial role in promotion of vascular inflammatory responses. However, the molecular basis for the early-phase responses of endothelial cells to pathogens has not been fully elucidated. We found that human aortic endothelial cells respond to several TLR ligands, including bacterial lipoteichoic acids and lipopeptides, but not lipopolysaccharides, to induce Weibel-Palade body exocytosis, accompanied by release of the storage component von Willebrand factor. In this mini-review, we briefly describe a novel function of endothelial TLRs and discuss its implications in aortic inflammation or atherogenesis.

Rec.10/26/2006, Acc.11/30/2006, pp112-116

* Correspondence should be addressed to:

Takeshi Into, Department of Oral Disease Research, National Institute for Longevity Sciences, National Center for Geriatrics and Gerontology, 36-3 Gengo, Morioka, Obu, Aichi 474-8522, Japan. e-mail: into@nils.go.jp.

Key words endothelial cell, exocytosis, Toll-like receptor, inflammation

Introduction

Excess inflammatory responses of vessel walls or endothelium to invading pathogens have recently been suggested to be linked to atherogenesis^{1,2)}. Several common bacterial infectious agents or pathogens, such as *Chlamydia pneumoniae*, *Helicobacter pylori*, *Porphyromonas gingivalis* and other commensal bacteria, have been detected in vessel walls or atherosclerotic lesions in humans³⁻⁶⁾. However, the linkage between endothelial innate recognition of such pathogens and endothelial inflammatory responses has not been fully elucidated. For the detection of

circulating bacteria in blood, several Toll-like receptors (TLRs) on endothelial cell surfaces are thought to identify molecular motifs that usually compose bacterial bodies or cell walls. TLR2 can detect a wide range of common bacterial constituents, such as lipoteichoic acids (LTA), peptidoglycans (PGN), bacterial di- or triacylated lipoproteins/lipopeptides, lipoarabinomannans, porins and fimbriae⁷⁾. TLR1 and TLR6 further participate in the accurate discrimination of these molecules by TLR2⁸⁾. TLR4 and TLR5 contribute to the recognition of lipopolysaccharides (LPS) and flagellin, respectively⁸⁾. After recognition of cognate ago-

Fig. 1 The endothelial cell-specific granule Weibel-Palade body

Weibel-Palade bodies contain several vasoactive substances as shown in the figure. The presence of the granules in HAEC could be detected by immunofluorescent staining (IF) using antibodies to VWF (left picture) and Ang-2 (right picture).

nists, endothelial TLRs activate a signaling pathway utilizing MyD88, IL-1R-associated kinase (IRAK) 1 and TNFR-associated factor (TRAF) 6, which ultimately activate I κ B and the release and translocation of active NF- κ B to the nucleus⁸). The endothelial NF- κ B signaling pathway downstream of TLRs is thought to participate in the development of vascular inflammatory diseases or atherogenesis through the promotion of expression of large number of proinflammatory mediators and adhesion molecules⁹⁻¹¹). However, it is still not known whether endothelial TLRs are primary initiators or modulators of vascular inflammatory diseases.

Induction of Weibel-Palade body exocytosis by bacterial constituents

We aimed to determine the early-phase inflammatory responses of human aortic endothelial cells (HAEC) to bacterial pathogens, especially their cell wall constituents. It is known that early endothelial activation involves dual phases: rapid translocation of P-selectin to the endothelial surface and slower synthesis and expression of adhesion molecules such as ICAM-1. The former process is accompanied by rapid exocytosis of Weibel-Palade bodies, endothelial cell-specific storage granules that contain vasoactive substances, including von Willebrand factor (VWF), P-selectin, IL-8, eotaxin-3, endothelin-1, CD63/lamp3, tissue plasminogen activator, angiotensin-2 and osteoprotegerin¹²) (Fig.1). These proteins are transported to the out-

Table 1 Recognition of bacterial constituents by TLRs

Component	TLR recognition
<i>E. coli</i> O26:B6 LPS	TLR4
<i>Salmonella minnesota</i> R595 LPS	TLR4
<i>Salmonella typhimurium</i> flagellin	TLR5
<i>S. aureus</i> LTA	TLR2/TLR6
<i>S. aureus</i> PGN	TLR2
Pam ₃ CSK ₄ (triacylated lipopeptide)	TLR2/TLR1
FSL-1 (diacylated lipopeptide)	TLR2/TLR6
MALP-2 (diacylated lipopeptide)	TLR2/TLR6

Fig. 2 VWF release by bacterial constituents

HAEC were stimulated with increasing concentrations of bacterial constituents shown in Table 1 for 60 min. The amount of VWF released into the culture medium was measured by ELISA.

side of the cell upon stimulation or vascular damage through Weibel-Palade body exocytosis and may control local or systemic biological effects, including vascular inflammation, thrombosis and atherogenesis¹²).

As bacterial constituents, we prepared LPS from *E. coli* and *Salmonella minnesota*, flagellin from *S. typhimurium*, PGN and LTA from *Staphylococcus aureus* and the synthetic analogs of bacterial lipoproteins Pam₃CSK₄ (*N*-palmitoyl-S-dipalmitoylglyceryl CSKKKK), FSL-1 (*S*-dipalmitoylglyceryl CGDPKHPKSF) and MALP-2 (*S*-dipalmitoylglyceryl

Fig. 3 Involvement of TLR2 and TLR6 in Weibel-Palade body exocytosis

HAEC transfected with specific siRNA oligonucleotides for mRNA of TLR2 or TLR6 were stimulated with the increasing concentrations of bacterial constituents shown in Table 1 for 60 min. The amount of VWF released into the culture medium was measured by ELISA.

Fig. 4 Regulation of TLR2 expression in HAEC

HAEC were pretreated with IFN- γ or precultured under laminar flow for 12 h. The cell surface TLR2 expression was measured by flow cytometry.

CGNNDENISFKK). These compounds are known to be recognized by cognate TLRs¹³ as shown in Table 1. We examined the release of VWF, an essential molecule for the formation of Weibel-Palade bodies¹⁴, after stimulation of HAEC with these bacterial constituents for 60 min. We found that LTA, Pam₃CSK₄, FSL-1, MALP-2 and, to a lesser extent, flagellin induced release of VWF in a dose-dependent manner (Fig.2). The stimulatory activity of PGN was very weak. Interestingly, the TLR4 agonist LPS did not activate Weibel-Palade body exocytosis despite its potent activity to induce IL-8 production after stimulation of HAEC for 4 h, suggesting that TLR4 lacks the ability to induce exocytosis. This may be related to evidence that TLR4 intracellularly localizes in artery endothelial cells¹⁵. We must further investigate the differential potential of endothelial TLRs in the induction of Weibel-Palade body exocytosis.

Induction of TLR2-mediated Weibel-Palade body exocytosis

We focused on TLR2-mediated Weibel-Palade body exocytosis, because several TLR2 agonists could activate potent re-

sponses (Fig.2). HAEC express TLR2 at a weak but evident level on their cell surface (Fig.4), whereas expression of TLR2 and responsiveness to TLR2 agonists have been reported in human endothelial cells from other endothelial beds such as HUVEC¹⁶. We found that ‘knockdown’ of TLR2 expression by siRNA almost completely suppressed the magnitude of VWF release by all TLR2 agonists in HAEC (Fig.3). TLR6 interference also significantly decreased the activities of LTA, FSL-1 and MALP-2 and even Pam₃CSK₄ (Fig.3). Therefore, endothelial recognition of bacterial compounds by TLR2 and TLR6 directly contributes to Weibel-Palade body exocytosis.

Regulation of TLR2-mediated Weibel-Palade body exocytosis

Recently, complete deficiency of TLR2 in atherosclerosis-prone LDLR-null mice was shown to lead to an apparent reduction in the formation of lesions even in the absence of known exogenous TLR2 agonists¹⁷. Interestingly, the progression of lesions was indicated to be probably caused by aortic endothelial cells, not cells from bone marrow¹⁷. Indeed, several proper-

ties of endothelial TLR2 have been implicated in the disease development. Specifically, endothelial TLR2 expression is enhanced by inflammatory stimuli, such as TNF- α , IFN- γ and LPS¹⁸⁾. Indeed, the expression level of TLR2 is known to be increased at atherosclerotic lesions in humans¹⁹⁾. We examined whether regulation of endothelial TLR2 expression modulates exocytosis. We found that increased endothelial TLR2 expression after IFN- γ treatment (Fig.4) increased the magnitude of TLR2-mediated exocytosis of Weibel-Palade body, suggesting enhanced responsiveness of endothelial cells to pathogens in an inflamed or atherosclerotic lesion. Physiologic fluid shear stress has been suggested to have atheroprotective effects *in vivo*, since atherosclerosis preferentially occurs in an area of disturbed flow or a low level of shear stress, whereas regions with steady laminar flow and physiologic shear stress are protected. Disturbed flow or a low level of shear stress has been reported to regulate expression of various regulatory molecules of endothelial activation, by which atherosclerotic processes may be accelerated in the sites. Physiologic fluid shear stress is known to decrease endothelial TLR2 expression via impaired activity of the transcriptional factor SP1²⁰⁾. Indeed, TLR2 expression slightly decreased in HAEC incubated under laminar flow (Fig.4). We found that laminar flow decreased TLR2-mediated exocytosis. Thus, TLR2 agonist-induced Weibel-Palade body exocytosis may be influenced by surface expression level of physiologically or pathologically regulated endothelial TLR2.

Conclusions

Our findings show a linkage between endothelial innate recognition of circulating bacteria and early-phase endothelial inflammatory responses. These findings may provide a new insight into the roles of endothelial TLRs in the initiation and modulation of the development of vascular inflammation and atherosclerosis.

References

- 1) Gibson FC 3rd, Hong C, Chou HH, Yumoto H, Chen J, Lien E, Wong J, Genco CA: Innate immune recognition of invasive bacteria accelerates atherosclerosis in apolipoprotein E-deficient mice. *Circulation*, 109: 2801-2806, 2004.
- 2) Ross R: Atherosclerosis--an inflammatory disease. *N Engl J Med*, 340: 115-126, 1999.
- 3) Desvarieux M, Demmer RT, Rundek T, Boden-Albala B, Jacobs DR Jr, Sacco RL, Papapanou PN: Periodontal microbiota and carotid intima-media thickness: the Oral Infections and Vascular Disease Epidemiology Study (INVEST). *Circulation*, 111: 576-582, 2005.
- 4) Haraszthy VI, Zambon JJ, Trevisan M, Zeid M, Genco RJ: Identification of periodontal pathogens in atheromatous plaques. *J Periodontol*, 71: 1554-1560, 2000.
- 5) Kuo CC, Shor A, Campbell LA, Fukushima H, Patton DL, Grayston JT: Demonstration of *Chlamydia pneumoniae* in atherosclerotic lesions of coronary arteries. *J Infect Dis*, 167: 841-849, 1993.
- 6) Stelzel M, Conrads G, Pankuweit S, Maisch B, Vogt S, Moosdorf R, Flores-de-Jacoby L: Detection of *Porphyromonas gingivalis* DNA in aortic tissue by PCR. *J Periodontol*, 73:868-870, 2002.
- 7) Kirschning CJ, Schumann RR: TLR2: cellular sensor for microbial and endogenous molecular patterns. *Curr Top Microbiol Immunol*, 270: 121-144, 2002.
- 8) Akira S, Takeda K: Toll-like receptor signalling. *Nat Rev Immunol*, 4: 499-511, 2004.
- 9) Hajishengallis G, Sharma A, Russell MW, Genco RJ: Interactions of oral pathogens with toll-like receptors: possible role in atherosclerosis. *Ann Periodontol*, 7: 72-78, 2002.
- 10) Michelsen KS, Doherty TM, Shah PK, Arditi M: TLR signaling: an emerging bridge from innate immunity to atherogenesis. *J Immunol*, 173: 5901-5907, 2004.
- 11) Mullaly SC, Kubes P: Toll gates and traffic arteries: from endothelial TLR2 to atherosclerosis. *Circ Res*, 95: 657-659, 2004.
- 12) Lowenstein CJ, Morrell CN, Yamakuchi M: Regulation of Weibel-Palade body exocytosis. *Trends Cardiovasc Med*, 15: 302-308, 2005.
- 13) Takeda K, Kaisho T, Akira S: Toll-like receptors. *Annu Rev Immunol*, 21: 335-376, 2003.
- 14) Denis CV, Andre P, Saffaripour S, Wagner DD: Defect in regulated secretion of P-selectin affects leukocyte recruitment in von Willebrand factor-deficient mice. *Proc Natl Acad Sci USA*, 98: 4072-4077, 2001.
- 15) Dunzendorfer S, Lee HK, Soldau K, Tobias PS: Toll-like receptor 4 functions intracellularly in human coronary artery endothelial cells: roles of LBP and sCD14 in mediating LPS responses. *FASEB J*, 18: 1117-1119, 2004.
- 16) Oude Nijhuis CS, Vellenga E, Daenen SM, Kamps WA, De Bont ES: Endothelial cells are main producers of interleukin 8 through Toll-like receptor 2 and 4 signaling during bacterial infection in leukopenic cancer patients. *Clin Diagn Lab Immunol*, 10: 558-563, 2003.
- 17) Mullick AE, Tobias PS, Curtiss LK: Modulation of atherosclerosis in mice by Toll-like receptor 2. *J Clin Invest*, 115:

3149-3156, 2005.

- 18) Faure E, Thomas L, Xu H, Medvedev A, Equils O, Arditi M: Bacterial lipopolysaccharide and IFN- γ induce Toll-like receptor 2 and Toll-like receptor 4 expression in human endothelial cells: role of NF- κ B activation. *J Immunol*, 166: 2018-2024, 2003.
- 19) Edfeldt K, Swedenborg J, Hansson GK, Yan ZQ: Expres-

sion of toll-like receptors in human atherosclerotic lesions: a possible pathway for plaque activation. *Circulation*, 105: 1158-1161, 2002.

- 20) Dunzendorfer S, Lee HK, Tobias PS: Flow-dependent regulation of endothelial Toll-like receptor 2 expression through inhibition of SP1 activity. *Circ Res*, 95: 684-691, 2004.