Application of HRT/ERT for Japanese women

Japan is going to face a tremendously rapid rise of postmenopausal women in the population in a few decades. In order to promote a better quality of life in Japanese elderly women HRT/ERT is expected to be a health promoting regimen. Because Japanese women are not so obese and are less likely to suffer from IHD compared with women in America and Europe, and because Leiden mutation of factor V and prothrombin abnormality which facilitate the incidence of venous thrombosis in HRT/ERT users have not been found in Japanese, it is presumed that HRT/ERT would prevent IHD in Japanese women with fewer severe adverse reactions such as pulmonary thrombus. However, in order to prevent breast cancer and uterine cancer as well as to raise bone mineral density and decrease IHD, application of selective estrogen receptor modulator (SERM) including raloxifene is awaited (11). Our experiences have taught us that the optimal prescription of HRT/ERT for Japanese women in terms of fewer adverse reactions on lipids and coagulation/fibrinolysis system is a half dose of CEE plus MPA, estradiol patch and estriol (12).

References

- 1) Mendelsohn ME, Karas RH. The protective effects of estrogen on the cardiovascular system. *N Engl J Med* **340**: 1801–1811, 1999.
- 2) Mosca L. The role of hormone replacement therapy in the prevention of postmenopausal heart disease. *Arch Intern Med* **160**: 2263–2272, 2000.
- 3) Grodstein F, Stampfer MJ, Colditz GA, et al. Postmenopausal hormone therapy and mortality. *N Engl J Med* **336**: 1769–1775, 1997.
- 4) Hu FB, Stampfer MJ, Manson JE, et al. Trends in the incidence of coronary heart disease and changes in diet and lifestyle in women. *N Engl J Med* **343**: 530–537, 2000.
- 5) Writing Group for the Women's Health Initiative Investigators: Risks and benefits of estrogen plus progestin in healthy postmenopausal women. Principal results from the Women's Health Initiative randomized controlled trial. *JAMA* **288**: 321–333, 2002.
- 6) Hodis HN, Mack WJ, Lobo R. What is the cardioprotective role of hormone replacement therapy. *Cur Atheroscler Report* **5**: 56–66, 2003.
- 7) Hulley S, Grady D, Bush T, et al. Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. *JAMA* **280**: 605–613, 1998.
- 8) Grady D, Herrington D, Bitter V, et al. Cardiovascular disease outcomes during 6.8 years of hormone therapy. Heart and Estrogen/progestin Replacement Study follow-up (HERS II). *JAMA* **288**: 49–57, 2002 (Erratum in *JAMA* **288**: 1064, 2002).
- 9) Hodis HN, Mack WJ, Robo RA, et al. Estrogen in the prevention of atherosclerosis: a randomized, double-blind, placebo-controlled trial. *Ann Intern Med* **135**: 939–953, 2001.
- 10) Byington RP, Furberg CD, Herrington DM, et al. Effect of estrogen plus progestin on progression of carotid atherosclerosis in postmenopausal women with heart disease: HERS B-mode substudy. *Arterioscler Thromb Vasc Biol* **22**: 1692–1697, 2002.
- 11) Barret-Connor E, Grady D, Sashegyi A, et al. Raloxifene and cardiovascular events in osteoporotic postmenopausal women: four-year results from the MORE (Multiple Outcomes of Raloxifene Evaluation) randomized trial. *JAMA* **287**: 847–857, 2002.
- 12) Ouchi Y ed. The Guideline for Hormone Replacement Therapy to Promote Health of Elderly Females (in Japanese). Medical Review Co., Tokyo, 2001.

3. Androgen Deficiency and Its Disorders in Men

Taiji TSUKAMOTO, Naoki ITOH and Shin-ichi HISASUE

Department of Urology, Sapporo Medical University School of Medicine, Sapporo

Key words: androgen, aging male, hypogonadism, HRT

Introduction

Hypergonadotropic and hypogonadotropic hypogonadism are representative diseases of androgen deficiency in men. In our recently aged society the number of aged men with hypogonadism and its associated disorders, that is andropause, has been increasing. The WHO raised this issue for discussion on the improved care of aged men (1). It is well accepted that the androgen level in men declines gradually with aging, which is contrast to that in women. A study already reported the diagnosis and treatment of andropause in

1944 (2). Unfortunately, after that, little progress was made in this area until recently. It was only 3 years ago that the American Endocrine Society held a consensus meeting on this issue and indicated that there were many issues remaining to be discussed (3).

In this review, we will summarize the current situation of diagnosis and treatment of partial androgen deficiency of the aging male (PADAM), andropause, in our country.

Decline of androgen with aging in men

Androgens, in particular, testosterone (T), decline gradually with aging. In our cross-sectional study, when free-testosterone (fT) was measured, it was found to reach a peak in the 20s, followed by a gradual decline with age. However,

Reprint requests should be addressed to Dr. Taiji Tsukamoto, Department of Urology, Sapporo Medical University School of Medicine, Sapporo, 060-8556

the extent of the decline differed and there was a wide variation in each individual. This is in contrast to that in women, all of whom show a constant and abrupt decline of estrogen (E) when they have menopause. The decline of T with aging is supported as well by the Baltimore Longitudinal Study of Aging, in which the decline was confirmed in each individual with a 10- to 30-year follow-up (4). The study estimated that hypogonadism was found in 10% of men in their 50s, 20% in their 60s and 30% in their 70s, when it was defined as a serum T level lower than 11.3 nmol/l (325 ng/dl).

Androgen decline with aging and associated disorders

It is clear that the androgen level of men declines with aging. However, the definition of the T level indicating hypogonadism in aged men is not concretely established. In addition, not all disorders of various organs or diseases caused by a decline of androgen with aging have been revealed. Although T has many target organs in addition to those of the prostate and testis, such as the brain, liver, kidney, cardiovascular systems, muscle and bone, it is not fully understood to what extent a decline of T affects the functions of these organs. This is in clear contrast to estrogen, which has been extensively studied in disorders and diseases associated with its deficiency.

While sporadic, a few studies have reported the influence of T decline with aging on the cardiovascular system and bone. In a meta-analysis of 30 cross-sectional case control studies, 18 suggested levels of androgens such as T, fT dihydroepiandrosterone (DHEA), DHEA-S or androstenedione in men were inversely correlated with the incidence of coronary heart diseases (CHD). Thus, a decline of the androgen level with aging may contribute to an increase of CHD in men (5). Unfortunately, we do not exactly know the mechanism by which decreases of androgen levels participate in the increase in the incidence of CHD.

The decline of androgen levels with aging in men is also speculated to participate in the loss of bone density in aged men. It is well accepted that young men with hypogonadism have a loss of bone density and testosterone replacement therapy dramatically improves the density. In addition, castration and LH-RH agonist treatment induce significant losses of bone density in patients with prostate cancer. However, the participation of estrogen has been indicated in the loss of bone density, although the direct involvement of T in the loss has not been fully understood. For example, the Rancho Bernard Study showed that bioavailable estrogen, as well as bioavailable T, was strongly involved in the loss (6). The Mayo Clinic Study reported a stronger influence of bioavailable estradiol (E2) than total T on bone density (7). Finally, in the Dubbo Osteoporosis Epidemiology Study, E2 was more influential than fT (8). In fact, synthetic E administration for patients with prostate cancer who received castration or an LH-RH agonist improved bone metabolism (9). Thus, these studies may emphasize the involvement of E as well as T in maintaining normal bone density in men.

Diagnosis and treatment of PADAM

How do we diagnose patients as having PADAM? The disease is defined as a syndrome consisting of biochemical abnormality (low T) and symptoms and signs associated with hypogonadism. However, measurement of T with biological activity and establishment of its normal range are not easy. The total T level does not always indicate the biological activity of T. The fT does not either. In addition, the dialysis method for the measurement of fT is ideal but not simple. It cannot be used as a routine measurement. Although the measurement of bioavailable T may reflect the biological activity of T, it is not practical. A recent study suggested that the calculated free T (CFT) estimated by total T, sex-hormone binding globulin and albumin, may be useful for the diagnosis of PADAM (10). We have evaluated the clinical relevance of CFT in patients who visited us complaining of symptoms related to PADAM. So far, we have found that the CFT levels declined parallel with increasing age and fT levels. Further study will be necessary to confirm these findings.

Patients with PADAM generally present psychological, physical and sexual function-related symptoms. Depression, irritation and anxiety are such symptoms. Physical symptoms consist of things such as joint and muscle-related ones, sweating, hot flashes, sleep disorders, and loss of memory and concentration capacities. Some patients may have biochemical abnormalities such as loss of bone density, lipid metabolism disturbance and excessive obesity. For these patients, symptoms, association of the systemic diseases, the external genitalia, the prostate (digital rectal examination), serum prostate-specific antigen, and the levels of androgens should be routinely examined. Bone density and lipid-related profiles should be assessed. For evaluation of symptoms related to PADAM, the St. Louis University ADAM questionnaire and aging males symptoms' rating scale are frequently used (11, 12). We have been validating these questionnaires that were translated into Japanese and. So far, however, we have found that the St. Louis University ADAM questionnaire seems to have a low specificity in diagnosing patients with PADAM.

Finally, we refer to androgen-replacement therapy. While patients with PADAM may generally be indicated for therapy, the best candidates for treatment are not yet known. Of particular, concerns are the clinical manifestations of prostate cancer and benign prostatic hyperplasia possibly caused by the therapy. It may be valid for aged men who have a total T level lower than 200 ng/dl (3). However, the replacement therapy should not be indicated for those without symptoms or signs related to PADAM even when they have a low testosterone level. Patients who have total T levels between 200 and 400 ng/dl should be evaluated for symptoms and signs related to PADAM. In this situation, measurement of fT and bioavailable T may be recommended, but the abnormal levels of these hormones are not necessarily established. Thus, androgen replacement therapy for PADAM must be assessed for risk and benefit.

Conclusion

The studies that have been done to date suggest that a decline of androgen with aging may produce various disorders more frequently than previously anticipated. However, recognition of these disorders in the medical community has just started. Without more extensive understanding of the physiological actions of androgens on various organs other than the testis and prostate, we would not be able to provide PADAM patients with the most appropriate diagnosis and treatment.

References

- 1) World Health Organization. Report of the World Health Organization: Active ageing: a policy framework. *Aging Male* **5**: 1–37, 2002.
- 2) Heller CG, Myers GB. The male climacteric; its symptomatology, diagnosis and treatment. *JAMA* **126**: 472–477, 1944.
- 3) Swerdloff RS. Summary of the consensus session from the 1st annual meeting. The Endocrine Society, 2000.
- 4) Harman SM, Metter EJ, Tobin JD, Pearson J, Blackman MR. Longitudinal effects of aging on serum total and free testosterone levels in healthy men. *J Clin Endocrinol Metab* **86**: 724–731, 2001.
- 5) Alexandersen P, Haarbo J, Christiansen C. The relationship of natural androgens to coronary heart disease in males: a review. *Atherosclerosis* **125**: 1–13, 1996.
- 6) Greendale GA, Edelstein S, Barrett-Corner E. Endogenous sex steroids and bone mineral density in older women and men: Rancho Bernard Study. *J Bone Miner Res* **12**: 1833–1843, 1997.
- 7) Khosla S, Melton LJ3rd, Atkinson EJ, O'Fallon WM, Klee GG, Riggs BL. Relationship of serum sex steroid levels and bone turnover markers with bone mineral density in men and women: a key role for bio-available estrogen. *J Clin Endocrinol Metabol* **83**: 2266–2274, 1998.
- 8) Center JR, Nguyen TV, Sambrook PN, Eisman JA. Hormonal and biochemical parameters in the determination of osteoporosis in elderly men. *J Clin Endocrinol Metabol* **84**: 3626–3635, 1998.
- 9) Scherr D, Pitts WR Jr, Vaughn ED Jr. Diethylstilbesterol revisited: androgen deprivation, osteoporosis and prostate cancer. *J Urol* **167**: 535–538, 2002.
- 10) Vermeulen A, Verdonck L, Kaufman JM. A critical evaluation of simple methods for the estimation of free testosterone in serum. *J Clin Endocrinol Metab* **84**: 3666–3672, 1999.
- 11) Morley JE, Charlton E, Patric KP, et al. Validation of a screening questionnaire for androgen deficiency in aging males. *Metabolism* **49**: 1239–1242, 2000.
- 12) Heinemann LAJ, Saad F, Thiele K, Wood-Dauphinees S. The aging males' symptoms (AMS) rating scale: cultural and linguistic validation into English. *Aging Males* **4**: 14–22, 2001.

4. Physiological Significance of Replacement Therapy of Dehydroepiandrosterone

Toshihiko YANASE

Department of Medicine and Bioregulatory Science, Graduate School of Medical Sciences, Kyushu University, Kyushu

Key words: DHEA, DHEA-S, aging, hormone replacement therapy

DHEA and aging

Dehydroepiandrosterone (DHEA) and its interconvertible sulfate ester, DHEA sulfate (DHEA-S) are mainly produced by the adrenal glands and their production decreases progressively with aging (1). The decrease of DHEA (-S) with aging is enzymatically explained by the relative reduction of 17, 20-lyase activity to 17 α -hydroxylase activity, both of which are catalyzed by a single polypeptide enzyme, cytochrome P450c17. The P450c17 activity or its mRNA is also decreased with aging and this may be related to the impairment of P450c17 by adrenal lipid peroxidation (1). The relatively selective reduction of 17,20-lyase activity with aging may be associated with the reduction of cytochrome P450 reductase

activity (1) or simply explained by the volume reduction (atrophy) of zona reticularis which is responsible for the production of DHEA (-S) (2).

Physiological significance of DHEA

DHEA and DHEA-S have recently attracted wide attention because their serum levels were shown to be one of the important markers for longevity (3) and because many investigators including us have clarified the beneficial effects of DHEA on dementia, obesity, lipid metabolism, diabetes mellitus, atherosclerosis, and osteoporosis, etc (Fig. 1) (1, 4–11). Sunderland et al first reported that the serum concentration of DHEA-S in Alzheimer's disease (AD) is decreased in comparison to controls (4). But, in our study, not only the patients with AD and but also the patients with cerebrovascular dementia (CVD) were found to have lower concentrations of DHEA-S compared with age- and gender-matched controls (5). A reduced concentration of the serum

Reprint requests should be addressed to Dr. Toshihiko Yanase, Department of Medicine and Bioregulatory Science, Graduate School of Medical Sciences, Kyushu University, Kyushu, 812-8582