Cyst Infection in Autosomal Dominant Polycystic Kidney Disease

I-kuan Wang¹⁻³, Chih-Chia Liang³, Yung-Fang Chen⁴ and Chiu-Ching Huang³

Key words: autosomal dominant polycystic kidney disease, pyocyst, urinary tract infection

(Intern Med 51: 1803, 2012)

(DOI: 10.2169/internalmedicine.51.7313)

Picture 1.

Picture 2.

ver and right flank pain lasting for 3 days. She had a history of autosomal dominant polycystic kidney disease (ADPKD), and chronic kidney disease stage V for several years. Urine analysis showed hematuria and pyuria. Urine culture was negative. An unenhanced computed tomography (CT) examination revealed multiple renal cysts, some of which showed hemorrhage (Picture 1). Despite antibiotic therapy, the patient's fever persisted for several days. MRI showed multiple renal cysts and a predominant 5.4 cm cyst in the upper pole of the right kidney with a fluid-fluid level suggestive of infected cyst on T2-weighted image (Picture 2).

It is difficult to diagnose an infected renal cyst in patients with ADPKD, due to limitations of conventional ultrasound and CT imaging (1). Isolation of an infected renal cyst by MRI has been reported, but experience is limited, as to date there are few reports in the literature (2).

The authors state that they have no Conflict of Interest (COI).

Acknowledgement

The present study was supported by Taiwan Department of Health Clinical Trial and Research Center of Excellence (DOH 101-TD-B-111-004).

References

1. Gupta S, Seith A, Sud K, et al. CT in the evaluation of complicated autosomal dominant polycystic kidney disease. *Acta Radiol* **41**: 280-284, 2000.
2. Chicoskie C, Chaoui A, Kuligowska E, Dember LM, Tello R. MRI isolation of infected renal cyst in autosomal dominant polycystic kidney disease. *Clin Imaging* **25**: 114-117, 2001.

A 22-year-old woman was admitted to the hospital for fe-

¹Graduate Institute of Clinical Medical Science, College of Medicine, China Medical University, Taiwan, ²Department of Internal Medicine, College of Medicine, China Medical University, Taiwan, ³Division of Nephrology, China Medical University Hospital, Taiwan and ⁴Department of Radiology, China Medical University Hospital, Taiwan

Received for publication January 14, 2012; Accepted for publication January 18, 2012

Correspondence to Dr. Chiu-Ching Huang, cch@mail.cmuh.org.tw

© 2012 The Japanese Society of Internal Medicine Journal Website: <http://www.naika.or.jp/imindex.html>