Intractable Hiccup in Demyelinating Disease with Anti-Myelin Oligodendrocyte Glycoprotein (MOG) Antibody

Tetsuya Akaishi¹, Masatoshi Konno², Ichiro Nakashima¹ and Masashi Aoki¹

Key words: anti-MOG antibody, intractable hiccup, cortical lesions, pulvinar lesions, medullary lesions

(Intern Med 55: 2905-2906, 2016)

(DOI: 10.2169/internalmedicine.55.7146)

Picture.

The patient was a 29-year-old man who presented with intractable hiccup at the onset of demyelinating disease. This was followed, in quick succession, by blurred vision, gait disturbance with myoclonic spasms and paresthesia in

both legs, and the inability to urinate. Brain and spinal MRI revealed lesions in the parasagittal cingulate gyri, pulvinar, cerebral peduncles, superior colliculi, middle cerebellar peduncles, inferior medulla, and upper cervical cord (Pic-

¹Department of Neurology, Tohoku University Graduate School of Medicine, Japan and ²Department of Neurology, Yamagata City Hospital Saiseikan, Japan

Received for publication January 18, 2016; Accepted for publication February 7, 2016

Correspondence to Dr. Ichiro Nakashima, nakashima@med.tohoku.ac.jp

ture). The patient was diagnosed with neuromyelitis optica spectrum disorder (NMOSD) based on the clinical findings (1). However, a serum analysis showed that the patient was negative for serum anti-aquaporin-4 (AQP4) antibody and positive for anti-myelin oligodendrocyte glycoprotein (MOG) antibody. We therefore considered that the correct diagnosis was acute demyelinating disease associated with anti-MOG-antibody. Two courses of steroid pulse therapy and oral prednisolone therapy alleviated the symptoms within two months. Although intractable hiccup is a characteristic manifestation of NMOSD with anti-AQP4-antibody (2), it can occur in patients with anti-MOG-antibody-associated diseases whose medullary lesions are not located in the area postrema.

The authors state that they have no Conflict of Interest (COI).

References

1. Wingerchuk DM, Banwell B, Bennett JL, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology* **85**: 177-189, 2015.
2. Misu T, Fujihara K, Nakashima I, Sato S, Itoyama Y. Intractable hiccup and nausea with periaqueductal lesions in neuromyelitis optica. *Neurology* **65**: 1479-1482, 2005.

The Internal Medicine is an Open Access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).