Vaginal Amebiasis Mimicking Cervical Cancer

Masayuki Ota and Naoki Yanagisawa

Key words: vaginal amebiasis, cervical cancer

(Intern Med 55: 3071-3072, 2016)

(DOI: 10.2169/internalmedicine.55.7289)

Picture 1.

Picture 2.

A 46-year-old woman presented with lower abdominal pain, diarrhea and irregular vaginal bleeding persisting for the past 2 months. She had a history of cervical cancer treated with concurrent chemoradiotherapy 6 years previously, which was successful. Magnetic resonance imaging with T2-weighted and diffusion images showed a high inten-

sity area in the cervix, suggestive of recurrent cervical cancer (Picture 1). However, a biopsy specimen of the cervix was compatible for amebiasis on hematoxylin/eosin and periodic acid-Schiff staining (Picture 2). A stool specimen demonstrated an ameba trophozoite, which was later confirmed to be *Entamoeba histolytica*. Blood tests were nega-

tive for HIV, syphilis, and hepatitis B, but positive for hepatitis C. Treatment with metronidazole resulted in a rapid clinical response. Vaginal amebiasis may mimic symptoms and imaging findings of cervical cancer (1, 2). Physicians must be aware of this rare disease, and a tissue biopsy is essential to make an accurate diagnosis.

The authors state that they have no Conflict of Interest (COI).

References

1. Arvind A, Minakshi B. Cervical amoebiasis mimicking cervical carcinoma, a rare presentation of a common infection. *J Infect Public Health* 2015 (in press).
2. Antony SJ, Lopez-Po P. Genital amoebiasis: historical perspective of unusual disease presentation. *Urology* **54**: 952-955, 1999.

The Internal Medicine is an Open Access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).