[PICTURES IN CLINICAL MEDICINE]

Multiple Hemorrhagic Cerebral Cortical Lesions in Neuro-Behçet's Disease

Kazuyuki Saito, Takashi Watanabe and Shuta Toru

Key words: neuro-Behçet's disease, intracranial hemorrhage, cerebral venous thrombosis, magnetic resonance imaging

(Intern Med 56: 2377-2378, 2017)

(DOI: 10.2169/internalmedicine.8632-16)

Picture A: DWI

Picture B: MR venography

Picture C: T1WI

Picture D: gadolinium T1WI

Picture

A 27-year-old Japanese man presented with erythema nodosum and consciousness opacity of four days in duration. He had panuveitis and fever. A cerebrospinal fluid (CSF) analysis revealed 100/mm³ pleocytosis (polynuclear cells, 87%) and the elevation of interleukin-6 (5,630 pg/mL). Brain MRI demonstrated multiple hemorrhagic cerebral cor-

tical lesions with gadolinium enhancement (Picture A, DWI; C, T1WI; D, gadolinium T1WI), but normal MR venography (Picture B). Cerebral venous thrombosis was not proven. The patient was positive for human leukocyte antigen-B51. He did not completely fulfil the International Study Group criteria for acute neuro-Behçet's disease

Department of Neurology, Nitobe Memorial Nakano General Hospital, Japan

Received: November 30, 2016; Accepted: January 18, 2017; Advance Publication by J-STAGE: August 10, 2017

Correspondence to Dr. Kazuyuki Saito, kaz_saito228@yahoo.co.jp

(NBD) (1) because he lacked oral ulceration; however, the acute onset of neurological symptoms and CSF pleocytosis of $>6.2/\text{mm}^3$ both matched the criteria. He was treated with prednisolone (1 mg/kg/day; the dose was tapered by 5 mg per month and reduced to 5 mg/day over two years) and infliximab (5 mg/kg bimonthly) for one year, which resulted in remission. Although cortical lesions are rare in NBD (2), it is important to consider NBD when such atypical lesions are encountered.

The authors state that they have no Conflict of Interest (COI).

References

1. Hirohata S, Kikuchi H, Sawada T, et al. Clinical characteristics of neuro-Behçet's disease in Japan: a multicenter retrospective analysis. *Mod Rheumatol* **22**: 405-413, 2012.
2. Farahangiz S, Sarhadi S, Safari A, et al. Magnetic resonance imaging findings and outcome of neuro-Behçet's disease: the predictive factors. *Int J Rheumat Dis* **15**: 3142-3149, 2012.

The Internal Medicine is an Open Access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

© 2017 The Japanese Society of Internal Medicine
Intern Med 56: 2377-2378, 2017