[PICTURES IN CLINICAL MEDICINE]

Meningococcal Bursitis

Rentaro Oda¹, Yoshiyuki Sekikawa² and Igen Hongo²

Key words: *Neisseria meningitidis*, meningococcemia, bursitis, septic bursitis

(Intern Med 56: 3403-3404, 2017)

(DOI: 10.2169/internalmedicine.9173-17)

Picture 1.

Picture 2.

Picture 3.

A healthy 72-year-old Japanese woman complained of a chilly sensation and arthralgia for the previous 4 days. She had a fever (37.7°C); a physical examination revealed bilateral swollen ankle bursae (Picture 1, 2). Thrombocytopenia ($3.1 \times 10^9/\mu\text{L}$) and an elevated C-reactive protein level (29.23 mg/dL) were observed. Gram staining of the bursa fluid obtained by aspiration showed Gram-negative cocci (Pic-

ture 3). Blood and bursa fluid cultures revealed *Neisseria meningitidis*. Meningococcemia and septic bursitis were diagnosed. No risk factors, such as complement deficiency or anatomic or functional asplenia, for meningococcal disease were noted. Ceftriaxone was administered for 14 days. Septic bursitis typically results from skin injury or trauma but rarely from bacteremia, as observed in this case. Septic bursitis is most commonly caused by *Staphylococcus aureus* (1, 2), and onset due to other bacteria is rare, with no cases of septic bursitis due to *Neisseria meningitidis* having been reported. Aspiration of the swollen bursa is necessary to confirm the diagnosis of septic bursitis.

The authors state that they have no Conflict of Interest (COI).

References

1. Zimmermann B 3rd, Mikolich DJ, Ho G Jr. Septic bursitis. *Semin Arthritis Rheum* **24**: 391-410, 1995.
2. Cea-Pereiro JC, Garcia-Mejide J, Mera-Varela A, Gomez-Reino JJ. A comparison between septic bursitis caused by *Staphylococcus aureus* and those caused by other organisms. *Clin Rheumatol* **20**: 10-14, 2001.

¹Department of Infectious Diseases, Tokyo Bay Urayasu Ichikawa Medical Center, Japan and ²Division of Infectious Diseases, Musashino Red Cross Hospital, Japan

Received: March 10, 2017; Accepted: April 26, 2017; Advance Publication by J-STAGE: October 11, 2017

Correspondence to Dr. Rentaro Oda, odarenmomovivi@hotmail.co.jp

The Internal Medicine is an Open Access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To

view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

© 2017 The Japanese Society of Internal Medicine
Intern Med 56: 3403-3404, 2017