[PICTURES IN CLINICAL MEDICINE]

Focal Eosinophilic Myositis

Akio Kawabe¹, Kazuhisa Nakano¹, Aya Nawata^{1,2} and Yoshiya Tanaka¹

Key words: focal eosinophilic myositis, eosinophilic myositis, eosinophils

(Intern Med 59: 141, 2020)

(DOI: 10.2169/internalmedicine.3602-19)

Picture.

A 72-year-old man presented to the hospital with a 2-day history of rapidly progressive, painful redness, swelling, and warmth of the right thigh (Picture A). His temperature was 38.9°C; the creatine kinase level was 2,139 U/L. Magnetic resonance imaging (Picture B) showed extensive hyperintensity in the muscle (arrows) and intermuscular fascia (arrowheads) on short-tau inversion recovery imaging. These findings suggested necrotizing fasciitis. However, a histological examination (Picture C, Hematoxylin and Eosin staining) revealed endomysial mononuclear cell infiltration with eosinophils (arrows) and muscle fiber necrosis with regeneration (arrowheads). Subsequently, the eosinophil count increased to 2,223/ μ L. The patient was diagnosed with focal eosinophilic myositis, a rare and self-limited disease with localized lower limb pain and swelling without other organ involvement (1). The thigh can also be affected (1, 2), but the etiologic factors were not identified in the patient. At one-month

follow-up after the presentation, the symptoms had spontaneously resolved, with normal laboratory values shown.

The authors state that they have no Conflict of Interest (COI).

References

1. Selva-O'Callaghan A, Trallero-Araguás E, Grau JM. Eosinophilic myositis: an updated review. *Autoimmun Rev* **13**: 375-378, 2014.
2. Kaufman LD, Kephart GM, Seidman RJ, et al. The spectrum of eosinophilic myositis. Clinical and immunopathogenic studies of three patients, and review of the literature. *Arthritis Rheum* **36**: 1014-1024, 1993.

The Internal Medicine is an Open Access journal distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

¹The First Department of Internal Medicine, School of Medicine, University of Occupational and Environmental Health, Japan and ²Department of Pathology, School of Medicine, University of Occupational and Environmental Health, Japan

Received: July 2, 2019; Accepted: July 8, 2019; Advance Publication by J-STAGE: August 21, 2019

Correspondence to Dr. Yoshiya Tanaka, tanaka@med.uoeh-u.ac.jp

© 2020 The Japanese Society of Internal Medicine. *Intern Med* 59: 141, 2020