[CASE REPORT]

Distribution of Deep Gray Matter Lesions on Magnetic Resonance Imaging in Lymphomatosis Cerebri

Zen Kobayashi¹, Sawako Sakai¹, Sakiko Itaya², Yoshiyuki Numasawa³, Kiyobumi Ota⁴,
Miho Akaza⁵, Yasuhiro Ueda⁶, Shinichi Ogawa⁷, Shoichiro Ishihara¹,
Hiroyuki Tomimitsu¹ and Shuzo Shintani¹

Abstract:

We herein report the distribution of gray matter lesions on magnetic resonance imaging (MRI) in two patients with lymphomatosis cerebri (LC). In our patients, the fluid-attenuated inversion recovery sequence of brain MRI demonstrated a bilateral and diffuse high signal intensity, not only in the white matter but also in the thalamus, globus pallidus, putamen, and hippocampus. Among the deep gray matter, the caudate head and putamen (striatum) were relatively spared when compared with the globus pallidus, thalamus, and hippocampus. Interestingly, we found seven previous reports of similar MRI findings, with relative sparing of the striatum, in patients with LC. This finding may be characteristic of LC and help facilitate its diagnosis. Further investigations of a larger number of LC patients are necessary to confirm these findings.

Key words: caudate head, deep gray matter, lymphomatosis cerebri, putamen, striatum

(Intern Med 60: 623-627, 2021)

(DOI: 10.2169/internalmedicine.5200-20)

Introduction

Lymphomatosis cerebri (LC) is a rare form of primary central nervous system lymphoma (PCNSL) that predominantly involves the white matter. Pathological examinations of LC patients demonstrate the diffuse infiltration of lymphoma cells into the nerve fibers (1). Immunohistochemical studies have revealed diffuse large B-cell lymphoma in most LC patients (2). However, why LC patients present with diffuse infiltration without mass formation remains unclear.

On magnetic resonance imaging (MRI), LC is characterized by diffuse, non-enhancing infiltrative lesions without mass formation or mass effect (1-9). The lesions of LC are likely to be most clearly demonstrated with a fluid-attenuated inversion recovery (FLAIR) sequence (2, 3). The lesions also show a high signal intensity on diffusion-weighted imaging (DWI), and most of the lesions exhibit an

increase in apparent diffusion coefficient (ADC) values on an ADC map (4), which may reflect the diffuse infiltration of non-cohesive malignant lymphoid cells (4). The lesions of LC are found in the cerebral white matter in 77.8%, corpus callosum in 42.2%, basal ganglia in 33.3%, thalamus in 26.7%, midbrain in 46.7%, pons in 44.5%, medulla oblongata in 31.1%, cerebellum in 26.6%, and spinal cord in 11.1% of patients (3). The cerebral cortices, cerebellar cortices, and cerebellar dentate nuclei are also rarely but occasionally involved (3).

However, the distribution of deep gray matter lesions on MRI characteristic of LC has not been reported. We herein report the distribution of MRI of deep gray matter lesions on the FLAIR sequence of MRI in two patients with LC and review the literature.

¹Department of Neurology, JA Toride Medical Center, Japan, ²Department of Neurology, Kanto Central Hospital, Japan, ³Aozora Clinic, Japan, ⁴Department of Neurology, Nakatsu Municipal Hospital, Japan, ⁵Department of Neurology and Neurological Science, Tokyo Medical and Dental University, Japan, ⁶Department of Neurosurgery, JA Toride Medical Center, Japan and ⁷Department of Hematology, JA Toride Medical Center, Japan

Received: April 30, 2020; Accepted: August 4, 2020; Advance Publication by J-STAGE: September 30, 2020

Correspondence to Dr. Zen Kobayashi, kobayashiz112@yahoo.co.jp

Figure 1. Fluid-attenuated inversion recovery (FLAIR) sequence of brain MRI (A-H, J-L) demonstrated a high signal intensity in the brainstem (A-F), left cerebellum (B, C), bilateral hippocampus (D, E), bilateral thalamus (G, H), bilateral globus pallidus (G, H, J), bilateral putamen (G, H, J), and bilateral cerebral white matter (F-H, J-L). The letter “R” indicates the right side. The putamen and caudate head (striatum) were relatively spared. Gadolinium-enhanced T1-weighted image (Gd-T1WI) (M-P) showed abnormal enhancement from the left hippocampus (M, arrow) to the globus pallidus (N-P, arrow).

Case Reports

Case 1

A 67-year-old woman gradually presented with bradykinesia, followed by appetite loss and difficulty walking, and was admitted to our hospital about 1 month after the onset. She had no medical history of note. A neurological examination demonstrated a mask-like face, dysarthria, dysphagia, bradykinesia, muscle rigidity, frontal lobe sign, and pyramidal sign as well as a wide-based and short-stepped gait.

On MRI, bilateral brain lesions were most clearly demonstrated with a FLAIR sequence, which showed a high signal intensity in the brainstem, cerebellum, hippocampus, thalamus, putamen, globus pallidus, and cerebral white matter (Fig. 1). Among the deep gray matter, the caudate head and putamen (striatum) were relatively spared when compared with the globus pallidus, thalamus, and hippocampus. There was no apparent mass effect. On DWI, the lesions showed a faint high signal intensity. In an ADC map, most lesions exhibited a slight increase of the ADC values. A gadolinium-

enhanced T1-weighted image (Gd-T1WI) showed abnormal enhancement from the right hippocampus to the globus pallidus (Fig. 1).

Blood tests demonstrated a soluble interleukin 2 receptor (sIL-2R) level of 264 U/mL (reference range: 145-519 U/mL). A lumbar puncture yielded watery clear cerebrospinal fluid (CSF). A CSF examination showed a cell count of 5/mm³, total protein of 50 mg/dL (reference range: 8-43 mg/dL), glucose of 65 mg/dL, sIL-2R of 63.1 U/mL (reference range: <85), and β 2 microglobulin of 5.50 mg/L (reference range: 0.69-1.29). Cytology of the CSF was negative.

A stereotactic brain biopsy of the right globus pallidus demonstrated the diffuse infiltration of large abnormal cells, which were immunopositive for CD20 and CD79a. A diagnosis of LC due to diffuse large B-cell lymphoma was established based on the MRI and biopsy findings.

Case 2

A 71-year-old man presented with progressive gait disorder and was admitted to our hospital about 5 months after the onset. His medical history included hypertension and chronic kidney disease. A neurological examination demon-

Figure 2. FLAIR sequence of brain MRI (A-H, J-M) demonstrated a high signal intensity in the brainstem (A-H), bilateral cerebellum (C, D), bilateral hippocampus (E-G), bilateral thalamus (J, K), bilateral caudate head (K), bilateral globus pallidus (K), bilateral putamen (L), and bilateral cerebral white matter (E-H, J-M). The striatum was relatively spared. Gd-T1WI (N-Q, S-V) showed abnormal enhancement around the inferior horn of the left lateral ventricle (N-Q) and in the genu of the corpus callosum (T-V).

strated time-related disorientation, miscalculation, frontal lobe sign, and upper limb apraxia as well as limb and truncal ataxia.

On MRI, bilateral brain lesions were most clearly demonstrated with a FLAIR sequence, which showed a high signal intensity in the brainstem, cerebellum, hippocampus, thalamus, caudate head, putamen, globus pallidus, and cerebral white matter (Fig. 2). Among the deep gray matter, the caudate head and putamen (striatum) were relatively spared when compared with the globus pallidus, thalamus, and hippocampus. There was no apparent mass effect. Gd-T1WI showed abnormal enhancement around the inferior horn of the left lateral ventricle and in the corpus callosum (Fig. 2).

Blood tests showed that the sIL-2R level was slightly elevated to 723 U/mL. A lumbar puncture yielded watery clear

CSF. The cell count was 9/ μ L, comprising mononuclear cells of 8/ μ L and polymorphonuclear cells of 1/ μ L. The total protein and glucose levels of the CSF were 51 and 61 mg/dL, respectively. The sIL-2R level of the CSF was elevated to 676 U/mL. Cytology of the CSF was negative.

A stereotactic brain biopsy from the genu of the corpus callosum demonstrated the diffuse infiltration of large abnormal cells with an angiocentric pattern, which were immunopositive for CD10, CD20, and CD79a. A diagnosis of LC due to diffuse large B-cell lymphoma was established based on the MRI and biopsy findings.

Discussion

On MRI of our patients, the FLAIR sequence clearly

Table. The Differential Diagnosis of Lymphomatosis Cerebri.

	Clinical course	MRI findings		
		Bilateral hemispheric lesions on the first MRI scan	Mass effect	Relative sparing of the striatum despite involvement of other deep gray matter on FLAIR
Posterior reversible encephalopathy syndrome	acute	common (posterior predominant)	none (or slight)	uncommon
Progressive multifocal leukoencephalopathy	subacute	uncommon	none (or slight)	uncommon
Lymphomatosis cerebri	subacute (or chronic)	common	none (or slight)	sometimes present
Glioma	subacute (or chronic)	uncommon	present	uncommon

Figure 3. Klüver-Barrera staining (myelin staining) of a coronal section of the human postmortem brain hemisphere at the plane of the internal segment of the globus pallidus shows that the striatum is hardly stained in comparison with other deep gray matter tissues, including the globus pallidus, thalamus, and hippocampus.

demonstrated bilateral diffuse brain lesions, which showed no apparent mass effect. Enhancement of the lesions on Gd-T1WI was shown only in a small area. These MRI findings were consistent with LC. The differential diagnosis of LC is shown in Table. Izquierdo et al. reported that 95.2% of LC patients have bilateral hemispheric lesions on the first MRI scan (2), and this finding is uncommon in other diseases considered in the differential diagnosis (2).

In our patients, the caudate head and putamen (striatum) were relatively spared on MRI compared with the globus pallidus, thalamus, and hippocampus. Interestingly, we found seven previous reports of similar MRI findings with relative sparing of the striatum despite involvement of other deep gray matter in patients with LC (2, 3, 5-9). This finding may be due to the fact that the nerve fibers are sparse in the striatum when compared with other deep gray matter. Fig. 3 shows the Klüver-Barrera staining (myelin staining) of a coronal section of the human postmortem brain hemisphere at the plane of the internal segment of the globus pallidus (an autopsy of this another patient was done at our hospital). This figure shows that the striatum is hardly stained compared with other deep gray matter tissues, in-

cluding the globus pallidus, thalamus, and hippocampus.

In conclusion, in addition to the finding of “bilateral hemispheric lesions on the first MRI scan (2)”, “relative sparing of the striatum despite involvement of other deep gray matter on FLAIR” may be characteristic of LC and provide a clue for the diagnosis of LC, although the absence of these findings does not necessarily exclude the possibility of LC. Further investigations of a larger number of LC patients are necessary to confirm these findings.

The authors state that they have no Conflict of Interest (COI).

Acknowledgement

The authors thank Takami Itoh, Yukinori Inadome, and Daisuke Kobayashi for contributing to the diagnosis and medical care of the patients.

References

- Bakshi R, Mazziotta JC, Mischel PS, Jahan R, Seligson DB, Vinters HV. Lymphomatosis cerebri presenting as a rapidly progressive dementia: clinical, neuroimaging and pathologic findings. *Dement Geriatr Cogn Disord* **10**: 152-157, 1999.
- Izquierdo C, Velasco R, Vidal N, et al. Lymphomatosis cerebri: a rare form of primary central nervous system lymphoma. Analysis of 7 cases and systematic review of the literature. *Neuro Oncol* **18**: 707-715, 2016.
- Li L, Rong JH, Feng J. Neuroradiological features of lymphomatosis cerebri: a systematic review of the English literature with a new case report. *Oncol Lett* **16**: 1463-1474, 2018.
- Yu H, Gao B, Liu J, et al. Lymphomatosis cerebri: a rare variant of primary central nervous system lymphoma and MR imaging features. *Cancer Imaging* **17**: 26, 2017.
- Raz E, Tinelli E, Antonelli M, et al. MRI findings in lymphomatosis cerebri: description of a case and revision of the literature. *J Neuroimaging* **21**: e183-e186, 2011.
- Imperiale D, Taraglio S, Atzori C, Testi R. Diffuse leukoencephalopathy due to lymphomatosis cerebri: a clinicopathological report. *Neurol Sci* **36**: 1071-1073, 2015.
- Kitai R, Hashimoto N, Yamate K, et al. Lymphomatosis cerebri: clinical characteristics, neuroimaging, and pathological findings. *Brain Tumor Pathol* **29**: 47-53, 2012.
- Yamaguchi N, Matsuda S, Yoshizawa T, Shigeeda R, Inoue T, Uchibori Y. A case of lymphomatosis cerebri rapidly confirmed by brain biopsy. *Rinsho Shinkeigaku (Clin Neurol)* **59**: 286-289, 2019 (in Japanese, Abstract in English).
- Watanabe M, Satoi H, Takahashi Y, Nishida N, Toda H, Matsumoto S. Remission of lymphomatosis cerebri induced by

corticosteroid and high-doses intravenous methotrexate. Rinsho Shinkeigaku (Clin Neurol) **52**: 486-490, 2012 (in Japanese, Abstract in English).

The Internal Medicine is an Open Access journal distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

© 2021 The Japanese Society of Internal Medicine
Intern Med 60: 623-627, 2021