[CASE REPORT]

High-output Heart Failure Caused by a Tumor-related Arteriovenous Fistula: A Case Report and Literature Review

Akihiro Tobe¹, Akihito Tanaka¹, Satoya Yoshida¹, Toru Kondo¹, Ryota Morimoto¹,
Kenji Furusawa¹, Takahiro Okumura¹, Yasuko K Bando¹,
Hideki Ishii^{1,2} and Toyoaki Murohara¹

Abstract:

High-output heart failure caused by a tumor-related arteriovenous fistula in adults is a rare clinical condition. We herein report a case of high-output heart failure caused by an arteriovenous fistula associated with renal cell carcinoma and a literature review of 29 published cases to date. Renal cell carcinoma seems to be the most common underlying tumor. For the diagnosis, right heart catheterization and enhanced computed tomography are considered useful. The removal of the underlying tumor and arteriovenous fistula is the best treatment for heart failure.

Key words: high-output heart failure, arteriovenous fistula, tumor, renal cell carcinoma

(Intern Med Advance Publication)

(DOI: 10.2169/internalmedicine.6962-20)

Introduction

High-output heart failure caused by an arteriovenous fistula associated with neoplasm in adults is rare but results in a critical condition. Its epidemiology, treatment, and clinical course, as well as the type of underlying tumor have not been fully elucidated. We herein report a case of high-output heart failure caused by an arteriovenous fistula associated with renal cell carcinoma along with a literature review.

Case Report

A 64-year-old man with lower leg edema and dyspnea on exertion visited a local doctor. His blood pressure and pulse rate were 145/79 mmHg and 93 beats per minute, respectively. Chest radiography showed cardiac enlargement with a cardiothoracic ratio (CTR) of 69%, bilateral pleural effusion, and nodule shadows in the right lung (Fig. 1A). Computed tomography (CT) incidentally revealed a right renal mass.

He was referred to the urology department of our hospital.

The results of a right kidney biopsy indicated clear cell renal cell carcinoma. Enhanced CT revealed a giant hypervascular mass arising from the inferior pole of the right kidney extending to the renal vein and inferior vena cava (IVC), as well as multiple metastases in the lungs. Taken together, these results suggested a clinical stage of T3b N0 M1 according to the TNM classification system (Fig. 2A-C).

Neo-adjuvant chemotherapy was started; however, the cardiac enlargement and lower leg edema worsened, and the patient was finally referred to a cardiologist. A systolic heart murmur was heard, but abdominal auscultation was not performed. Transthoracic echocardiography revealed mild hypokinesis with a left ventricular ejection fraction of 45%, 4-chamber dilatation, moderate mitral valve regurgitation, and severe tricuspid valve regurgitation with an estimated right ventricular pressure of 40 mmHg (Table 1). The patient was diagnosed with congestive heart failure due to unknown cause and received more intensive treatment including oral diuretics (tolvaptan 15 mg/day, furosemide 60 mg/day, and

¹Department of Cardiology, Nagoya University Graduate School of Medicine, Japan and ²Department of Cardiology, Fujita Health University Bantane Hospital, Japan

Received: December 21, 2020; Accepted: February 12, 2021; Advance Publication by J-STAGE: March 29, 2021

Correspondence to Dr. Akihito Tanaka, akihiro17491194@gmail.com

Figure 1. Transition of chest X-ray findings. CTR: cardiothoracic ratio, BW: body weight, NT-proBNP: N-terminal pro-brain natriuretic peptide, BNP: brain natriuretic peptide

Figure 2. Enhanced CT findings in the arterial phase. A: Axial plane. Arrowheads indicate a hypervascular mass in the right kidney. B: Coronal plane. The proximal part of the IVC is enlarged and enhanced. C: Axial plane. Multiple veins drain into the IVC. D: An enlarged view of dotted area of the picture C. An arrow indicates the renal cancer extending to the IVC. IVC: inferior vena cava, pIVC: proximal part of IVC, dIVC: distal part of IVC, Ao: aorta, #: renal artery, *: veins draining into IVC

spironolactone 25 mg/day) and an intravenous diuretic (furosemide 20 mg/day). However, dyspnea on exertion, cardiomegaly, and a high level of brain natriuretic peptide (BNP; 259 pg/mL) persisted (Fig. 1B). Right heart catheterization showed a high cardiac output of 8.68 L/min, a cardiac index of 4.72 L/min/m² (Fick), and slightly elevated pressure in each chamber (Table 1). Blood gas sampling showed high

oxygen saturation at the proximal site to renal veins in the IVC (90.3%), suggesting an arteriovenous shunt in the systemic circulation. Coronary angiography revealed normal coronary arteries. Reassessment of enhanced CT images revealed enhancement of the right renal veins and IVC in the arterial phase, indicating an arteriovenous fistula (Fig. 2D). The patient was finally diagnosed with high-output heart

Table 1. Echocardiographic and Right Heart Catheterization Data before and after Surgery.

	Before surgery	After surgery
<i>Echocardiography</i>		
Interventricular septum thickness, mm	8.3	8.9
Posterior left ventricular wall thickness, mm	9.7	8.9
Left ventricular end-diastolic dimension, mm	59.8	53.7
Left ventricular end-systolic dimension, mm	44.7	40.8
Left ventricular ejection fraction, %	49.06	47.41
Stroke volume, ml	87.63	66.14
Cardiac output, L/min	6.40	4.43
Left atrial dimension, mm	57.0	46.9
IVC, mm	35.4	12.6
Mitral regurgitation	Moderate	Mild-moderate
Tricuspid regurgitation	Severe	Moderate
Estimated right ventricular pressure, mmHg	40.0	30.1
<i>Right heart catheterization</i>		
Blood pressure (systolic/diastolic/mean), mmHg	109/59/82	122/60/84
Heart rate, /min	73	60
Cardiac output, L/min (Fick method)	8.68	4.92
Cardiac output, L/min (Thermo method)	10.97	5.59
Cardiac index, L/min/m ² (Fick method)	4.72	2.79
Cardiac index, L/min/m ² (Thermo method)	5.96	3.18
Pulmonary capillary wedge pressure, mmHg	16	7
Mean Pulmonary artery pressure, mmHg	26	17
Right atrial pressure, mmHg	12	4
Stroke volume, mL	122	82
Systemic vascular resistance, dyne-sec/cm ⁵	645	1301
Pulmonary vascular resistance, dyne-sec/cm ⁵	92	163
Oxygen saturation of IVC, %	90.3	83.9
Oxygen saturation of SVC, %	61.4	72.2
Oxygen saturation of RA, %	82.2	75.3

IVC: inferior vena cava, SVC: superior vena cava, RA: right atrium

failure due to an arteriovenous fistula associated with renal cell carcinoma.

Radical nephrectomy with a lower midline abdominal incision was performed. During the surgery, the central venous pressure was 12-16 mmHg before ligation of the renal artery, and it decreased to 7-8 mmHg immediately after the ligation. Multiple enlarged veins draining directly into the IVC were observed and then ligated and dissected, along with the renal artery. After clamping the IVC, main right renal vein was excised, and the right kidney was removed together with the extending part in the IVC. A histopathological examination confirmed clear cell renal cell carcinoma.

The postoperative course was uneventful. The cardiomegaly and lower leg edema dramatically improved, and the dose of diuretics was gradually reduced. Treatment was thus discontinued 13 days after the surgery. Postoperative right heart catheterization after the discontinuation of diuretics showed normalization of the cardiac output, pressure, and oxygen saturation in the IVC (Table 1).

At the 1-month follow-up visit, the patient was doing well. Chest radiography showed a CTR of 46%, and he had a BNP level of 28 pg/mL without diuretics (Fig. 1C). He

then received postoperative adjuvant chemotherapy.

Discussion

We herein report a case of high-output heart failure caused by an arteriovenous fistula associated with renal cell carcinoma, which can be considered a tumor-related heart disease. An arteriovenous fistula is known to cause high-output heart failure by increasing the venous return; (1) however, a tumor-related arteriovenous fistula causing heart failure is uncommon. Table 2 summarizes previously published cases of heart failure caused by a tumor-related arteriovenous fistula in the systemic circulation (2-30).

Underlying tumors

In the majority of published cases, the tumors causing heart failure due to an arteriovenous fistula in the systemic circulation were renal cancers (22 out of 30 cases) (2-4, 6, 7, 9, 11-19, 21, 22, 24, 25, 28, 29). including mostly renal cell carcinoma. Cases of clear cell carcinoma, which is the most common type of renal cell carcinoma, frequently involve a mutated von Hippel-Lindau tumor suppressor gene.

Table 2. Summary of Previously Published Cases Representing Heart Failure Caused by Tumor-related Arteriovenous Fistula.

Author (Year)	Age	Sex	CC	Cardiac murmur	Bruit of tumor	HT	Location of tumor	Type of tumor	Diagnosis of AVF	Treatment of tumor	Course of HF
Hamilton (1953) ²	29	m	HF	+	+	+	kidney	adenocarcinoma	bruit	surgery	improved
Scheifley (1959) ³	39	f	Both	+	+	+	kidney	hypernephroma	bruit, AG	surgery	improved
Abbot (1961) ⁴	62	f	Tumor	+	+	+	kidney	renal cell carcinoma	bruit, AG	surgery	improved
Crawford (1963) ⁵	51	m	HF	+	+	+	Ao and IVC	malignant mesenchymoma	bruit, AG	surgery	improved
Nicoloff (1964) ⁶	74	f	Both	-	+	+	kidney	hypernephroma	bruit, AG	surgery	improved
Thomas (1965) ⁷	60	m	Tumor	+	+	+	kidney	clear cell carcinoma	bruit, AG	surgery	improved
Vetto (1966) ⁸	73	m	Tumor	-	+	+	scapula, pelvis	renal cell carcinoma	bruit, AG, RHC	surgery	improved
Norris (1971) ⁹	39	m	Tumor	-	+	+	bilateral kidneys	adenocarcinoma	bruit, AG, RHC	surgery, RT	improved
Damascelli (1972) ¹⁰	33	f	Tumor	ND	+	ND	thyroid, lymph node	small medullary carcinoma	bruit, AG	surgery	improved
Sondag (1973) ¹¹	67	f	HF	+	-	+	kidney	hypernephroma	AG	chemo	ND
Pickens (1973) ¹²	51	f	HF	+	+	+	kidney	clear cell carcinoma	RHC, bruit, AG	surgery	improved
Curtiss (1974) ¹³	38	f	Tumor	+	-	+	kidney	clear cell carcinoma	AG	surgery	improved
Rodgers (1975) ¹⁴	63	f	HF	+	+	+	kidney	clear cell carcinoma	RHC, LVG	surgery	improved
Bass (1975) ¹⁵	60	m	Tumor	-	ND	+	kidney	clear cell carcinoma	AG	surgery	improved
Ricketts (1978) ¹⁶	53	f	Tumor	+	+	ND	kidney	possible hypernephroma	bruit, AG	TAE	improved
Crawford (1979) ¹⁷	45	f	HF	ND	+	+	bilateral kidneys	renal cell carcinoma	bruit, AG	surgery	improved
Thomas (1981) ¹⁸	63	f	Tumor	ND	ND	ND	bilateral kidneys	possible hypernephroma	AG	chemo	got worse
Nakada (1983) ¹⁹	62	m	Tumor	ND	-	-	kidney	clear cell carcinoma	AG	surgery	improved
Waki (1984) ²⁰	59	m	HF	ND	+	ND	liver	hepatocellular carcinoma	AG	TAE	improved
Selli (1986) ²¹	40	f	Tumor	+	ND	-	kidney	renal cell carcinoma	sonogram, AG	surgery	improved
Holmes (1991) ²²	63	m	HF	ND	ND	+	kidney	renal cell carcinoma	AG	surgery	improved
Lee (1993) ²³	44	f	Tumor	ND	ND	ND	intrapelvis	leiomyoma	TEE, AG	surgery	improved
Takahashi (1995) ²⁴	76	m	Tumor	ND	ND	+	kidney, pharynx	clear cell carcinoma	AG	surgery	improved
Piggott (2000) ²⁵	62	f	HF	+	-	-	kidney	renal cell carcinoma	infer	surgery	improved
Szucs-Farkas (2002) ²⁶	47	m	Both	ND	ND	ND	iliac artery	malignant fibrous histiocytoma	eCT, AG	surgery	improved
Ozaki (2003) ²⁷	53	f	HF	+	+	-	intrapelvis	possible alveolar soft part sarcoma	bruit, eCT, RHC, AG	-	no change
Hayek (2014) ²⁸	58	m	HF	+	ND	+	kidney	clear cell carcinoma	RHC, MRI, AG	surgery	improved
Watanabe (2015) ²⁹	64	f	Both	ND	+	-	kidney	clear cell carcinoma	eCT	TAE, surgery	improved
Ando (2018) ³⁰	78	f	HF	ND	ND	+	iliac artery	leiomyosarcoma	eCT	surgery	ND
Tobe (2020)	64	m	HF	+	ND	+	kidney	clear cell carcinoma	RHC, eCT	surgery	improved

Listed reports are written in English or Japanese. Such cases are not listed: patients' age<18, pulmonary arteriovenous fistula, hypertension or cardiomegaly without heart failure, cardiac tumors, hematological malignancies, angioma or angiosarcoma.

CC: chief complaint, HF: heart failure-related symptoms, Tumor: tumor-related symptoms, Both: both heart failure- and tumor- related symptoms, ND: not described, HT: hypertension, AVF: arteriovenous fistula, AG: angiography, RHC: right heart catheterization, LVG: left ventricular angiogram, TEE: transesophageal echocardiography, eCT: enhanced computed tomography, MRI: magnetic resonance imaging, RT: radiation therapy, TAE: transcatheter arterial embolization

The gene inhibits angiogenesis and cell growth (31), so most renal cell carcinoma cases show hypervascularity. Microscopically, vascular invasion and microembolism of neoplastic cells are suspected of causing vascular fragility, resulting in arteriovenous connection (29, 32). Furthermore, renal cell carcinoma commonly creates spaces that are cov-

ered in a layer of endothelium and directly communicating with both capsular arteries and veins (33). These features may lead to the formation of an arteriovenous fistula. Other underlying tumor types, such as unclassified malignant mesenchymoma (5), small medullary carcinoma of the thyroid (10), hepatocellular carcinoma (20), leiomyoma (23),

malignant fibrous histiocytoma (26), and possible alveolar soft part sarcoma (27) and leiomyosarcoma (30), have also been reported. Arteriovenous fistulas associated with tumors might not be uncommon; the incidence of the involvement of arteriovenous fistula in renal cell carcinoma is reported to be 12%-50% (34, 35). However, it seems quite rare for congestive heart failure to develop because of this. The type or location of tumors accompanied by arteriovenous fistulas that cause heart failure might thus be limited.

Symptoms at the first visit

The chief complaint at the first visit was heart failure-related symptoms (2, 5, 11, 12, 14, 17, 20, 22, 25, 27, 28, 30), symptoms related to underlying tumors (4, 7-10, 13, 15, 16, 18, 19, 21, 23, 24), and both (3, 6, 26, 29) in 13, 13, and 4 previously published cases, respectively. About half of the patients presented with symptoms associated with heart failure at the first visit. Renal cancer, which is a major underlying cause, is often asymptomatic and it tends to be found incidentally. The famous triad of hematuria, flank pain, and a palpable mass is no longer a common presentation, and such a condition therefore suggests the presence of advanced stage disease (36). Therefore, heart failure-related symptoms might appear as the first symptoms. When a patient presents with high-output heart failure due to unknown cause, we may need to consider a tumor-associated arteriovenous fistula.

The diagnosis

• Physical examinations

In most cases, an arteriovenous fistula bruit and/or a systolic heart murmur were heard (2-14, 16, 17, 20, 21, 25, 27-29). A systolic heart murmur indicates relative outflow tract obstruction caused by the high-output state. The bruit and murmur disappeared after resection of the underlying tumors and arteriovenous fistulas. Arteriovenous fistulas that cause high-output heart failure might have a large shunt flow, resulting in an audible arteriovenous fistula bruit. However, the bruit was not detected in some of the previous cases (11, 13, 19, 25), and the absence of bruit is not always indicative of the absence of an arteriovenous fistula. Systolic hypertension, which also results from an arteriovenous fistula because of an increase in cardiac output (37), was observed in most cases (2-9, 11-15, 17, 22, 24, 28, 30).

• Right heart catheterization

Preoperative right heart catheterization was performed in 12 cases (5, 8, 9, 12-14, 16, 17, 23, 27, 28), and follow-up right heart catheterization was performed in 6 cases after treatment (5, 9, 13, 16, 28). The pre-treatment cardiac output was 8.14-15.2 L/min (5, 9, 13, 17, 27, 28), and an increase in the oxygen saturation level was noted in 8 cases (5, 8, 9, 12, 14, 27, 28). Interestingly, right heart catheterization revealed the presence of an arteriovenous fistula in some cases (14). The post-treatment follow-up data showed a decrease in cardiac output in all reported cases (5, 9, 13,

16, 28).

• Imaging examinations

Imaging is essential to confirm the presence of an arteriovenous fistula. In the past, angiography was frequently performed, but more recently, enhanced CT has been most commonly used. Magnetic resonance imaging (MRI) and echography can be also helpful.

To detect a tumor-related arteriovenous fistula, bruit over the site of the tumor and systolic heart murmur are helpful findings. To confirm the presence of an arteriovenous fistula, angiography was performed in the past, but enhanced CT or MRI is currently recommended. For a definitive diagnosis of arteriovenous shunt and the assessment of heart failure, right heart catheterization is helpful.

Treatment and clinical course

Resection of the underlying tumor was performed in most cases. After the removal of tumors and arteriovenous fistula, heart failure improved in almost all cases. In two cases, embolization of the arteriovenous fistula was performed, which resulted in the improvement of heart failure (16, 20). In one case, radiation therapy brought temporary relief of heart failure symptoms (9). When treating malignant tumor, direct therapies, including resection, embolization, or radiation, are usually considered, and these can also be used to treat heart failure. Furthermore, even if the tumor is benign and does not require invasive treatment for itself, we should consider performing such therapies in patients with heart failure that is uncontrollable by medical therapy due to severe tumor-related arteriovenous fistula.

Conclusion

High-output heart failure caused by a tumor-related arteriovenous fistula in adults is rare, but its incidence may not be negligible. A careful examination can lead to the correct and prompt diagnosis. Renal cancer, mainly renal cell carcinoma, seems to be the most common cause of underlying tumors. The removal of the underlying tumor and arteriovenous fistula is the best treatment for heart failure.

Author's disclosure of potential Conflicts of Interest (COI).

Takahiro Okumura: Honoraria, Ono Pharmaceutical Co., Ltd., Medtronic, Pharma K. K., and Otsuka Pharma Ltd.; Research funding, Ono Pharmaceutical Co., Bayel Pharmaceutical Co., Ltd., Daiichi-Sankyo Co., Ltd. and Astellas Pharma Inc. Hideki Ishii: Honoraria, Astellas Pharma Inc., Astrazeneca Inc., Daiichi-Sankyo Pharma Inc. and MSD K. K. Yasuko Bando: Honoraria, Daiichi-Sankyo Co., Ltd., MSD K. K., Mitsubishi Tanabe Pharma Co., Nippon Boehringer Ingelheim Co., Ltd., Pfizer Japan Inc., Eily Lily Co., Ltd. and Bristol-Squibb Mayer Co., Ltd. Toyooki Murohara: Honoraria, Bayel Pharmaceutical Co., Ltd., Daiichi-Sankyo Co., Ltd., Dainippon Sumitomo Pharma Co., Ltd., Kowa Co., Ltd., MSD K. K., Mitsubishi Tanabe Pharma Co., Nippon Boehringer Ingelheim Co., Ltd., Novartis Pharma K. K., Pfizer Japan Inc., Sanofi-aventis K. K. and Takeda Pharmaceutical Co., Ltd.; Research funding, Astellas Pharma Inc.,

Daiichi-Sankyo Co., Ltd., Dainippon Sumitomo Pharma Co., Ltd., Kowa Co., Ltd., MSD K. K., Mitsubishi Tanabe Pharma Co., Nippon Boehringer Ingelheim Co., Ltd., Novartis Pharma K. K., Otsuka Pharma Ltd., Pfizer Japan Inc., Sanofi-aventis K. K., Takeda Pharmaceutical Co., Ltd. and Teijin Pharma Ltd.

References

- Reddy YNV, Melenovsky V, Redfield MM, Nishimura RA, Borlaug BA. High-Output Heart Failure: A 15-Year Experience. *J Am Coll Cardiol* **68**: 473-482, 2016.
- Hamilton GR, Getz RJ, Jerome S. Arteriovenous fistula of the renal vessels; case report and review of the literature. *J Urol* **69**: 203-207, 1953.
- Scheifley CH, Daugherty GW, Greene LF, Priestley JT. Arteriovenous fistula of the kidney; new observations and report of three cases. *Circulation* **19**: 662-671, 1959.
- Abbott CE, Poutasse EF. Renal arteriovenous fistula: occurrence in renal-cell carcinoma. Report of a case. *Cleve Clin Q* **28**: 283-288, 1961.
- Crawford ES, Turell DJ, Alexander JK. Aorto-inferior vena caval fistula of neoplastic origin. Hemodynamic and coronary blood flow studies. *Circulation* **27**: 414-421, 1963.
- Nicoloff DM. Renal Arteriovenous Fistula: Occurrence in Renal Cell Carcinoma. *Am J Surg* **108**: 82-84, 1964.
- Thomas CV, Feint JF, Nayman J. An arteriovenous shunt associated with an adenocarcinoma of the kidney. *Br J Surg* **52**: 943-946, 1965.
- Vetto RM, Bigelow JC, Dueger WC Jr. Arteriovenous fistula in pulsatile metastatic carcinoma. Report of chronic heart failure relieved by resection. *Am Surg* **32**: 557-560, 1966.
- Norris AD, Murray M, Mantell BS, Singh M. Carcinoma of both kidneys presenting as arteriovenous fistulae. *Br J Urol* **43**: 395-398, 1971.
- Damascelli B, Preda S, LaMonica G, Veronesi U. Giant arteriovenous fistula in thyroid tumour inducing cardiac failure--selective angiography. *Br J Radiol* **45**: 531-534, 1972.
- Sondag TJ, Patel SK, Petasnick JP, Chambliss J. Hypernephromas with massive arteriovenous fistulas. *Am J Roentgenol Radium Ther Nucl Med* **117**: 97-103, 1973.
- Pickens S. Hypernephroma presenting as cardiomegaly. *Br Med J* **3**: 678-679, 1973.
- Curtiss EI, Shaver JA, Boehnke MA. Left to right shunt due to arteriovenous fistula formation in a renal cell carcinoma. *Arch Intern Med* **134**: 951-953, 1974.
- Rodgers MV, Moss AJ, Hoffman M, Lipchik EO. Arteriovenous fistulae secondary to renal cell carcinoma. Clinical and cardiovascular manifestations: report of a case. *Circulation* **52**: 345-350, 1975.
- Bass EM, Lipper MH. Malignant renal arteriovenous fistulae. A cause of reversible cardiac failure. *S Afr Med J* **49**: 1361-1362, 1975.
- Ricketts RR, Finck E, Yellin AE. Management of major arteriovenous fistulas by arteriographic techniques. *Arch Surg* **113**: 1153-1159, 1978.
- Crawford ED, Skinner DG, Mallamo JT. Renal arteriovenous fistulas secondary to bilateral renal cell carcinoma. *Urology* **13**: 525-528, 1979.
- Thomas JL, Bernardino ME. Neoplastic-induced renal vein enlargement: sonographic detection. *AJR Am J Roentgenol* **136**: 75-79, 1981.
- Nakada G, Machida T, Masuda F, et al. [A case of arteriovenous fistulae secondary to renal cell carcinoma accompanied by congestive heart failure]. *Hinyokika Kyo* **29**: 901-905, 1983.
- Waki M, Neyatani H, Yoshida M, Kitanaka H. [Arteriovenous shunt with congestive heart failure in primary hepatocellular carcinoma: case report]. *Rinsho Hoshasen* **29**: 1513-1515, 1984.
- Selli C, Bartolozzi C, Lizzadro G, Petacchi D. Arteriovenous fistula associated with renal cell carcinoma: demonstration by magnetic resonance imaging. *Urol Radiol* **8**: 190-193, 1986.
- Holmes SA, Ball AJ. Arteriovenous fistula associated with adenocarcinoma of the kidney. *Urol Int* **47**: 81-83, 1991.
- Lee VS, Thompson NW, Cho KJ, Goldblum JR. High-output cardiac failure: an unusual manifestation of intravenous leiomyomatosis. *Surgery* **113**: 466-470, 1993.
- Takahashi Y, Uno H, Kuriyama M, Miyata H, Shimokawa K, Kawada Y. [Pharyngeal metastasis and arteriovenous fistula of renal cell carcinoma--report of a case]. *Gan To Kagaku Ryoho* **22**: 133-136, 1995.
- Piggott M, Farrugia D, Otite U, Oliver RT, Crake T. Renal cell carcinoma presenting with cardiac failure. *Hosp Med* **61**: 804-805, 2000.
- Szucs-Farkas Z, Toth J, Szollosi Z, Peter M, Bartha I. Pseudoaneurysm and ilio-caval fistula caused by malignant fibrous histiocytoma of the aorta--CT diagnosis and angiographic confirmation. *Eur Radiol* **12**: 450-453, 2002.
- Ozaki K, Oda H, Hatada K, et al. Highly vascular pelvic tumor causing high-output heart failure because of massive arteriovenous shunting: a case report. *Circ J* **67**: 554-555, 2003.
- Hayek S, Kung R, Barb I, Master V, Al S, Clements S. Digging deep: high output heart failure in renal cell carcinoma. *Am J Med* **127**: 22-24, 2014.
- Watanabe D, Horiguchi A, Isono M, et al. [Renal Cell Carcinoma Presenting with High-Output Heart Failure Due to Arteriovenous Fistula]. *Nihon Hinyokika Gakkai Zasshi* **106**: 35-39, 2015.
- Ando T, Goto H, Date K, Takeda M. Surgical Experience of a Case of Primary Leiomyosarcoma of the Left Common Iliac Artery That Presented as Acute Heart Failure Involving an Arteriovenous Fistula. *Ann Vasc Dis* **11**: 243-247, 2018.
- Cohen HT, McGovern FJ. Renal-cell carcinoma. *N Engl J Med* **353**: 2477-2490, 2005.
- Maldonado JE, Sheps SG, Bernatz PE, Deweerd JH, Harrison EG Jr. Renal Arteriovenous Fistula. A Reversible Cause of Hypertension and Heart Failure. *Am J Med* **37**: 499-513, 1964.
- Ichijo S. Vascular Architecture of Renal Adenocarcinoma. *Tohoku J Exp Med* **102**: 273-281, 1970.
- Imagawa A, Yuasa M. [Arteriovenous fistula associated with renal cell carcinoma of the kidney (author's transl)]. *Nihon Hinyokika Gakkai Zasshi* **67**: 103-106, 1976.
- Onishi T, Masuda F, Sasaki T, et al. [Clinical evaluation of 6 arteriovenous fistulae secondary to renal cell carcinoma (author's transl)]. *Nihon Hinyokika Gakkai Zasshi* **73**: 316-325, 1982.
- Gray RE, Harris GT. Renal Cell Carcinoma: Diagnosis and Management. *Am Fam Physician* **99**: 179-184, 2019.
- Chobanian AV. Clinical practice. Isolated systolic hypertension in the elderly. *N Engl J Med* **357**: 789-796, 2007.

The Internal Medicine is an Open Access journal distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).