[CASE REPORT]

An Autopsy Case of Disseminated Varicella Zoster Virus Infection During the Treatment of Nephrotic Syndrome

Katsuhisa Kuwano¹, Yuki Ota¹, Kiyokazu Tsuji¹, Kenta Torigoe¹, Ayuko Yamashita¹, Kumiko Muta¹, Mineaki Kitamura¹, Hiroshi Yamashita¹, Tadashi Uramatsu¹, Masato Tashiro², Hiroko Hayashi³, Koichi Izumikawa², Hiroshi Mukae⁴ and Tomoya Nishino¹

Abstract:

A 68-year-old woman developed systemic blisters while receiving treatment for nephrotic syndrome. As she also developed marked liver dysfunction and disseminated intravascular coagulation, she was admitted to our hospital. She was diagnosed with varicella zoster virus (VZV) infection. Treatment was administered in the intensive-care unit, but the patient died on day 24 post-admission after severe VZV infection. A post-mortem examination showed micro-abscesses and necrosis caused by varicella zoster infection in multiple organs, including the liver, kidneys, and gastrointestinal tract. Because VZV infection can become severe in immunocompromised patients, careful consideration is needed for the prevention and treatment of the viral infection.

Key words: varicella zoster virus, nephrotic syndrome, immunosuppressive therapy

(Intern Med Advance Publication)

(DOI: 10.2169/internalmedicine.7332-21)

Introduction

Herpes zoster is caused by the reactivation of varicella zoster virus (VZV), which remains latent in the dorsal root ganglia. Herpes zoster is highly prevalent in Japan, with an occurrence rate of 4.15 per 1,000 individuals (1). Herpes zoster is more common in older adults than in younger ones. Those with reduced T-cell immunity (e.g. recipients of an organ transplant or hematopoietic stem cells; patients receiving immunosuppressive therapy; and those with lymphoma, leukemia, or HIV infection) are at high risk of exacerbation (2).

In general, a patient with herpes zoster infection develops blisters on one side of the body along a single nerve pathway. Disseminated herpes zoster, which is a rash with disseminated lesions, may lead to visceral complications (3). Among patients who have undergone hematopoietic stem cell transplantation, 10% of those who develop disseminated

herpes zoster die (4). However, the mortality rate in those with renal disease is unknown.

We herein report a recent autopsy case of disseminated herpes zoster infection, in which the deceased patient had been receiving immunosuppressive therapy for nephrotic syndrome.

Case Report

A 68-year-old woman developed generalized blisters while receiving treatment for nephrotic syndrome. As she also developed marked liver dysfunction, renal dysfunction, and disseminated intravascular coagulation, she was admitted to our hospital as an emergency case. Three years earlier, she had shown the onset of nephrotic syndrome associated with cryoglobulinemic vasculitis and received treatment for remission with 30 mg/day of prednisolone (PSL) and cryofiltration. Thereafter, as proteinuria improved, the PSL dose was tapered to 5 mg/day, and the patient received

¹Department of Nephrology, Nagasaki University Hospital, Japan, ²Department of Infectious Diseases, Nagasaki University Graduate School of Biomedical Sciences, Japan, ³Department of Pathology, Nagasaki University Graduate School of Biomedical Sciences, Japan and ⁴Department of Respiratory Medicine, Nagasaki University Graduate School of Biomedical Sciences, Japan

Received: February 14, 2021; Accepted: June 29, 2021; Advance Publication by J-STAGE: August 13, 2021

Correspondence to Dr. Yuki Ota, yukiota.nagasaki@gmail.com

Table. Laboratory Findings.

[Blood test]		TP	3.4 g/dL	VZV IgG (EIA)	55.8
WBC	6,800 / μ L	Alb	1.6 g/dL	VZV IgM (EIA)	0.89
RBC	266 $\times$ 10 ⁴ / μ L	BUN	35 mg/dL	VZV-DNA (EIA)	4.0 $\times$ 10 ⁶ copies/mL
Hb	8.8 g/dL	Cr	1.31 mg/dL	[Urinalysis]	
Hct	24 %	eGFR	31.8 mL/min/1.73m ²		
PLT	7.7 $\times$ 10 ⁴ / μ L	Na	132 mEq/L		
PT-INR	3.0	K	4.7 mEq/L		
APTT	88.9 sec	Cl	86 mEq/L		
FDP	177 μ g/mL	Glu	78 mg/dL		
D-dimer	74.3 μ g/mL	LDL-C	122 mg/dL		
T.Bil	1.6 mg/dL	CRP	1.67 mg/dL		
AST	2,670 U/L	IgG	280 mg/dL		
ALT	747 U/L	IgA	97 mg/dL		
ALP	820 U/L	IgM	107 mg/dL		
γ GTP	257 U/L	C3	20 mg/dL		
LDH	6,713 U/L	C4	3.2 mg/dL		
CK	199 IU/L				
				Protein	(3+)
				Glucose	(-)
				Urobilinogen	(-)
				Bilirubin	(-)
				Ketone	(-)
				Occult blood	(3+)
				RBC	>100 /HPF
				WBC	10-19 /HPF

Alb: albumin, ALT: alanine transaminase, ALP: alkaline phosphatase, AST: aspartate aminotransferase, APTT: activated partial thromboplastin time, BUN: blood urea nitrogen, Cl: chloride, Cr: creatinine, CK: creatine kinase, CRP: C-reactive protein, eGFR: estimated glomerular filtration rate, FDP: fibrin/fibrinogen degradation products, γ GTP: gamma-glutamyl-transferase, Glu: glucose, Hb: hemoglobin, Hct: hematocrit, IgA: immunoglobulin A, IgG: immunoglobulin G, IgM: immunoglobulin M, K: potassium, LDL-C: low density lipoprotein Cholesterol, LDH: lactate dehydrogenase, Na: sodium, PLT: platelet, PT-INR: prothrombin time-international normalized ratio, RBC: red blood cell, T.Bil: total bilirubin, TP: total protein, VZV: varicella zoster virus, WBC: white blood cell

maintenance therapy. Three months earlier, the patient had had recurrence of nephrotic syndrome and been admitted to the previous hospital where she received steroid pulse therapy (methylprednisolone 500 mg/day, 3 days). Thereafter, treatment with 30 mg/day of PSL and 100 mg/day of mizoribine (MZR) had been commenced.

Two days before admission to our hospital, she developed diffuse blisters on the right side of her body. Thereafter, she experienced severe acute chest pain, and the blisters spread throughout her body. The patient tested positive on the Tzanck test and was diagnosed with herpes infection. Treatment with acyclovir (ACV) 1,200 mg/day was initiated. Our hospital took over the treatment.

The patient's physical findings on admission were as follows: Glasgow Coma Scale score, E3V4M5; body height, 148 cm; body weight, 39 kg; blood pressure, 81/56 mmHg; heart rate, 119/min; percutaneous oxygen saturation (SpO₂), 80% (10 L/min of oxygen); and body temperature, 38.7 °C. A large number of half-rice-grain-sized to red-bean-sized blisters accompanied by purple erythema were observed throughout the patient's body. They included blood blisters and purpura. Bilateral coarse crackles were heard, and generalized edema was also observed. Laboratory findings showed that the patient had anemia, thrombocytopenia, abnormal coagulation, liver dysfunction, renal dysfunction, and hypoalbuminemia as well as high VZV-DNA levels (Table).

As the blister lesions tested positive for the VZV antigen, we considered the cause of organ injury to be disseminated herpes zoster infection. Urinalysis results indicated proteinuria (urinary plasma creatinine ratio: 6.68 g/gCr) and hema-

turia (>100/high-power field). Thoracoabdominal contrast-enhanced computed tomography showed an enlarged liver, spleen, and kidneys. Organ injury was observed, as indicated by inhomogeneous contrast enhancement. Pleural effusion and ascites were present, and consolidation accompanied by an air bronchogram was observed in the bilateral lower lobes (Fig. 1).

The patient was in a condition of shock with respiratory failure and disseminated intravascular coagulation. Mechanical ventilation via endotracheal intubation was initiated. In addition, as the patient developed anuric acute kidney injury, continuous hemodiafiltration was commenced, and the patient was admitted to the intensive-care unit. A bone marrow test showed marked hemophagocytic findings, which we considered to be hemophagocytic lymphohistiocytosis syndrome (HLS) accompanying severe viral infection. We continued the administration of ACV, which had been started at the previous hospital. As the patient was suspected of having pneumonia-associated sepsis, empirical treatment was also commenced with the broad-spectrum antibacterial agents meropenem and vancomycin. No additional treatments, such as immunosuppressant therapy, were provided for cryoglobulinemic vasculitis because treatment of severe infection was prioritized. MZR was discontinued, and PSL was gradually reduced and then discontinued. Following treatment initiation, the blisters crusted, but the patient's general condition remained poor. ACV was administered at a dosage of 1,200 mg/day, but the dose was reduced to 250 mg/day due to crusting of the blisters. Pancytopenia due to HLS progressed, and the patient required red blood cell and plate-

Figure 1. Computed tomography findings. The liver, spleen, and kidneys were found to be enlarged, and inhomogeneous contrast enhancement was seen. Pleural effusion and ascites were present, and consolidation accompanied by air bronchogram was observed in the bilateral lower lobes.

Figure 2. Clinical course. ACV: acyclovir, MEPM: meropenem, VCM: vancomycin, mPSL: methylprednisolone, IVIG: Intravenous immunoglobulin, WBC: white blood cell, Hb: hemoglobin, PLT: platelet, AST: aspartate aminotransferase, ALT: alanine transaminase, LDH: lactate dehydrogenase, CRP: C-reactive protein, VZV: varicella zoster virus

let transfusion over the course of several days. She died on post-admission day 24 (Fig. 2).

A post-mortem examination was conducted after obtaining consent from the patient's family. Infiltration of inflammatory cells and coagulative necrosis accompanied by intranuclear inclusion bodies were observed in the liver; findings of pulmonary edema as well as micro-abscesses, which were considered the foci of disseminated viral infection, were also observed. In addition, severe neutrophil infiltration was seen in the left pleurae, suggesting purulent pleuritis. Findings of myocardial necrosis accompanied by infiltration of inflammatory cells were observed in the heart; micro-abscesses and necrosis were also seen in the spleen and pancreas. Destructive lesions of the glomeruli accompanied by fibrin deposition and acute tubular necrosis were observed in the kidneys (Fig. 3). Furthermore, ulcers, which were considered to have been caused by viral infection, were observed in the stomach and both the ascending and descend-

ing colon. Hemophagocytosis and necrotic foci were observed in the bone marrow.

Taken together, the above findings indicated VZV infection-induced necrosis of almost all the viscera. Although VZV immunostaining was not performed, we considered these observations to be pathological findings associated with VZV infection because VZV viremia and nuclear inclusion bodies were observed. The cause of death was therefore established as multiple organ failure from disseminated herpes zoster.

Discussion

This patient developed disseminated herpes zoster while receiving immunosuppressive therapy for nephrotic syndrome. Her condition worsened to multiple organ failure, which led to her death. The present case suggests that VZV infection should be considered when treating a patient with

Figure 3. Pathological anatomical findings. a) Lungs (macroscopic findings). Left lung 395 g, right lung 520 g (left is right lung). The cut surface is yellowish-brown with edema and hyperemia. Blood and fibrin retention can be observed between the lobes in the left lung (arrow). b) Lungs (microscopic findings). In the bilateral lungs, pulmonary edema was found, and the foci of fibrin precipitation and inflammatory cell infiltration were partially observed as shown in the photograph (left: $\times 20$). Eosinophilic intranuclear inclusion bodies were observed in a small number of macrophages with halos around them (arrows, central: $\times 40$). Multinucleated giant cells with eosinophilic intranuclear inclusion bodies were also observed (right: $\times 40$). c) Liver (macroscopic findings). The liver weighed 785 g. Necrosis with a yellowish-brown tone was observed. d) Liver (microscopic findings). Confluent necrosis is observed, which accounts for 20%-30% of the total liver. The necrosis was surrounded by cells showing ground-glass-like nuclei (left: $\times 4$, right: $\times 40$). e) Spleen (macroscopic and microscopic findings). The spleen weighed 110 g. Adhesion of yellowish-brown moss-like material is observed on the film. Necrosis can be observed on the cut surface (arrow). Extensive necrosis and blistering were observed in the spleen ($\times 20$). f) Kidneys (macroscopic and microscopic findings). Left kidney 146 g, right kidney 132 g (the right kidney is shown on the left). The cortex is reddish-brown, and the medulla is greenish-brown. In the glomerulus, a crescent formation was observed. It was covered with cells showing ground-glass-like nuclei, and an inclusion body-like structure was also observed (arrow, $\times 40$).

kidney disease who requires immunosuppressive therapy as carefully as patients with blood disorders and transplantation. This case study is novel because, to our knowledge, there have been no reports of disseminated VZV infection in patients with cryoglobulinemic vasculitis.

While the epidemiology and prognosis of disseminated herpes zoster is widely recognized in the fields of blood disease and transplantation (5, 6), it is not widely recognized in nephrology. Despite the importance of being adequately informed on herpes zoster in order to perform disease treatment, few nephrologists have sufficient knowledge (7). The onset and exacerbation of herpes zoster is associated with reduced T-cell immunity (2). When treating kidney diseases, including nephrotic syndrome and glomerulonephritis, drugs that suppress T cell immunity, such as corticosteroids, are primarily used. Therefore, treatment using such drugs may increase the risk of VZV infection. Mizoribine, an immunosuppressant, also selectively suppresses the proliferation of lymphocytes by inhibiting the purine synthesis pathway of nucleic acids. Therefore, it was considered that the drug had a negative impact on the patient's immunity by inhibiting the proliferation of T and B cells.

We confirmed disseminated lesions caused by VZV in a post-mortem examination. There have been similar post-mortem studies on patients who died from disseminated herpes zoster infection (8-13). However, few reports on post-mortems of patients with kidney disease who died from disseminated herpes zoster infection have been published. In our patient, a pathological analysis showed VZV infection-induced micro-abscesses and necrosis in the liver, lungs, pleurae, heart, stomach, colon, kidneys, spleen, pancreas, and bone marrow. The patient's VZV-DNA content in the blood was also extremely high. It was considered that the virus spread through the body via the bloodstream, resulting in various organ injuries. In particular, the liver was severely necrotic, which may pathologically explain the marked elevation of hepatic enzymes. Furthermore, broad necrosis was observed in the spleen. In the lungs, eosinophilic inclusion bodies were observed in a small number of macrophages with halos around them. In the kidneys, crescent formation, glomerular necrosis, and fibrin deposition were observed. Intracellular inclusion body-like structures were also observed. These findings were therefore considered to be pathological findings related to VZV infection rather than those of cryoglobulinemic vasculitis, which was the underlying disease. The central nervous system was not evaluated in the present study. However, as the patient exhibited impaired consciousness, she might have developed VZV infection-induced encephalomyelitis and meningitis.

Among patients with herpes zoster, those who are ≤ 50 years old and have moderate to severe rash or pain, facial and ocular lesions, and acute infective complications and are immunodeficient are at a high risk of exacerbation. Aggressive antiviral therapy is indicated for such patients, and it is recommended that therapy be commenced within 72 h after the onset (2). Early symptoms of disseminated herpes zoster

include severe pain in the abdomen, back, and lower back caused by visceral infection. These symptoms may appear prior to rash. Given the high mortality rate, early detection is key to providing effective treatment. In the present case, the patient had severe chest pain before developing generalized blisters. The patient's risk factors were old age and immune deficiency. In addition, treatment was commenced after more than 72 h had passed since the symptom onset. These factors were considered to be the causes of the VZV-induced progression observed in the visceral lesions.

VZV loads remained abnormally high even after ACV therapy was commenced. As ACV resistance was observed during treatment, the patient might have had ACV-resistant VZV infection. Studies have reported the effectiveness of foscarnet for ACV-resistant VZV infection, although the drug has not been approved for the treatment of herpes zoster in Japan (2, 14).

To prevent the onset and worsening of herpes zoster, vaccine inoculation against the disease is also important. At present, the available herpes zoster vaccines are recombinant zoster vaccine (RZV) and zoster vaccine live (ZVL). The Advisory Committee on Immunization Practices recommends that RZV be administered to adults ≥ 50 years old with normal immunocompetence, irrespective of the presence of comorbidities (15, 16). As ZVL is a live vaccine, it is, in principle, contraindicated for patients with compromised immunity. However, one study reported the safety of the live vaccine in patients with nephrotic syndrome receiving immunosuppressive therapy (17). Whether or not the patient in the present case had received a varicella vaccine was unclear. In children with steroid-sensitive nephrotic syndrome, live vaccines should be avoided if prednisone doses are 1 mg/kg daily (<20 mg/day) or 2 mg/kg every other day (<40 mg every other day), in which case corticosteroid-sparing immunosuppressive agents and varicella zoster globulin should be administered in case of contact with VZV infection for nonimmune children on immunosuppressive agents (18). No consensus has been reached concerning adult cases. When treating a patient with kidney disease, it is necessary to ask them about their vaccination history and consider inoculation against herpes zoster prior to immunosuppressive therapy.

We reported an autopsy case of disseminated herpes zoster. The patient died from the disease while receiving immunosuppressive therapy for nephrotic syndrome. Patients with severely compromised immunity, including those who have undergone hematopoietic stem cell transplantation, may develop life-threatening disseminated herpes zoster. This can also occur in immunocompromised renal disease patients. Nephrologists may often consider preventing common bacterial (e.g. *Mycobacterium tuberculosis*) and fungal infections, but they are rarely aware of viral infection. It is important to detect the disease at an early stage, so patients exhibiting chest or abdominal pain and presenting with blisters while receiving immunosuppressive therapy should be closely evaluated for the possibility of herpes zoster infection.

The authors state that they have no Conflict of Interest (COI).

Acknowledgements

We thank the laboratory staff at the Department of Pediatrics, Nagasaki University Hospital, for performing real-time polymerase chain reaction.

Authorship statement: TN was responsible for the organization and coordination of the report. KK and YO drafted the manuscript. KK, YO, HY, TU, MT and KI were involved in the treatment of the patient. HH helped describe the anatomy of the patient and analyze the pathological findings. KT, KT and AY contributed to the acquisition of the data. KI, HM and TN helped treat the patient and write the manuscript. All authors contributed to the writing of the final manuscript.

References

1. Toyama N, Shiraki K, Amano M, et al. Epidemiology of herpes zoster and its relationship to varicella in Japan: A 10-year survey of 48,388 herpes zoster cases in Miyazaki Prefecture. *J Med Virol* **81**: 2053-2058, 2009.
2. Cohen JI. Herpes Zoster. *N Engl J Med* **369**: 255-263, 2013.
3. Locksley RM, Flournoy N, Sullivan KM, Meyers JD. Infection with varicella-zoster virus after marrow transplantation. *J Infect Dis* **152**: 1172-1181, 1985.
4. Atkinson K, Meyers JD, Storb R, Prentice RL, Thomas ED. Varicella-zoster virus infection after marrow transplantation for aplastic anemia or leukemia. *Transplantation* **29**: 47-50, 1980.
5. Koc Y, Miller KB, Schenkein DP, et al. Varicella zoster virus infections following allogeneic bone marrow transplantation: Frequency, risk factors, and clinical outcome. *Biol Blood Marrow Transplant* **6**: 44-49, 2000.
6. Rommelaere M, Maréchal C, Yombi JC, Goffin E, Kanaan N. Disseminated varicella zoster virus infection in adult renal transplant recipients: Outcome and risk factors. *Transplant. Proc* **44**: 2814-2817, 2012.
7. Furuto Y, Kawamura M, Namikawa A, Takahashi H, Shibuya Y. Successful management of visceral disseminated varicella zoster virus infection during treatment of membranous nephropathy: A case report. *BMC Infect Dis* **19**: 5-9, 2019.
8. Tsuji H, Yoshifuji H, Fujii T, Matsuo T, Nakashima R, Imura Y, et al. Visceral disseminated varicella zoster virus infection after rituximab treatment for granulomatosis with polyangiitis. *Mod Rheumatol* **27**: 155-161, 2017.
9. Yamashiro N, Nagasaka T, Ooishi N, et al. An autopsy case of meningoencephalitis and cerebral infarction that developed with Ramsay Hunt syndrome and disseminated herpes zoster. *Brain and Nerve* **70**: 253-258, 2018.
10. Matsuda Y, Nonaka K, Seki A, et al. Disseminated varicella zoster virus infection and acute hepatitis: Autopsy-proven cases from a single institute. *Pathol Int* **69**: 378-380, 2019.
11. Akiyama M, Yoshifuji K, Fukuda T, et al. Fulminant visceral disseminated varicella-zoster virus infection without skin involvement in a patient with autoimmune hemolytic anemia on prednisolone therapy. *Rinsho Ketsueki* **57**: 467-471, 2016.
12. Mori M, Imamura Y, Maegawa H, Yoshida H, Naiki H, Fukuda M. Cytology of pleural effusion associated with disseminated infection caused by varicella-zoster virus in an immunocompromised patient: A case report. *Acta Cytol* **47**: 480-484, 2003.
13. Yamazaki E, Kamijoh A, Taguchi J, et al. Fatal acute visceral disseminated varicella-zoster virus infection in a patient with chronic graft-versus-host disease. *Rinsho Ketsueki* **45**: 1053-1057, 2004.
14. Safrin S, Berger TG, Gilson I, et al. Foscarnet therapy in five patients with AIDS and acyclovir-resistant varicella-zoster virus infection. *Ann Intern Med* **115**: 19-21, 1991.
15. Dooling KL, Guo A, Patel M, et al. Recommendations of the Advisory Committee on Immunization Practices for Use of Herpes Zoster Vaccines. *MMWR Morb Mortal Wkly Rep* **67**: 103-108, 2018.
16. Freedman MS, Hunter P, Ault K, Kroger A. Advisory Committee on Immunization Practices Recommended Immunization Schedule for adults aged 19 years or older - United States, 2020. *MMWR Morb Mortal Wkly Rep* **69**: 133-135, 2020.
17. Kamei K, Miyairi I, Ishikura K, et al. Prospective study of live attenuated vaccines for patients with nephrotic syndrome receiving immunosuppressive agents. *J Pediatr* **196**: 217-222.e1, 2018.
18. Radhakrishnan J, Cattran DC. The KDIGO practice guideline on glomerulonephritis: Reading between the (guide) lines-application to the individual patient. *Kidney Int* **82**: 840-856, 2012.

The Internal Medicine is an Open Access journal distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).