[CASE REPORT]

Juvenile Hepatocellular Carcinoma in a Healthy Liver: A Case Report

Sachiyo Onishi¹, Masahiro Tajika¹, Tsutomu Tanaka¹, Keisaku Yamada¹, Tomoyasu Kamiya¹,
Seiji Natsume², Yasuhiro Shimizu² and Yasumasa Niwa¹

Abstract:

Primary hepatocellular carcinoma (HCC) in patients <30 years old is extremely rare. In younger patients, HCC develops against a background of persistent hepatitis B virus infection. We herein report a 23-year-old woman with HCC with all-negative hepatitis virus markers developing in an apparently healthy liver. Imaging studies showed a 50-mm hypervascular mass in segment 4 of the left liver lobe, compatible with HCC. The patient underwent surgical resection. A histological examination showed the presence of poorly differentiated HCC. The patient was diagnosed with HCC developing in a healthy liver. This is an extremely rare case of non-B non-C HCC.

Key words: hepatocellular carcinoma, non-B non-C, healthy liver, juvenile hepatocellular carcinoma

(Intern Med Advance Publication)

(DOI: 10.2169/internalmedicine.8305-21)

Introduction

In Japan, primary hepatocellular carcinoma (HCC) often develops against a background of chronic hepatitis, resulting from persistent infection with hepatitis B virus (HBV) or hepatitis C virus (HCV). Individuals over 50 years old are considered the most susceptible. However, the onset of primary HCC at a younger age is often associated with persistent HBV infection (1).

We herein report a 23-year-old woman with primary HCC who showed no findings of chronic hepatitis and for whom all hepatitis-related markers remained consistently negative.

Case Report

A 23-year-old healthy Japanese woman had normal vaginal delivery at 21 years old. One year before this presentation, she noticed an abdominal mass, visited a local doctor, and underwent a follow-up examination. On admission because of dysuria, a tumor was detected by abdominal ultrasonography (US) in her upper abdomen, and she was referred to our hospital for a further examination of the mass.

She had no family history of HCC or any other malignancy. Furthermore, she was a non-smoker and non-drinker, with no history of medications including pills or supplements. The patient also had no history of blood transfusion or trauma, such as hepatic trauma. She did not have any metabolic abnormalities, including her thyroid function based on the findings obtained either at the clinic of her own obstetrician at the time of her first child's birth or at our hospital.

On visiting our hospital, her physical characteristics were as follows: height, 159 cm; weight, 35 kg; and body mass index (BMI), 13.8 kg/m². She had a soft abdomen with no palpable liver or spleen, no abdominal distension, no tenderness, and no edema. Blood test results showed normal results for liver function tests (Table 1). However, some tumor markers were elevated: alpha fetoprotein, 941.8 ng/mL; protein induced by vitamin K absence-II, 1,170 mAU/mL. These findings suggested HCC (Table 1). All HBV markers, including HBsAg, HBsAb, and HBcAb, were negative (Table 1). Results for anti-HCV antibody were also negative. Serum concentrations of IgG, IgM, copper, and ferritin were all normal (Table 1). The laboratory findings for the patient were negative for antinuclear antibody and anti-mitochondrial antibody (Table 1). The albumin level, total

¹Department of Endoscopy, Aichi Cancer Center Hospital, Japan and ²Department of Gastroenterological Surgery, Aichi Cancer Center Hospital, Japan

Received: July 11, 2021; Accepted: September 2, 2021; Advance Publication by J-STAGE: October 19, 2021

Correspondence to Dr. Sachiyo Onishi, soonishi@aichi-cc.jp

Table 1. Results of Laboratory Blood Examination on Admission.

TP	7.8 g/dL	AFP	941.8 ng/mL
Albumin	4.8 g/dL	AFP-L3	1.80 %
BUN	11 mg/dL	PIVKA-II	1,170 mAU/mL
Cre	0.49 mg/dL	CEA	0.5 ng/mL
AST	18 U/L	CA19-9	13.2 U/mL
ALT	12 U/L	HBsAg	(-)
γ -GTP	16 U/L	HBsAb	(-)
ALP	216 U/L	HBcAb	(-)
T-bil	0.6 mg/dL	HCVAb	(-)
LDH	161 U/L	IgG	1,179 mg/dL
Na	140 mmol/L	IgM	294 mg/dL
K	3.8 mmol/L	ANA	<40
Cl	106 mmol/L	AMA	<20
ChE	263 U/L		
HbA1c	5.3 %	Fe	35.8 ng/mL
WBC	9,320 / μ L	Cu	107 μ g/dL
RBC	498 $\times$ 10 ⁴ / μ L	TC	204 mg/dL
Hb	13.8 g/dL	TG	55 mg/dL
Plt	26.4 $\times$ 10 ⁴ / μ L		
PT-INR	1.06 %		
APTT	42.2 s		

TP: total protein, BUN: blood urea nitrogen, Cre: creatinine, AST: aspartate aminotransferase, ALT: alanine aminotransferase, γ -GTP: γ -glutamyl transferase, ALP: alkaline phosphatase, T-bil: total bilirubin, LDH: lactate dehydrogenase, ChE: cholinesterase, WBC: white blood cell, RBC: red blood cell, Hb: hemoglobin, Plt: platelet, PT-INR: prothrombin time-international normalized ratio, APTT: activated partial thromboplastin time, AFP: alpha-fetoprotein, PIVKA-II: protein induced by vitamin K absence or antagonist II, HBsAg: hepatitis B surface antigen, HBsAb: hepatitis B surface antibody, HBcAb: hepatitis B core antibody, HCVAb: hepatitis C virus antibody, ANA: antinuclear antibody, AMA: anti-mitochondrial antibody, TC: total cholesterol, TG: triglyceride

bilirubin level, prothrombin time, and indocyanine green (ICG) R15 value were within normal ranges, suggesting a good hepatic reserve (Table 1).

Abdominal US showed a solitary hypoechoic tumor with clear boundaries in segment 4 of the left liver lobe. Abdominal computed tomography (CT) showed a 5-cm tumor in S4 of the liver (Fig. 1a). Multiphase CT showed a strong uptake of the contrast agent in the hepatic arterial phase (Fig. 1b), followed by washout in the portal venous phase (Fig. 1c) and the equilibrium phase (Fig. 1d). However, CT showed no findings of a central scar. Magnetic resonance imaging (MRI) showed decreased and increased signal intensities for the lesion on T1- and T2-weighted imaging, respectively, whereas diffusion restriction with a high signal intensity and a low apparent diffusion coefficient were apparent on diffusion-weighted imaging. Dynamic contrast-enhanced MRI showed strong enhancement in the hepatic artery phase with subsequent washout in the portal venous and equilibrium phases and typical hypointensity in the hepatobiliary phase in segment 4 of the liver. Based on these findings, HCC of cT2N0M0 cStage II (Union for International Cancer Control Seventh Edition, UICC 7th edition)

was diagnosed.

The patient underwent surgical resection of segment 4, as her hepatic reserve was considered adequate for resection. The operation and postoperative clinical course were uneventful, and the patient was discharged on postoperative day 7.

A postoperative examination of the tumor specimen showed a 50 $\times$ 30-mm lesion with expansive growth (Fig. 2a). A histological examination of the non-tumorous portions of the specimen showed no fibrosis (Fig. 2b), and no infiltration of inflammatory cells into the portal areas or hepatic lobules was evident (Fig. 2b). No fatty deposits were identified in hepatocytes (Fig. 2b). The pathological report indicated poorly differentiated HCC, as follows: fc (capsule formation) (+), fc-inf (infiltration) (+), sf (septum formation) (+), s0, n0, Vp1, Vv1, Va0, b0, p0, sm (-) pT2 N0 M0 p Stage II (UICC, 7th edition) (Fig. 2c).

She gave birth to her second child 15 months after surgery, and at the time of writing this manuscript, at 5 years post-operation, she has had no recurrence of the disease. We conducted follow-up investigations such as blood tests, including tumor marker evaluations, every month, abdominal US every three months, and abdominal CT every six months.

Discussion

Approximately 80% of primary liver cancers are HCC, and 80% of all HCC patients have chronic hepatitis or liver cirrhosis with persistent HBV or HCV infection and a history of alcohol intake (1-6). Recently, the incidence of non-B non-C-type HCC, which may be due to alcoholic hepatitis, nonalcoholic steatohepatitis, hemochromatosis, primary biliary cirrhosis (PBC), or drugs such as oral contraceptive pill (7-11), has been reported to be increasing. In Japan, HCV infection often causes chronic hepatitis or liver cirrhosis, and HCC caused by HCV often develops in middle-aged patients (12, 13). The frequency of HCC in patients <35 years old is 0.89%, and only 0.14% are <25 years old (6, 12). In carcinogenesis of HCC in young patients, the roles of HBV DNA integration or the HBV HBx protein have been reported (14, 15). HCC that occurs unrelated to hepatitis or liver cirrhosis is extremely rare (3.2%) in Japan (12). Fibrolamellar HCC (FLC) is a special type of liver cancer (16, 17) that develops in young patients and is unrelated to hepatitis or liver cirrhosis. It is characterized by a large tumor with calcification and a central scar (17, 18).

In the present case, the patient was a young woman, at only 23 years old. Furthermore, in this case, hepatitis virus-related markers such as HBsAg, HBsAb, HBcAb, HCVAb were all negative, and neither HBV-DNA nor HCV-RNA was detected. In the postoperative examination of pathological specimens, the background liver appeared normal and no findings suggestive of chronic hepatitis were detected and the pathological findings were also different from FLC. As far as we search in Pubmed and medical journal 'Igaku

Figure 1. Multiphase computed tomography (CT) on admission showing juvenile hepatocellular carcinoma that developed in a normal liver. a) Image taken before the administration of contrast medium shows a tumor in the S4 segment of the liver. b-d) Findings of contrast-enhanced CT. A solid hepatic mass is seen in the S4 segment of the liver. The uptake of the contrast agent in the hepatic arterial phase (b) is strong, with subsequent washout in the portal venous phase (c) and the equilibrium phase (d).

Figure 2. A postoperative examination shows a tumor 50×30 mm in size with expansive growth. a) Resected tumor. b) Hematoxylin and Eosin (H&E) staining of the noncancerous portions showing no fibrosis and no inflammatory infiltration to the portal areas and hepatic lobules. c, d) H&E staining of the tumor showing partial fatty changes, a solid nest structure, and a false duct structure, indicating poorly differentiated hepatocellular carcinoma. The tumor cells show invasion into a capsule with partial vascular invasion.

Table 2. Juvenile Hepatocellular Carcinoma without Hepatitis Infection: Summary of Case Reports.

No.	Age (years)	Sex	HBsAg/ Ab	HBcAb	HBV-DNA	HCV-Ab	Family history	Past history	Drinking/ smoking
1	Fujita ²² (1995)	25	F	-/-	-	-	-	Mycoplasma pneumonia	-/-
2	Sato ²³ (1996)	23	F	-/-	-	-	Chronic hepatitis	Congenital cataract	-/-
3	Churiki ⁸ (1997)	17	M	-/-	-	-	-	Bone fracture	-/-
4	Nakano ²⁴ (2002)	28	M	-/-	ND	-	-	-	-/+
5	Fukui ²⁵ (2011)	20	M	-/-	-	-	-	-	ND/ND
6	Kaibori ²⁶ (2007)	20	M	-/-	-	-	-	Precocious puberty	ND/ND
7	Okamoto ²⁷ (2010)	25	F	-/-	-	ND	ND	Epilepsy	ND/ND
8	Kitagawa ²⁸ (2014)	25	F	-/-	-	-	-	Ovarian cyst	-/-
9	Carla ²⁹ (2009)	26	F	-/-	-	ND	ND	ND	ND
10	Tanaka ³⁰ (1996)	39	F	-/-	-	-	ND	ND	-/-
11	Herath ³¹ (2017)	41	M	-/-	-	-	-	-	-/-
12	Sanada ³² (2004)	31	F	-/-	-	-	ND	Pyogenic spondylitis	ND
13	Kubo ³³ (2007)	31	M	-/-	-	-	-	Appendicitis	+/-
14	Zubin ³⁴ (2014)	40	F	-/-	ND	-	-	-	-/ND
15	Sunagawa ³⁵ (2013)	26	M	-/-	-	-	-	-	+/+
16	Abe ³⁶ (2019)	30	M	-/-	-	-	-	Asthma	-/ND
17	Hashimoto ³⁷ (2021)	20's	M	-/-	-	-	-	-	-/-
18	Onishi	23	F	-/-	-	-	-	-	-/-

No.		Transfusion recording	Drug	AFP	PIVKaII	Treatment	Recurrence	Outcome (after treatment)
1	Fujita ²² (1995)	-	ND	35,200	14.4	TAE	No recurrence	Alive 11 months
2	Sato ²³ (1996)	-	-	10,320	Normal	TAE	Brain metastasis (2 years)	Died 2 years
3	Churiki ⁸ (1997)	+	-	<3.0	<1.0	TAE, Ope	No recurrence	Alive 19 months
4	Nakano ²⁴ (2002)	-	-	37,000	18,100	ND	No recurrence	Died 16 days
5	Fukui ²⁵ (2011)	-	-	181.8	75,000	Ope, TAE, PEIT	Intrahepatic (7 months)	Alive 36 months
6	Kaibori ²⁶ (2007)	ND	ND	25,100	292	Ope	No recurrence	Alive 30 months
7	Okamoto ²⁷ (2010)	ND	Phenytoin	5,011	6,022	TAE, Ope	No recurrence	Alive 21 months
8	Kitagawa ²⁸ (2014)	-	-	1.2	329	Ope	No recurrence	Alive 108 months
9	Carla ²⁹ (2009)	ND	ND	>300	ND	Palliative chemo	No recurrence	Died 3 months
10	Tanaka ³⁰ (1996)	-	-	Normal	Normal	Ope	ND	ND
11	Herath ³¹ (2017)	-	-	ND	ND	BSC	ND	Died 3 months
12	Sanada ³² (2004)	ND	-	4.1	39	Ope	No recurrence	Alive 31 months
13	Kubo ³³ (2007)	-	-	4.8	41	Ope	ND	Alive 24 months
14	Zubin ³⁴ (2014)	+	ND	ND	ND	BSC	ND	Died 8 days
15	Sunagawa ³⁵ (2013)	-	-	Normal	Normal	Ope	ND	ND
16	Abe ³⁶ (2019)	-	-	2.3	24	ope	-	Alive 10 months
17	Hashimoto ³⁷ (2021)	ND	-	76,860	74,849	Chemotherapy	ND	Died 9 months
18	Onishi	-	-	941.8	1,170	Ope	-	Alive 60 months

M: male, F: female, ND: not described, TAE: transcatheter arterial embolization, Ope: operation, PEIT: percutaneous ethanol injection therapy, BSC: best supportive care

Chuo Zasshi (ICHUSHI)' up to 2021, only 17 cases of non-B non-C HCC without chronic hepatitis or liver cirrhosis in the liver have been reported (Table 2). Although approximately half of those cases involved patients who were under thirty years old, only two cases showed HCC without previous history, family history, blood transfusion history, drinking history or oral administration other than this patient. The patient was definitively underweight and BMI was low. However, her weight did not change preoperatively, and she had not experienced any changes in her body shape during her life. Further, blood testing revealed no evidence of malnutrition, and no evidence of fatty liver from malnutrition was seen in the background liver on postoperative pathology, so we do not believe that underweight status affected the development of HCC in this case. Therefore, we consider this case as extremely rare.

Recently, methods for genome-wide association analyses have advanced dramatically, and some reports have provided genomic information for HCC. However, HCC is a heterogeneous disease, with characteristics not only due to the cause, but also due to the genetic characteristics of the tumor, as the process of multicentric carcinogenesis differs between individual cases. In addition to identifying gene mutations for whole-genome sequencing, viral integration can also be identified. In recent years, not only the HBV genome, but also the adeno-associated virus type 2 genome has been reported to become integrated into HCC, and many cases without known risk factors have been reported. In our case, genome sequencing thus appears important for clarifying the genetic characteristics of carcinogenesis.

In general, HCC that develops in young patients often progresses and shows a low degree of differentiation on histology (19, 20). Recurrence within one year or pathological vascular invasion at the time of initial resection is said to be a factor contributing to poor prognosis at the time of recurrence (21) and follow-up within the short term using various modalities is desirable after surgical resection. In our case, vascular invasion (Vp1, Vv1) was seen in the postoperative pathological tissue, so the risk of recurrence was considered high. We conducted follow-up observations such as blood tests including tumor markers every month, abdominal ultrasonography every three months, and abdominal CT every 6 months. As of 5 years after surgery, she is undergoing follow-up with no recurrence, and we will continue follow-up observation in the future.

In conclusion, we experienced present an extremely rare case of non-B non-C HCC occurred in normal liver in at 23 years old. No recurrence has been observed for 5 years, however, careful follow-up is required due to the risk of recurrence.

The authors state that they have no Conflict of Interest (COI).

References

- Herbst DA, Reddy KR. Risk factors for hepatocellular carcinoma. *Clin Liver Dis* **1**: 180-182, 2012.
- European Association, For The, Study Of, The Liver. European Organisation For Research And Treatment Of Cancer. EASL-EORTC clinical practice guidelines: management of hepatocellular carcinoma. *J Hepatol* **56**: 908-943, 2012.
- Tomasz K. Nowicki, Karolina Markiet, Edyta Szurowska. Diagnostic Imaging of Hepatocellular Carcinoma - A Pictorial Essay. *Curr Med Imaging Rev* **13**: 140-153, 2017.
- Forner A, Llovet JM, Bruix J. Hepatocellular carcinoma. *Lancet* **379**: 1245-1255, 2012.
- El-serag HB. Hepatocellular carcinoma: an epidemiologic view. *J Clin Gastroenterol* **35**: S72-S78, 2002.
- Shiratori Y, Yoshida H, Omata M. Different clinicopathological features of hepatocellular carcinoma: recent trends in Japan. *Gastroenterology* **127**: S17-S26, 2001.
- Farinati F, De Maria N, Marafin C, et al. Hepatocellular carcinoma in alcoholic cirrhosis: Is sex hormone imbalance a pathogenetic factor? *Eur J Gastroenterol Hepatol* **7**: 145-150, 1995.
- Farinati F, Floreani A, De Maria N, et al. Hepatocellular carcinoma in primary biliary cirrhosis. *J Hepatol* **21**: 315-316, 1994.
- Kosaka A, Takahashi H, Yajima Y, et al. Hepatocellular carcinoma associated with anabolic steroid therapy: report of a case and review of the Japanese literature. *J Gastroenterol* **31**: 450-454, 1996.
- Tavani A, Negri E, Parazzini F, et al. Female hormone utilization and risk of hepatocellular carcinoma. *Br J Cancer* **67**: 635-637, 1993.
- Bosetti C, Turati F, La Vecchia C. Hepatocellular carcinoma epidemiology. *Best Pract Res Clin Gastroenterol* **28**: 753-770, 2014.
- Liver Cancer Study Group of Japan. The 19th national primary liver cancer follow-up report (2006-2007).
- Churiki M, Ikai I, Yamamoto M, et al. A case of Juvenile Hepatocellular Carcinoma without Hepatitis Infection. *Nihon Shokakibyo Gakkai Zasshi (Journal of Japanese Society of Gastroenterology)* **30**: 2292-2296, 1997.
- Ali A, Abdel-Hafiz H, Suhail M, et al. Hepatitis B virus, HBx mutants and their role in hepatocellular carcinoma. *World J Gastroenterol* **20**: 10238-10248, 2014.
- Minor MM, Slagle BL. Hepatitis B virus HBx protein interactions with the ubiquitin proteasome system. *Viruses* **6**: 4683-4702, 2014.
- Lafaro KJ, Pawlik TM. Fibrolamellar hepatocellular carcinoma: current clinical perspectives. *J Hepatocell Carcinoma* **2**: 151-157, 2015.
- Ganesan D, Szklaruk J, Kundra V, Kaseb A, Rashid A, Elsayes KM. Imaging features of fibrolamellar hepatocellular carcinoma. *AJR Am J Roentgenol* **202**: 544-552, 2014.
- Sergi CM. Hepatocellular carcinoma, fibrolamellar variant: diagnostic pathologic criteria and molecular pathology update. *A primer Diagnostics (Basel)* **6**: 3, 2016.
- Ng IO, Ng MM, Lai EC, Fan ST. Pathologic features and patients survival in hepatocellular carcinoma in relation to age. *J Surg Oncol* **61**: 134-137, 1996.
- Hanazaki K, Kajikawa S, Shimozaawa N, et al. Survival and recurrence after hepatic resection of 386 consecutive patients with hepatocellular carcinoma. *J Am Coll Surg* **191**: 381-388, 2000.
- Fujita N, Yoshikawa T, Uemura M, et al. A case of juvenile hepatocellular carcinoma arising from normal liver. *Nihon Shokakibyo Gakkai Zasshi* **92**: 1092-1097, 1995 (Abstract in English).
- Sato S, Tanioka H, Nagata H, et al. A case of non-B, non-C, juvenile hepatocellular carcinoma with brain metastasis. *Nihon Shokakibyo Gakkai Zasshi* **93**: 758-762, 1996 (Abstract in English).
- Nakano M, Ohtsuka Y, Sugaya H, Terano A, Sasai T, Monma T. A case of rapidly progressive hepatocellular carcinoma in a juvenile patient with no serum hepatitis virus marker. *Dokkyo J Med Sci* **29**: 177-181, 2002 (Abstract in English).
- Fukui K, Taniguchi H, Ikoma H, et al. A case with juvenile hepa-

- tocellular carcinoma without hepatitis underwent right hepatic trisegmentectomy. *Jpn J Cancer Chemother* **38**: 2475-2477, 2011 (Abstract in English).
26. Kaibori M, Ishizaki M, Uchida I, et al. A case of non-B, non-C, juvenile hepatocellular carcinoma. *Nihon Shokakibyō Gakkai Zasshi* **40**: 617-622, 2007 (Abstract in English).
 27. Okamoto N, Yamafuji K, Kubochi K, et al. A case of ruptured hepatocellular carcinoma having neither hepatitis B nor hepatitis C virus infection in a young adult patient. *Nihon Shokakibyō Gakkai Zasshi* **43**: 929-934, 2010 (Abstract in English).
 28. Kitagawa H, Tsuda S, Tuboi K, et al. A case of well-differentiated hepatocellular carcinoma that developed in a normal liver on a young woman. *Nihon Rinsho Gega Igakkai Zasshi* (Journal of Japanese society for Clinical Surgery) **75**: 190-196, 2014 (Abstract in English).
 29. Assed C, Marchiori E, Zanetti G, et al. Pulmonary metastases from primary hepatocellular carcinoma in a 26-year-old patient: a case report. *Cases J* **2**: 6256, 2009.
 30. Tanaka T, Hoshino Y, Hayashi S, et al. A case of hepatocellular carcinoma in normal liver with no evidence of HBV or HCV infection. *Hepatogastroenterology* **43**: 1390-1394, 1996.
 31. Herath HM, Kulatunga A. Large hepatocellular carcinoma in a non-cirrhotic liver with peritoneal and omental metastasis in a healthy man: a case report. *J Med Case Rep* **11**: 34, 2017.
 32. Sanada H, Sueda T, Nakai S, et al. Hepatocellular carcinoma in a young woman which occurred in the liver without virus infection. *Nihon Rinsho Gega Igakkai Zasshi* **65**: 1918-1923, 2004.
 33. Kubo Y, Itubo M, Koike K, et al. A young male patient with nonB-nonC hepatocellular carcinoma and concomitant focal nodular hyperplasia developed in the normal liver. *Kanzo* **48**: 233-239, 2007 (Abstract in English).
 34. Sharma ZD, Gokhale VS, Chaudhari N, Kakrani AL. An acute unusual presentation of hepatocellular carcinoma. *J Cancer Res Ther* **10**: 401-403, 2014.
 35. Sunagawa H, Ogura K, Orokawa T, et al. A case of juvenile hepatocellular carcinoma without viral hepatitis. *Liver Cancer* **19**: 23-27, 2013 (Abstract in English).
 36. Abe Y, Nojiri K, Mogaki M, et al. A case of hepatocellular carcinoma (with the maximum diameter of 12 cm) arisen in the normal liver of a 30-year-old man. *Nihon Rinsho Gega Igakkai Zasshi* **80**: 2067-2072, 2019 (Abstract in English).
 37. Hashimoto A, Hattori A, Tanaka T, et al. A case of young hepatocellular carcinoma with non-B, non-C, non-cirrhotic background. *Nihon Shokakibyō Gakkai Zasshi* **118**: 348-357, 2021.

The Internal Medicine is an Open Access journal distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).