Life Style Related Diseases —From Pathogenesis to Prevention—

Moderator: Masato KASUGA

The Second Department of Internal Medicine, Kobe University School of Medicine, Hyogo

Moderator: Yuji MATSUZAWA

Professor and Chairman, Department of Internal Medicine and Molecular Science, Graduate School of Medicine, Osaka University, Osaka

1. Diabetes Mellitus and Lifestyle

Masashi KOBAYASHI

The First Department of Medicine, Toyama Medical and Pharmaceutical University, Toyama

Key words: Obesity, insulin resistance, free fatty acids, TNF- α

Type 2 diabetes mellitus is caused by genetic and environmental background. The latter is mostly influenced by lifestyle

because intervention of lifestyle could prevent diabetes and Japanese Americans show an increased occurrence of type 2 diabetes compared with Japanese who live in Japan. Among the various factors of lifestyle, diet and exercise are the most important factors causing obesity which leads to the occurrence (1, 2). At present, the incidence of diabetes is 60 fold the incidence at the time right after World War II; this trend is closely

Figure 1. Insulin resistance by FFA.

Figure 2. Effect of TNF- α on insulin signal transduction.

*Presented at the 98th Annual Scientific Meeting of the Japanese Society of Internal Medicine, April 14, 2001, Tokyo
Reprint requests should be addressed to Dr. Masashi Kobayashi, the First Department of Medicine, Toyama Medical and Pharmaceutical University, 2630 Sugitani, Toyama 930-0194

related to fat intake the increase in the number of automobile registrations in Japan. These phenomena all lead to obesity which is closely related to insulin resistance.

Obesity is associated with increased adipocyte size and these cells can secrete free fatty acids and TNF- α which cause insulin resistance. Free fatty acids (FFA), as shown in Fig. 1, decrease insulin action by increasing PKC activity which leads to increased phosphorylation of serine/threonine of IRS-1. FFA also suppresses the function of the machinery of the glucose transporter. TNF- α decreases tyrosine phosphorylation of insulin receptors and IRS-1 and also decreases expression of Glut 4 protein (Fig. 2) (3). Thereby, these agents can decrease insulin action which leads to insulin resistance. Another polypeptide called "resistin" which is secreted from enlarged adipocytes, was recently found to suppress insulin action, but the details of the mechanism of resistin action remain to be clarified.

The advancement of molecular biology in the field of insulin signal transduction is remarkable and the knowledge acquired through the recent research can be applied to the development of new antidiabetic drugs. There are several serine/threonine kinases and tyrosine phosphatases which can decrease insulin action in the diabetes state, and adipocytokines, which are produced from enlarged adipocytes, may affect insulin signal transduction (4, 5). To prevent these proteins and cytokines from suppressing insulin action, specific molecular targets can be identified and agents can be developed for the normal insulin signaling; the development of new antidiabetic drugs is on-

going at the present time.

The insulin secretory capacity of Japanese is lower than that of the Western population. Therefore, the ability of compensatory insulin secretion to cope with insulin resistance in Japanese is much less than that in the Western population. Thus, Japanese people are vulnerable to insulin resistance due to their genetic background for insulin secretory ability. These phenomena appear to be prevailing among the Asian population in which the incidence of diabetes is currently increasing due to the westernized style of life.

References

- 1) Marshall JA, Hoag S, Shetterly S, Hamman RF. Dietary fat predicts conversion from impaired glucose tolerance to NIDDM. The San Luis Valley Diabetes Study. *Diabetes Care* **17**: 50-56, 1994.
- 2) Eriksson KF, Lindgarde F. Poor physical fitness, and impaired early insulin response but late hyperinsulinemia, as predictors of NIDDM in middle-aged Swedish men. *Diabetologia* **39**: 573-579, 1996.
- 3) Iwata M, Haruta T, Usui I, et al. Pioglitazone ameliorates tumor necrosis factor- α -induced insulin resistance by a mechanism independent of adipogenic activity of peroxisome proliferator-activated receptor- γ . *Diabetes* **50**: 1083-1092, 2001.
- 4) Takano A, Usui I, Haruta T, et al. Mammalian target of rapamycin pathway regulates insulin signaling via subcellular redistribution of insulin receptor substrate 1 and integrates nutritional signals and metabolic signals of insulin. *Mol Cell Biol* **21**: 5050-5062, 2001.
- 5) Wada T, Sasaoka T, Funaki M, et al. Overexpression of SH2-containing inositol phosphatase 2 results in negative regulation of insulin-induced metabolic actions in 3T3-L1 adipocytes via its 5'-phosphatase catalytic activity. *Mol Cell Biol* **21**: 1633-1646, 2001.

2. Insulin Resistance and Hypertension in Adolescents and Adults

Ikuo SAITO

Health Center, Keio University, Tokyo

Key words: obesity, leptin, sympathetic activity

Insulin resistance and the concomitant hyperinsulinemia have been implicated in the pathogenesis of hypertension and dyslipidemia.

In this review, I will discuss the relation of insulin resistance, hypertension and obesity and examine the role of the sympathoadrenal system in the development of hypertension in the general Japanese population.

Obesity and hypertension

Obesity is associated with an increased risk of hypertension. In our cross-sectional study, 3.5% of male students and 2.5% of female students were obese. Up to 30% of the obese male students and 8% of the obese female students had concomitant hypertension.

In contrast, 5.3% of non-obese male students and 0.6% of non-obese female students had hypertension. Our longitudinal studies of junior high school boys, university students, and middle-aged men showed that a higher baseline body weight and weight gain raised blood pressure, while weight loss reduced blood pressure (1). The mechanisms leading to hyper-