Original Article

Correlation Between Mercury Concentrations in Hair and Dental Amalgam Fillings

Koufuchi Ryo ¹⁾, Atsuko Ito ¹⁾, Rie Takatori ¹⁾, Yoshinori Tai ²⁾, Junji Tokunaga ¹⁾, Kazumune Arikawa ²⁾, Takashi Yamada ¹⁾, Keiko Shinpo ¹⁾, Hiroshi Yasuda ³⁾, Ichiro Saito ¹⁾

1) Department of Pathology, Tsurumi University School of Dental Medicine

2) Department of Geriatric Dentistry, Tsurumi University School of Dental Medicine

3) Research Laboratory, La Belle Vie Inc.

Abstract

To examine possible association between dental amalgam fillings and mercury burden, we measured the mercury concentrations in scalp hair samples from 74 female Japanese subjects aged 26-64 years who had a dietary habit of little fish-consumption. The hair mercury concentrations in the subjects with multiple amalgam fillings were significantly higher than those in the subjects with no amalgam fillings, and correlated with the number of amalgam fillings in the following regression equation: $\log \text{Hg} = 0.084 \text{ amalgam fillings} + 3.377$ ($r = 0.339$, $p = 0.003$). In addition, multiple regression analysis demonstrated that the hair mercury concentrations were significantly correlated with the amalgam fillings ($r = 0.292$; $p = 0.014$) and age ($r = 0.284$; $p = 0.017$) but not significantly with the body mass index ($r = 0.060$). A regression equation of $\log \text{Hg} = 0.071 \text{ amalgam fillings} + 0.011 \text{ age} + 0.006 \text{ BMI} + 2.72$ was obtained ($r = 0.447$), with 0.200 of the determination coefficient. These findings suggest that the dental amalgam fillings positively influence mercury accumulation in the hair and also, probably, in the body.

KEY WORDS: amalgam, hair, mercury, aging, body mass index (BMI)

Introduction

Anti-Aging Medicine is defined as the medicine which makes maintaining health and youth possible while growing older. The aim of this medicine is not simply to extend the life span, but to prevent age-associated physical and mental decline and maintain a high quality of life thus enabling continued social productivity.

The dentistry field also has a deep involvement in toxic metal contamination. It is becoming evident the influence on aging which dental materials used in dental treatment. The medical field is also hoping that dentists will make active efforts to diagnose and improve these conditions. It is clear that the Anti-Aging Medicine region in which dentistry is involved is very diverse. Close cooperation between the dentistry and medical fields will make it possible to implement the above methods thus resulting in a high level of Anti-Aging Medicine.

Mercury is well known as a toxic metal in the human body and at high levels, it causes liver and kidney damage as well as neurological symptoms/disorders ¹⁾. This toxic metal not only inhibits various enzymatic reactions and metabolic processes, but also enhances lipid peroxidation, progression of atherosclerosis and the risk of myocardial infarction and/or stroke, leading to death ²⁻⁵⁾.

Recently, an increasing interest in the health hazards associated with mercury exposure has been raised, and the mercury release from dental amalgam fillings has been blamed for a variety of health complaints ⁶⁻¹⁰⁾. In particular, prenatal exposure to mercury may affect the blood pressure and ability to respond to sensory stimuli in the exposed fetus and in infants later in life ¹⁰⁻¹⁴⁾. Thus,

dental amalgam fillings and restorations have become increasingly recognized as a potentially significant source of inadvertent chronic exposure to mercury, particularly in populations with no dietary dependency on foods of marine origins. Over the past decade, in the Western countries with little mercury intake from fish consumption, mercury release from dental amalgams has been blamed for a variety of health complaints ⁶⁻¹⁰⁾.

The purpose of this study was to examine the possible relation between dental amalgam fillings and mercury concentration in hair, a suitable biomarker of mercury burden ^{3,15)}. In this study, scalp hair mercury concentrations were examined in 74 female Japanese adults aged 26-65 years who had a dietary habit of little fish-consumption. The relation between the number of dental amalgam fillings and hair mercury concentration was examined using regression analysis, and the contribution of amalgam fillings to mercury burden was discussed.

Materials and Methods

Subjects and Materials

Scalp hair samples, well known as a suitable specimen for monitoring human exposure to mercury, were used in this study. On the basis of informed consent, scalp hair samples (the first 2-3cm proximal to the scalp, total weight of about 200mg) from 74 female subjects aged 26-65 years old who had 0-4 amalgam fillings and a dietary habit of little fish-consumption were collected at the outpatient clinic of Tsurumi University School of Dental Medicine and collaborative dental clinics (*Table 1*). Dentists counted the number of amalgam fillings for each patient. The data of body weight and height of each subject were obtained using a questionnaire and the corresponding body mass index (BMI) was calculated. The study protocol was approved by the ethical committee of Tsurumi University.

Table 1 Subject characteristics

Amalgam Filling Teeth	Number of Subjects	Age(year)		BMI	
		Mean	Range	Mean	Range
0	10	48.4	39-62	23.0	18.9-27.1
1	13	45.1	25-59	19.9	15.8-25.6
2	26	51.7	39-61	22.2	17.9-26.9
3	12	52.8	45-64	21.3	18.2-24.6
4	13	52.1	40-63	21.5	17.7-34.8

Analysis of hair mercury

The hair samples (75 mg each) were weighed, placed in 50-mL plastic tubes, and washed with acetone and with 0.01% Triton solution, in accordance with the procedures recommended by the Hair Analysis Standardization Board¹⁶⁾. Each washed hair sample was then mixed with 10 mL of 6.25% tetramethylammonium hydroxide (TMAH, Tama Chemical, Kawasaki-city, Kanagawa) and 50 μ L of 0.1% gold solution (SPEX Certi Prep, Metuchen, NJ, USA), then dissolved at 75°C by shaking for 2 hours. After cooling the solution to room temperature and adjusting its gravimetric volume, the obtained solution was subjected to mercury analysis. The mercury concentrations were measured using inductively coupled plasma mass spectrometry (ICP-MS, Agilent-7500c, California, USA) by the internal standard method, as previously reported¹⁷⁻¹⁹⁾, and were expressed as ng/g hair (ppb).

Statistical Analysis

The hair mercury concentrations were distributed in a nearly log-normal manner, and so converted to their logarithmic values for statistical analysis. The geometric mean (geomean) rather than the arithmetic mean was used to represent the hair mercury concentrations. Statistical significance was determined by one-way ANOVA, and the difference between individual groups was estimated using Scheffe test and Fisher's LSD test. The associations between hair mercury concentration and the number of amalgam fillings, subject age and BMI were determined using a multiple regression analysis.

Results

A scattered plot showing the association between the number of amalgam fillings and hair mercury concentrations is shown in *Figure 1*. The hair mercury concentrations in the subjects with 2 or more amalgam fillings were significantly higher than those in the subjects with no amalgam fillings (*Table 2*). In addition, the hair mercury concentration was significantly correlated with the number of amalgam fillings (AF) in the following regression equation of $\log \text{Hg} = 0.084 \text{ AF} + 3.377$ ($r = 0.339$, $p = 0.003$). Using this regression equation, we estimated one increment of amalgam fillings associated with a 1.2-fold increase in the hair mercury level.

Fig. 1. Relation between the number of dental amalgam fillings and hair mercury concentrations in 74 female Japanese subjects. The relation between the number of amalgam fillings and hair mercury level was examined using simple regression analysis ($r = 0.339$, $p = 0.003$).
 **: Significantly different ($p < 0.01$) from no amalgam filling group;
 *: $p < 0.05$

Table 2 Influence of amalgam fillings on hair mercury concentration

Amalgam Filling Teeth	Hair Mercury Concentration (ppb)		Scheffe Test p value	Fisher's Test p value
	Geomean	Log Hg $\pm$ S.D.		
0	1,687	3.2270 $\pm$ 0.2163		
1	3,095	3.4906 $\pm$ 0.3041	0.311	0.0306*
2	4,278	3.6312 $\pm$ 0.3134	0.009**	0.0003***
3	4,529	3.6560 $\pm$ 0.2463	0.021*	0.0007***
4	3,983	3.6003 $\pm$ 0.2168	0.055	0.0023**

*: $p < 0.05$
 **: $p < 0.01$
 ***: $p < 0.001$

The results of a multiple regression analysis showed that the hair mercury concentration was significantly correlated with the number of amalgam fillings ($r = 0.292$, $p = 0.014$) and patient age ($r = 0.284$, $p = 0.017$) but was not significantly correlated with their BMI ($r = 0.060$, $p = 0.620$) (*Figure 2* and *Table 3*). Thus, the regression equation of $\log \text{Hg} = 0.071 \text{ AF} + 0.011 \text{ Age} + 0.006 \text{ BMI} + 2.72$ was obtained ($r = 0.447$, $p < 0.001$), with the determination coefficient of 0.200.

Fig. 2. Relation between aging and hair mercury concentrations in 74 female Japanese subjects.
The relation between the age and hair mercury level was examined using a simple regression analysis ($r = 0.353$, $p = 0.002$).

Table 3 Multiple regression analysis: Relation of hair mercury concentration with amalgam fillings, age and BMI in female subjects

	Partial R	Simple R	T value	p value
Amalgam	0.292	0.339	2.516	0.014*
Age	0.284	0.353	2.443	0.017*
BMI	0.060	0.111	0.496	0.621

Determination Coefficient = 0.200

Partial R: Partial regression coefficient with multiple regression analysis

Simple R: Simple regression coefficient with simple regression analysis

*: There is a significant ($p < 0.05$) relationship with hair mercury concentration.

Discussion

Hair mineral analyses have been used in forensic medicine, in screening populations for toxic metal poisoning and in monitoring environmental pollutants²⁰. Mercury, especially organic methyl mercury, is well known to be a toxic metal and accumulates in human kidneys, liver and fat-rich tissues such as brain and adipose tissue. In addition, it accumulates with age-dependency and gender-difference (1.5-fold higher in male than in female) in Japanese population²¹⁻²³.

In the present study, a significant, positive relationship between the number of dental amalgam fillings and hair mercury levels in Japanese women was demonstrated (Figure 1 and Table 2, 3). These results are compatible with the findings in pregnant Italian women reported by Luglie *et al.*⁹. They have demonstrated that the number of amalgam fillings positively influenced the mercury concentration in the amniotic fluid of pregnant Italian women⁹. In an experimental study, Takahashi *et al.*²⁴ reported that mercury vapor released from amalgam fillings in pregnant rats was distributed to maternal and fetal organs in significant amounts depending on the number of amalgam fillings.

Some epidemiological studies have shown that prenatal exposure to mercury may affect the blood pressure and responsibility to sensory stimuli in the infants in later life¹⁰⁻¹⁴. Therefore, pregnant women should avoid mercury exposure.

Recently, Yasuda *et al.*¹⁹ reported that in Japanese men the hair mercury concentration is related to the BMI, which is well known to be associated with overall mortality. Higher BMI is reported to be a high risk factor for various diseases such as metabolic syndromes and cancers²⁵⁻²⁷, and also to be one of the strongest predictors of death from cardiovascular disease in men (relative risk, 2.90)²⁸. Thus, mercury exposure seems to be correlated with not only obesity, but also with aging¹⁹. These findings suggest the possibility that dental amalgam fillings may play some role in the pathogenesis of cardiovascular diseases, cancer and physiological senescence.

The present study demonstrated a significant relation between dental amalgam fillings and hair mercury concentrations in Japanese women who had a dietary habit of little fish-consumption. These findings suggest that in addition to dietary mercury intake, dental amalgam fillings also contribute to mercury accumulation in the body.

Acknowledgments

This work was partially supported by grants-in-aid for scientific research from the Ministry of Education, Culture, Sports, Science and Technology of Japan (18592292). The authors sincerely appreciate the members of the Japanese Society of Anti-Aging Dental Medicine for collaborating in this study.

References

- 1) Harada M: Minamata disease: methylmercury poisoning in Japan caused by environmental pollution. *Crit Rev Toxicol* 25; 1-24: 1995
- 2) Guallar E, Sanz-Gallardo MI, van't Veer P, et al: Mercury, fish oils, and the risk of myocardial infarction. *N Engl J Med* 347; 1747-1754: 2002
- 3) Salonen JT, Seppanen K, Lakka TA, et al: Mercury accumulation and accelerated progression of carotid atherosclerosis: a population-based prospective 4-year follow-up study in men in eastern Finland. *Atherosclerosis* 148; 265-273: 2000
- 4) Salonen JT, Seppanen K, Nyyssonen K, et al: Intake of mercury from fish, lipid peroxidation, and the risk of myocardial infarction and coronary, cardiovascular, and any death in eastern Finnish men. *Circulation* 91; 645-655: 1995
- 5) Virtanen JK, Voutilainen S, Rissanen TH, et al: Mercury, fish oils, and risk of acute coronary events and cardiovascular disease, coronary heart disease, and all-cause mortality in men in eastern Finland. *Arterioscler Thromb Vasc Biol* 25; 228-233: 2005
- 6) Berglund A, Molin M: Mercury vapor release from dental amalgam in patients with symptoms allegedly caused by amalgam fillings. *Eur J Oral Sci* 104; 56-63: 1996
- 7) Bratel J, Haraldson T, Meding B, et al: Potential side effects of dental amalgam restorations. (I). An oral and medical investigation. *Eur J Oral Sci* 105; 234-243: 1997
- 8) Ganss C, Gottwald B, Traenckner I, et al: Relation between mercury concentrations in saliva, blood, and urine in subjects with amalgam restorations. *Clin Oral Investig* 4; 206-211: 2000
- 9) Luglie PF, Campus G, Chessa G, et al: Effect of amalgam fillings on the mercury concentration in human amniotic fluid. *Arch Gynecol Obstet* 271; 138-142: 2005
- 10) Razagui IB, Haswell SJ: Mercury and selenium concentrations in maternal and neonatal scalp hair: relationship to amalgam-based dental treatment received during pregnancy. *Biol Trace Elem Res* 81; 1-19: 2001
- 11) Davidson PW, Myers GJ, Weiss B, et al: Prenatal methyl mercury exposure from fish consumption and child development: a review of evidence and perspectives from the Seychelles Child Development Study. *Neurotoxicology* 27; 1106-1109: 2006
- 12) Grandjean P, Murata K, Budtz-Jorgensen E, et al: Cardiac autonomic activity in methylmercury neurotoxicity: 14-year follow-up of a Faroese birth cohort. *J Pediatr* 144; 169-176: 2004
- 13) Steuerwald U, Weihe P, Jorgensen PJ, et al: Maternal seafood diet, methylmercury exposure, and neonatal neurologic function. *J Pediatr* 136; 599-605: 2000
- 14) Vahter M, Akesson A, Lind B, et al: Longitudinal study of methylmercury and inorganic mercury in blood and urine of pregnant and lactating women, as well as in umbilical cord blood. *Environ Res* 84; 186-194: 2000
- 15) Suzuki T, Hongo T, Yoshinaga J, et al: The hair-organ relationship in mercury concentration in contemporary Japanese. *Arch Environ Health* 48; 221-229: 1993
- 16) Cranton E, Bland J, Chatt A, et al: Standardization and interpretation of human hair for elemental concentrations. *J Holistic Med* 4; 10-20: 1982
- 17) Yasuda H, Yonashiro T, Yoshida K, et al: Mineral imbalance in children with autistic disorders. *Biomed Res Trace Elem* 16; 285-291: 2005
- 18) Yasuda H, Yonashiro T, Yoshida K, et al: Relationship between body mass index and minerals in male Japanese adults. *Biomed Res Trace Elem* 17; 316-321: 2006
- 19) Yasuda H, Yoshida K, Fukuchi K, et al: Association of aging with minerals in male Japanese adults. *Anti-Aging Medicine* 4; 38-42: 2007
- 20) Batzevich VA: Hair trace element analysis in human ecology studies. *Sci Total Environ* 164; 89-98: 1995
- 21) Yasuda H, Yonashiro T, Yoshida K, et al: High toxic metal levels in scalp hairs of infants and children. *Biomed Res Trace Elem* 16; 39-45: 2005
- 22) Yasutake A, Matsumoto M, Yamaguchi M, et al: Current hair mercury levels in Japanese: survey in five districts. *Tohoku J Exp Med* 199; 161-169: 2003
- 23) Yonei Y, Mizuno Y, Kido M, et al: Research on toxic metal levels in scalp hair of Japanese. *Anti-Aging Med Res* 2; 11-20: 2005
- 24) Takahashi Y, Tsuruta S, Arimoto M, et al: Placental transfer of mercury in pregnant rats which received dental amalgam restorations. *Toxicology* 185; 23-33: 2003
- 25) Murphy TK, Calle EE, Rodriguez C, et al: Body mass index and colon cancer mortality in a large prospective study. *Am J Epidemiol* 152; 847-854: 2000
- 26) Patel AV, Rodriguez C, Bernstein L, et al: Obesity, recreational physical activity, and risk of pancreatic cancer in a large U.S. Cohort. *Cancer Epidemiol Biomarkers Prev* 14; 459-466: 2005
- 27) Rodriguez C, Patel AV, Calle EE, et al: Body mass index, height, and prostate cancer mortality in two large cohorts of adult men in the United States. *Cancer Epidemiol Biomarkers Prev* 10; 345-353: 2001
- 28) Calle EE, Thun MJ, Petrelli JM, et al: Body-mass index and mortality in a prospective cohort of U.S. adults. *N Engl J Med* 341; 1097-1105: 1999