Meet the Professor I

Tracheal tumors

Douglas J. Mathisen

Harvard Medical School, Boston, USA

Tracheal tumors are uncommon. From 1962 to 2002 we have seen 360 tracheal tumors. Adenoid cystic and squamous cell carcinomas were equally common (N=135). The remaining 90 tumors were an assortment of low grade or benign neoplasms, many one of a kind.

The most common symptoms were dyspnea, cough, hemoptysis, wheezing and stridor. For squamous carcinomas, hemoptysis was the most common symptom (60%) compared to 29% for adenoid cystic carcinoma. For adenoid cystic carcinoma dyspnea and wheezing were more common (65% and 44%) compared to squamous cell carcinoma (50% and 27%). Because adenoid cystic carcinoma is much slower growing, symptom duration is often much longer (19 months v. 7).

Resectability rates have risen from 68% to 82% and operative mortality rates have fallen 21% to 3% over the 40-year period of our experience. Resectability rates are slightly higher for adenoid cystic carcinomas (75%) compared to squamous cell carcinomas (66%). The most common type of resection was sleeve resection of the trachea (53%) and next most common was carinal resection (31%) with and without lung resection.

Clear resection margins for both squamous and adenoid cystic carcinomas were important for survival. Lymph node positivity was more common with squamous cell carcinoma (27%) than adenoid cystic carcinoma (13%).

Multivariate analysis revealed that complete resection, negative margins, adenoid cystic histology and age were statistically significant variables. Surgical resection is superior to palliative therapy.