113**SURGICAL TREATMENT OF 102 CASES WITH PRIMARY TRACHEAL TUMORS**

General Thoracic Surgical Department,
Shanghai Chest Hospital, 241 Huai Hai Rd(W),
Shanghai, P.R. China.
David H. Chao, Yun-Zhong Zhou and OuLin Huang

A total of 102 patients with primary tracheal tumors were operated at the Shanghai Chest Hospital from 1962 through 1993. Tumor located at cervical trachea in 30 cases at thoracic trachea in 72. The lengths of tumor were 1.0-9.5cm.

Pathological diagnosis of squamous-cell carcinoma 38, adenocarcinoma 4, adenoid cystic carcinoma 38, carcinoid 3, and others 6.

All tumors were under gone surgical resections. 38 cases received tracheal circumferential resection and end to end anastomosis, fenestration and tumor resection in 17, carinal resection and reconstruction in 14, partial tracheal wall resection and reconstruction in 6, insertion of "T" tube in 5 cases, Partial tracheal and carinal resection and carinal replacement of bifurcation prosthesis in 2, replacement of partial trachea with artificial vascular prosthesis in 1, using inversive of main bronchus to substitute trachea in 2.

Operative mortality and complication rate were 13.7% and 23.1%. The 5-years survival rate were 46.4%, 3-years were 48.3%. Complications including bleeding 4, infection 3, anastomotic separation 2, anastomotic stenosis 2, ARDS 2, surgical shock 2. The authors believe that surgical resection is the best method for the primary tracheal tumors. If the length of tumors is less than 5cm, end to end anastomosis are more satisfactroy.

114**RESECTION AND RECONSTRUCTION OF THE TRACHEAL CARINA AMONG 114 PRIMARY LUNG CANCER CASES**

General Thoracic Surgical Department,
Shanghai Chest Hospital, 241 Huai Hai Rd(W),
Shanghai, P.R. China.
Lin Qiang, Yun-Zhong Zhou and OuLin Huang

This paper mainly study the treatment effects on 114 lung cancer cases with the method of resection and reconstruction of the tracheal carina. The data collected from clinic during 1978 to 1993. The age range 27-72 years old, male 82, female 32. The modes of operation were as follows: right upper lobectomy with reconstruction of the carina in 52 cases, right sleeve pneumonectomy in 25 cases and left pneumonectomy with carina resection in 37. The survival rates of the remainder were 97%(111/114), 88%(110/114), 57%(64/114), 39%(44/114), 19%(22/114), at 6 months, 1, 3, 5, and 10 years respectively. The results indicate that tracheobronchoplasty is useful when the carina is invaded by cancer. The operation is benifit to treat lung cancer and therefore expanded the surgical indication. At the same time, the 5-years survival rate is satisfactory.

115**CLOSURE OF BRONCHOPLEURAL FISTULA BY USING INTRAPLEURAL TETRACYCLINE HYDROCHLORID SOLUTION.**

Institute of Diseases of The Chest and Hospital,
Dhaka,Bangladesh.
AKM.Shariful islam, MN.Alam Khan, Shafiquel Ahsan,
Md.Zillur Rahman, Md.Nurul Islam.

To close bronchopleural fistulae without any magor surgical procedure a prospective study on closure of bronchopleural fistulae with intrapleural Tetracycline Hydrochloride solution wash in 240 consecutive patients was carried out at the Institute of Diseases of the Chest and Hospital from January 1990 to June 1994. 91.6% of patients with small bronchopleural fistulae were cured by this method (P=.001). Where as only 33.3% of patients with large fistulae showed improvement only. There were no serious side effects of the procedure. Persistant air leak longer than seven to ten days after thoracostomy tube drainage were considered as a case of bronchopleural fistula and selected for Tetracycline Hydrochloride solution wash. Before introducing thoracostomy tube confirmation was made by radiology. .5% lignocaine 20ml, 2gm. of Tetracycline Hydrochloride solution (20 ml) and distilled water 20 ml were introduced through the thoracostomy tube by a syringe and clamped for 6 to 8 hours. Same intervention continued upto five times in every 8 to 10 days interval when required. It is concluded that intrapleural Tetracycline Hydrochloride solution wash is very easy, low coast and effective technique to treat bronchopleural fistulae irrespective of etiology.

116**VIDEO-ASSISTED THORACOSCOPIC SURGERY: KOREA UNIVERSITY EXPERIENCE**

Korea University Medical Center, Department of Chest Surgery, Seoul, Korea
Kwang Taik Kim, Sung Ho Lee, In Sung Lee,
Hyoung Mook Kim.

We report our experience with VATS for treatment and diagnosis of thoracic disease. From March 1992 to October 1994, a VATS procedure was performed in 211 patients. Indications were spontaneous pneumothorax (155), pleural effusion (9), mediastinal tumor (17), pulmonary nodule (9), diffuse lung disease (10), pulmonary metastases (2), traumatic thoracic injury (3), hyperhydrosis (2), esophageal leiomyoma (2), Boerhaave's syndrome (1), pericardial effusion (1). The procedures were bullectomy and pleurodesis with 50% glucose, apical pleurectomy, thoracic sympathectomy lung biopsy, pleural biopsy, excision of pericardium, wedge resection of pulmonary nodule, excision of mediastinal mass, excision of esophageal mass. The complication rate was 5.8%(9/155)in spontaneous pneumothorax patients and 3%(2/56) in other procedures. The mortality within postoperative 30 days was 1% (2/211), including 1 acute renal failure and respiratory failure. The conversion to thoracotomy during VATS were 7 cases, including 2 immediate postoperative massive air leakage, 1 giant bullae, 2 mediastinal tumor, 1 pulmonary malignancy, 1 esophageal tumor. The recurrence rate of pneumothorax after VATS was 8%(13/155). The success rate of VATS in other thoracic disease except pneumothorax was 85%.

In conclusion, although prospective trials should be progressed to define the role of VATS, the VATS carries a low morbidity and mortality and high success rate and can be effectively applicated to the surgical treatment of the various thoracic problems.