DOMINANCE ORDER IN CRICKETS:
ORTHOPTERA: GRYLLIDAE
AND THE EFFECTS OF SOCIAL INVERSION

PETER O. PERETTI

Kennedy-King College

BOICE, HUGHES and COBB (3) found that both gerbils and hamsters used aggressiveness in gaining dominance at the watering spot. With hamsters, aggression was positively correlated with tunnel dominance, body weight, and watering spot dominance. The position in the dominance hierarchy for male baboons depended upon his fighting ability in conjunction with his physical condition and relationships with other baboons (grooming, etc.). Social control in baboons was based on physical dominance (6).

In a study with crayfish, BOUVBJERG (5) found that size correlated with aggressiveness. The larger the animal the more aggressive the organism. Generally the larger, older animals were dominant. Visual and tactual senses played a role in furnishing the cues for dominance. If, in fact, both senses were lost, the ability to receive and transmit cues to dominance to other crayfish was lost (4).

Other conditions can affect dominance. Handled rats (handled while young) were found to be more dominant in competition with non-handled rats. Socially raised rats were also found to dominate non-socially raised rats (2).

ALEXANDER (1) formulated five levels in order to classify crickets according to their aggression levels. Using ALEXANDER's method, crickets could be observed fighting; rated on their aggressiveness; who won or who lost, and then placed in a dominance hierarchy according

to the way in which they scored on the previous behavioral measures.

BOUVBJERG (4) using crayfish, allowed them to build up a dominance order. Once the order was completed, the dominant crayfish of the group was removed while observation remained on the other three animals. After five days the emergent dominant crayfish was removed and reintroduced with the isolated original dominant. Crayfish were removed from the original group and then placed into the inverted group until the original was terminated. The results showed that the first and the second ranking individuals of the original group (with one exception) reestablished their original position through the inversion. BOUVBJERG concluded that the establishment of a dominance order implied that there were intrinsic aggressive differences between individuals. He stated the important implication of his experiment was that sensory perception of the individual crayfish and the retention of the recognition, as a form of learning, would follow in the social inversion method.

METHOD

Subjects: Four adult, male crickets of the same species (*Orthoptera: Gryllidae*). All of the animals were of the same size and housed in a large terrarium with separate quarters. Each quarter was equipped with its own food and water from which the crickets could partake on an *ad lib* basis. Individual crickets were marked for recognition by painting a number (I, II, III, or IV) on the thorax of each animal with a light-colored nail polish.

Materials: The terrariums used in the research had outside measurements of 20 in. length x 10½ in. in width x 12 in. in height. The floors of the units were covered with earth, sand, grass, and twigs to depth of 3 in. to simulate a "natural" environment. Loosely packed, the base of the terrariums allowed the crickets to dig and form shelters, if they desired.

Procedure: Dominance in the order of the crickets was determined by the Alpha, or first position; Beta, the second position; Gamma (third position), and the fourth position which was Delta. Each of the four Ss had not ever been in social contact with the others. The crickets were purchased individually from

the same supply house and allowed one week to adjust to the laboratory atmosphere.

After the week in separate quarters, two of the crickets were placed in a neutral terrarium, the same size as the "home unit", and the behavior that followed was observed. This procedure was followed until all four crickets had encountered each other five times. Encounters were coded similar to the way BOUVBJERG's (4) explanation of aggression in crayfish. There were four types of tension contact: avoidance, threat, strike and fight (C s D means that animal C struck animal D).

A won-loss record was obtained for each animal. A loss was determined by the retreat of one of the animals or by throwing the other back a few inches, flipping the other into the air, or by turning the other on his side. A dominance hierarchy was thus established.

The method of social inversion was then used on all of the Ss. The method involved the removal and isolation of the dominant (Alpha) cricket. Observations on the other three crickets remained for twenty-four hours. After this time the emergent dominant cricket of the original group was removed and reintroduced with the the isolated original dominant. Observations on both groups continued and again, after twenty-four hours, the emergent dominant cricket of the original group was removed and introduced into the inverted group. After another twenty-four hours the fourth cricket was removed and introduced into the inverted group. The original group was now completely terminated.

When one cricket from the original group was placed into the inverted group, five encounters with each other cricket took place. During all the time of the total experiment the Ss lived in their individual quarters. Fighting was done in the neutral terrarium. A won-loss record for each individual S was tabulated for fights in the inverted group as well as in the original group.

RESULTS

A record of each S's encounters was made. Table 1 shows the record of the encounters; the winner being specified in each column. Each S fought all other Ss five times. A record of the total win-loss for each S is shown in Table 2. Table 2 also indicates a percentage of total wins. Table 3 indicates the performance of each cricket with the other crickets (win-loss). Tables 1, 2, and 3 are *before* social

Table 1: Results of encounters, each cricket with all others. The winner *S* specified in the column for each encounter.

Subjects		Encounters				
		1st	2nd	3rd	4th	5th
I	vs II	I	II	I	I	II
I	vs III	III	III	I	III	I
I	vs IV	I	IV	I	I	I
II	vs III	II	II	III	II	II
II	vs IV	IV	II	II	II	IV
III	vs IV	III	IV	III	III	III

Table 2: The total win-loss record for each cricket for all encounters. Percentage of total wins is also recorded. 15 fights for each cricket pere performed.

Subject	Win	Loss	% Wins
I	9	6	60.0
II	9	6	60.0
III	7	8	46.6
IV	5	10	33.3

Table 3: A record of the wins and losses of each cricket to all other crickets. Five encounters with each cricket. I vs II equals 3 (wins) -2 (losses).

Subject	I	II	III	IV
I	—	3-2	2-3	4-1
II	2-3	—	4-1	3-2
III	3-2	1-4	—	3-2
IV	1-4	2-3	2-3	—

inversion. Tables 4, 5, and 6 are comparable to the first three tables, respectively, as they indicate similar data. The former tables differ from the latter in that Tables 4, 5, and 6 denote performances *after* social inversion. Of note was the fact that *S* performance *before* and *after* social inversion was similar.

Figure 1 shows the percentage of total wins for each cricket for both *before* and *after* the social inversion process had taken place. Arithmetic differences between the two percentages is also shown.

Figure 2 shows the dominance hierarchy *before* and *after* social inversion.

CONCLUSIONS AND

DISCUSSION

Crickets I and II were the dominant *S*s in the original group. The dominance order in the original group was not completely a straight-line dominance order. *S* I was dominated in the encounters with *S* III three out of five

times but it dominated Ss II and IV. S II, whose win-loss record and percentage of wins was similar to S I's, showed dominance over Ss III and IV. Thus S I dominated Ss II and IV while S II dominated Ss III and IV. S I dominated S II.

Cricket I was affected by the social inversion process by increasing its number of total wins (by 6.6%). Conversely, Ss II and III decreased in the number of their total wins (20% and 13%, respectively). Similarly to Cricket I, S IV increased in the number of total wins. Cricket IV went from 33% total wins to 60% total wins. This was an increase of 26%.

After the social inversion process it can be seen that S I remained in

the Alpha position (this time alone) with a 66 percentage of wins. The main difference in the hierarchy which differed from the original group's was due to S IV's increased percentage of wins (by 26%). Under the original group's condition, animal IV was dominated by all

Table 4 : Results of encounters, each cricket with all others. The winner being specified in the column for each encounter.

Subjects			Encounters				
			1st	2nd	3rd	4th	5th
I	vs	II	I	II	I	II	I
I	vs	III	I	I	I	III	III
I	vs	IV	I	IV	I	I	I
II	vs	III	II	II	III	III	II
II	vs	IV	IV	II	IV	IV	IV
III	vs	IV	IV	IV	III	IV	IV

Table 5 : The total win-loss record for each cricket for all encounters. The percentage of total wins is also recorded. 15 fights for each cricket were performed.

Subject	Win	Loss	% Wins
I	10	5	66.6
II	6	9	40.0
III	5	10	33.3
IV	9	6	60.0

Table 6 : A record of the wins and losses of each cricket to all other crickets. Five encounters with each cricket.

Subject	I	II	III	IV
I	—	3-2	3-2	4-1
II	2-3	—	3-2	1-4
III	2-3	2-3	—	1-4
IV	1-4	4-1	4-1	—

and had won 33% of his total encounters. After the social inversion process, animal IV moved into the Beta position (II and III moved into third and fourth positions, respectively) and maintained 60%

Fig. 1. Percentages of total wins per cricket—Before shaded; After blank social inversion. Arithmetic difference is shown below the Figure for dominance performance before and after.

wins of all encounters. S IV completely dominated (four wins to one loss) Ss II and III and was completely dominated, in turn, by S I (four losses to one win). S I was dominant over animals II and III (with three win and two losses to each animal). After the social inversion process the dominance hierarchy was of the straight-line type with S I as Alpha, S IV as Beta, S II as Gamma, and S III as Delta (see Fig. 2).

The *before* and *after* social inversion data do suggest that the dominance order in the original group and the dominance

order in the inverted group were similar. The Alpha S (I) retained

Fig. 2. The order of dominance *Before* and *After* the social inversion process had taken place. Point of arrow shows the direction of dominance (i.e. A→B equals A dominates B). Roman numerals indicate the animal.

its position in each group while the other Ss generally retained their individual statuses. Why did S IV improve in the inverted group? There could be these possibilities: First, cricket IV was the last to be inverted. It was in the place of dominance in the original group for a while. While S IV was alone in the original group, Ss I, II and III were taking part in their encounters. Ss II and III were thus dominated by animal I. Secondly, S IV was paired against S I primarily in the encounters—once S IV was brought into the inverted group. S IV lost the first four of its encounters and won the final one (with S I). Being in isolation (alone in the original group) and having just won an intense battle (last encounter with S I), S IV had the definite advantage in the upcoming encounters. These two aspects could also explain why S IV did so well in the inverted group as compared to its performance in the original group. Thirdly, while not explaining why S IV's performance increased in the inverted group, it was found that animal IV was in the process of molting (shedding of shell) all the time while the encounters of the original group were taking place. Thus, S IV would retreat immediately, because he had no protective covering, rather than risking serious injury. Its five wins were recorded when: (a) it was cornered and had to fight (the condition of cornering gives the cornered animal the advantage, (1)); (b) it was toward the end of the molting period. Perhaps if animal IV was not molting during the primary (original) stage his win-loss record would have been more like his win-loss record of the second (inverted) stage.

SUMMARY

An experiment which analyzed the dominance order of four male crickets and the influence of social inversion. The Ss were paired with each other in encounters in order to obtain a dominance order. Once the order was determined the Ss were taken out from the group, starting with the Alpha and going down to the Delta animal, until all animals were inverted into the new group. The inverted group remained

dominated by the same animal as in the original group. One cricket (IV) showed an increase of 26% in the amount of total wins. The increase was explained as a result of : 1) IV was the last animal to be inverted ; 2) going into battle with II and III, IV was coming off from isolation and a previous win-an advantage in his favor, and 3) IV was molting during the time of the encounter. Results indicated that the dominance order in the original group and the order in the inverted group were similar although certain differences were noted.

REFERENCES

1. ALEXANDER, R. D. Aggressiveness, territoriality and sexual behavior in field crickets (*Orthoptera : Gryllidae*). *Behavior*, 1961, 17, 130-223.
2. BECKER, G. and EZINGA, G. Early banding and social rearing effects on dominant-subordination behavior in the adult rat. *Psychon. Sci.*, 1969, 15, 27-28.
3. BOICE, R., HUGHES, D. and COBB, C. Social Dominance in gerbils and hamsters. *Psychon. Sci.*, 1969, 16, 127-128.
4. BOUVBJERG, J. Dominance order in crayfish : *Orconectus virilis* (Hagen). *Physiol. Zool.*, 1953, 26, 163-78.
5. BOUVBJERG, J. Some factors affecting aggressive behavior in crayfish. *Physiol. Zool.*, 1956, 29, 127-136.
6. WASHBURN, S. L. and DEVORE, I. Social life of baboons. In *Psychobiology : The biological bases of behavior*. San Francisco: W. H. Freeman and Company, 1966, 10-19.

(1974年2月1日 原稿受付)