Localization of Quantitative Trait Loci for Yield Components in a Cross Oriental × Occidental Barley Cultivar (*Hordeum vulgare* L.)

Mohammad SAMERI and Takao KOMATSUDA*

Plant Genome Research Unit, National Institute of Agrobiological Sciences (NIAS)
(Tsukuba, Ibaraki 305–8602, Japan)

Abstract

Ninety-nine recombinant inbred lines (RILs) were grown for three years and yield components were evaluated. Composite interval mapping showed that the *vrs1* and *uzu* genes consistently controlled grain weight per plant. The analyses also identified a new QTL of kernel length on chromosome 3H, a QTL for kernel width on chromosome 6H and a QTL for thousand-kernel weight on chromosome 2H. The QTL generated in this study may be a useful target for marker-assisted selection in breeding programs.

Disciplines: Biotechnology / Plant breeding

Additional key words: composite interval mapping, dense spike (*dsp1*), six-rowed spike (*vrs1*), *uzu* or semibrachytic (*uzu1*)

Introduction

Oriental and occidental barley cultivars are genetically divergent from each other and they harbor genes characteristic of each region²¹. We developed recombinant inbred lines (RILs) from a cross between ‘Azumamugi’ and ‘Kanto Nakate Gold’. Azumamugi was bred from a cross between two Japanese landraces (‘Tochigi Sekitori 1’ and ‘Torano-o Sai 1’). Kanto Nakate Gold was bred by the following pedigree: Golden Melon/Shikoku//Kinki Shu, where Golden Melon is a two-rowed erectum-type cultivar (origin Northern Europe), Shikoku is a Japanese landrace, and Kinki Shu is a selection from Golden Melon. Because Azumamugi and Kanto Nakate Gold belong to oriental and occidental cultivars respectively, the two genotypes are highly polymorphic for morphological, physiological, and molecular markers^{7,8,19}. The objective of the present study is to identify QTLs for yield components, especially grain size in barley. We estimated genetic effects of known major genes and considered to detect new genes even with minor effects on these traits.

Materials and methods

Ninety-nine RILs (F_{12} and F_{13}) were developed from F_2 plants of Azumamugi × Kanto Nakate Gold by single seed descent. Azumamugi is six-rowed (*vrs1.a*), winter type (*Sgh1/sgh2*) and *uzu* or semi-brachytic (*uzu1*). Kanto Nakate Gold is two-rowed (*Vrs1.b*), spring type (*sgh1/Sgh2*) and non-*uzu* (*Uzu1*). Plants were sown in a field of NIAS, Tsukuba, in October 2000, 2001 and 2002, and grown until maturity in summer of 2001, 2002 and 2003 (thereafter referred to as ‘Year 2001’, ‘Year 2002’ and ‘Year 2003’, respectively). Each RIL was represented by a row of ten seeds and spaced 20 cm apart. Rows were spaced 80 cm apart. Four central plants of each RIL were harvested as a bundle of single plants, air-dried for one week at 30°C, and used for measuring quantitative traits. For seed sizes, 10 seeds per RIL were measured using a projector Movias NAC 5200 (Image Technical Center, Yokohama, Japan).

The heritabilities (h^2) were estimated by the formula: $h^2 = \delta_g^2 / (\delta_g^2 + \delta_e^2/r)$, where $\delta_g^2 = (MS_{\text{genotype}} - MS_{\text{error}})/r$, $\delta_e^2 = MS_{\text{error}}$ and r is number of replications. The correlation coefficients were computed using the computer program SPSS (Version 10.01, SPSS Corp., Chicago, IL).

Molecular linkage map^{11,12} and base map¹³ of these

*Corresponding author: e-mail takao@affrc.go.jp

Received 12 June 2006; accepted 26 October 2006.

RILs were constructed previously. Map distances were estimated by using Kosambi's map function⁹. Composite interval mapping (CIM) was performed using the computer program QTL Cartographer² version 1.14. A LOD score threshold of 3.0 was used to identify QTLs. The position of detected QTLs were compared with the ones of former QTL studies^{1,3,10,15} consulting integrated maps^{6,16}.

Results and discussion

RILs showed a wide range of variation for all the traits (Table 1). ANOVA showed that genetic variances for all the traits were significant, and heritability ranged from 0.73 to 0.99 (Table 2). Correlation coefficients between two years ranged from 0.61 to 0.98, which were

significant for all the traits (Table 3). Grain number per spike (GNS) and grain number per plant (GNP) were positively correlated with grain weight per plant (GWT_g), but negatively correlated with thousand-kernel weight (KWT_g) (Table 3).

GWT was controlled predominantly by *vrs1* and *uzu* as detected previously^{3,5,15,23}. The allele of Azumamugi at the *vrs1* locus increased GWT by 9.15–11.75 g (Table 4). KWT of six-rowed lines was 74% of two-rowed lines, but GNP of six-rowed lines was 2.7 times of two-rowed lines, therefore GWT of six-rowed lines was 1.5 times of two-rowed lines, showing six-rowed lines have a potential of higher yield than two-rowed lines (Fig. 1). The allele of Azumamugi at the *uzu* locus decreased GWT by 9.9–10 g (Table 4). However, the *uzu* locus did not control the GNP. Another QTL for GWT was detected on chromo-

Table 1. Means, coefficient of variations and standard deviations for agronomic traits observed in Azumamugi, Kanto Nakate Gold and their 99 RILs

Traits	GWT (g)	STW (g)	HI (%)	KLN (mm)	KWD (mm)	KS	KA (mm ²)	GNS	GNP	KWT (g)
Year 2002										
Azumamugi	61.3	52.3	53.9	7.5	3.4	2.3	25.1	92.1	1,953	35.2
Kanto Nakate Gold	80.0	106.3	43.0	10.0	3.8	2.7	37.3	39.0	1,278	53.0
RIL – mean	54.9	70.9	43.1	8.6	3.5	2.5	29.9	66.4	1,864	38.5
RIL – SD ¹⁾	22.5	24.6	9.5	0.7	0.3	0.3	3.5	30.2	905	7.3
RIL – minimum	12.5	25.0	18.8	7.1	3.0	1.8	22.9	25.8	399	24.4
RIL – maximum	117.5	145.0	62.5	10.4	4.2	3.5	37.4	108.0	3,933	54.4
RIL – CV ²⁾	41.0	34.6	22.1	8.5	7.6	11.1	11.6	45.5	48.6	18.9
Year 2003 ³⁾										
Azumamugi	50.0	25.0	66.7	7.6	3.3	2.3	25.2	96.0	3,192	34.0
Kanto Nakate Gold	47.5	46.3	50.7	9.9	3.7	2.7	36.6	36.0	1,593	49.6
RIL – mean	43.1	57.4	42.3	8.2	3.6	2.3	29.6	62.1	1,759	33.4
RIL – SD	17.6	17.6	10.8	0.7	0.2	0.2	3.6	27.5	826	6.9
RIL – minimum	6.3	20.0	10.4	6.8	3.0	1.8	22.4	21.0	578	18.0
RIL – maximum	87.5	127.5	61.9	9.9	4.2	3.4	37.2	105.0	3,600	47.6
RIL – CV	40.9	30.6	25.5	8.6	6.7	10.5	12.0	44.3	47.0	20.8

1): Standard deviation.

2): Coefficient of Variation (%).

3): Data in italic were scored in the Year 2001.

Table 2. Analysis of variance for agronomic traits in the RILs from the cross of Azumamugi × Kanto Nakate Gold

Source	df	GWT	STW	HI	KLN	KWD	KS	KA	GNS	GNP	KWT
Genotype	98	693.67**	721.08**	178.19**	0.93**	0.11**	0.12**	21.04**	1,647.30**	1,406,931.00**	95.70**
Year	1	6,913.64**	9,102.58**	35.33ns	7.18**	0.68**	1.53**	5.33ns	932.75**	540,248.70*	1,311.57**
Error	98	124.40	191.11	29.00	0.10	0.02	0.01	3.74	20.98	95,717.71	5.57
Heritability		0.82	0.73	0.84	0.89	0.82	0.90	0.82	0.99	0.93	0.94

*, **: significant at the 5% and 1% level respectively.

ns: not significant.

Table 3. Correlation coefficients between agronomic traits in the RILs (Year 2002)

	GWT	STW	HI	KLN	KWD	KS	KA	GNS	GNP	KWT
GWT	<u>0.717**</u>									
STW	0.530**	<u>0.614**</u>								
HI	0.620**	-0.281**	<u>0.726**</u>							
KLN	0.279**	0.504**	-0.150	<u>0.805**</u>						
KWD	-0.376**	0.093	-0.488**	0.047	<u>0.688**</u>					
KS	0.485**	0.333**	0.221*	0.724**	-0.649**	<u>0.830**</u>				
KA	-0.057	0.418**	-0.433**	0.752**	0.691**	0.091	0.698**			
GNS	0.476**	-0.189	0.700**	-0.256**	-0.688**	0.283**	-0.644**	<u>0.979**</u>		
GNP	0.701**	0.100	0.668**	-0.067	-0.681**	0.429**	-0.508**	0.903**	<u>0.876**</u>	
KWT	-0.208*	0.293**	-0.455**	0.422**	0.732**	-0.189	0.794**	-0.814**	-0.722**	<u>0.891**</u>

*, **: significant at the 5% and 1% level respectively. Correlation coefficients between two years are underlined.

some 3HL, but it may coincide with *denso* (*sdw1*)^{3,6,16}.

The major factor of GNP, GNS and KWT was the *vsr1* locus (Table 4). Six-rowed lines decrease KWT but considerably increase GNP, thus six-rowed spikes increase the yield of grains (Fig. 1). The *vsr1* locus contributed 25 to 80% of phenotypic variance for KWT in separate studies^{1,4,5,14,24} and 63–72% in this study. In spite of the major effect by the *vsr1* locus on KWT, we detected a new QTL for KWT named *qKWT.ak-2H* on the short arm of chromosome 2H (Table 4). The allele of Azumamugi at this QTL increased KWT.

The *vsr1*, *uzu* and *dsp1* have been reported to control the kernel sizes^{1,20} and our result was in agreement with these reports. Behind the pronounced effects of these major genes, we detected a new QTL for kernel length (KLN_{mm}) named *qKLN.ak-3H* on chromosome 3HS and a new QTL for kernel width (KWD_{mm}) named *qKWD.ak-6H* on chromosome 6H (Table 4). The QTL of KWD detected on chromosome 7H can be identical with *qDHE.ak-7H*, which controlled days to heading¹⁸. Kernel shape ($KS = KLN_{mm}/KWD_{mm}$) was controlled by *vsr1*, *uzu* and *dsp1*, but kernel area ($KA_{mm}^2 = KLN_{mm} \times KWD_{mm}$) was controlled by *vsr1* and *uzu*.

QTL for straw weight per plant (STW_g) was detected on chromosome 3HL, at the *uzu* locus, and the allele of Azumamugi decreased STW by 6–11.8 g (Table 4). Another QTL for STW was detected on chromosome 3HL (may coincide with *denso*), together with *Ppd-H1* (*Eam1*) and *Sgh2* (spring growth habit²²), which primarily controlled days to heading^{10,17}. The QTL of harvest index ($HI_g = 100 \times GWT/(GWT + STW)$) detected on chromosome 5HL could correspond to *qTPN.ak-5H*, which was a QTL for triplet number¹⁸.

Our study detected new QTLs for controlling seed size. These QTLs are useful for modifying spike architecture moderately rather than the major genes of strong pleiotrophic effects. The QTLs may be introduced to cul-

Fig. 1. Effects of the six-rowed spike gene (*vsr1*) and *uzu* gene on grain weight per plant (GWT), grain number per plant (GNP) and thousand-kernel weight (KWT) for 'Year 2002'.

Error bars represent standard error. Any two means having a common letter are not significantly different at the 5% level of significance in the Student's *t* test distribution.

Table 4. QTLs of yield components detected in the RILs of Azumamugi × Kanto Nakate Gold

Trait	Chr.	2002					2003					Locus ³⁾
		Flanking markers	Pos. cM	LOD	AE ¹⁾	PVE ²⁾	Flanking markers	Pos. cM	LOD	AE	PVE	
GWT (g)	2HL	MWG801 - <i>vrs1</i>	77.56	9.80	11.75	0.262	<i>vrs1</i> - MWG503	79.56	9.90	9.15	0.243	<i>vrs1</i>
	3HL	<i>uzu</i> - <i>e06m30.10.1</i>	65.53	7.10	-10.00	0.175	<i>uzu</i> - <i>e06m30.10.1</i>	65.53	8.75	-9.90	0.218	<i>uzu</i>
	3HL	MWG973 - <i>e06m30.8.3</i>	111.39	3.88	7.33	0.100	ND ⁴⁾					<i>sdw1</i>
GNP	2HL	<i>vrs1</i> - MWG503	79.56	36.14	789.63	0.738	<i>vrs1</i> - MWG503	81.56	26.98	801.39	0.620	<i>vrs1</i>
	3HL	<i>e06m30.8.3</i> - <i>e12m19.9.2</i>	116.56	3.10	150.54	0.026	ND					<i>sdw1</i>
	5HL	<i>e07m25.3</i> - <i>e12m19.9.1</i>	110.13	4.42	-190.11	0.043	<i>e07m25.3</i> - <i>e12m19.9.1</i>	110.13	3.61	-194.03	0.054	<i>Sgh2</i>
GNS	2HL	<i>vrs1</i> - MWG503	79.56	70.47	29.99	0.857	<i>vrs1</i> - MWG503	79.56	66.86	28.01	0.911	<i>vrs1</i>
	2HS	<i>e13m31.7.1</i> - <i>e15m19.8.1</i>	57.49	4.09	1.59	0.043	ND					<i>qKWT.ak-2H</i>
	2HL	<i>vrs1</i> - MWG503	79.56	32.88	-6.57	0.715	<i>vrs1</i> - MWG503	79.56	25.52	-5.95	0.629	<i>vrs1</i>
KWT (g)	5HL	<i>e07m25.3</i> - <i>e12m19.9.1</i>	112.13	4.42	2.02	0.049	ND					<i>Sgh2</i>
	2HL	ND					<i>vrs1</i> - MWG503	83.56	7.18	-0.25	0.113	<i>vrs1</i>
	3HS	ND					<i>ABG070</i> - <i>e07m25.7.1</i>	0.01	4.87	-0.18	0.062	<i>qKLN.ak-3H</i>
KLN (mm)	3HL	<i>uzu</i> - <i>e06m30.10.1</i>	65.53	17.96	-0.52	0.461	<i>uzu</i> - <i>e06m30.10.1</i>	67.53	22.02	-0.50	0.450	<i>uzu</i>
	7HS	<i>cMWG704</i> - <i>e11m17.10.2</i>	86.02	7.96	-0.30	0.160	<i>cMWG704</i> - <i>e11m17.10.2</i>	84.02	11.40	-0.32	0.200	<i>dsp1</i>
	2HL	<i>vrs1</i> - MWG503	79.56	20.08	-0.20	0.520	<i>vrs1</i> - MWG503	79.56	22.81	-0.19	0.523	<i>vrs1</i>
KWD (mm)	6H	<i>e11m18.12</i> - <i>e15m19.9.2</i>	35.78	4.52	0.08	0.078	ND					<i>qKWD.ak-6H</i>
	7HS	ND					<i>ABG701</i> - <i>cMWG704</i>	76.07	3.66	0.06	0.055	<i>dsp1</i>
	7H	<i>e11m17.10.2</i> - <i>e12m19.10.3</i>	97.54	3.90	0.08	0.089	ND					<i>qDHE.ak-7H</i>
KS	2HL	<i>vrs1</i> - MWG503	79.56	7.56	0.11	0.145	<i>vrs1</i> - MWG503	79.56	3.55	0.06	0.062	<i>vrs1</i>
	3HS	ND					<i>ABG070</i> - <i>e07m25.7.1</i>	0.01	3.27	-0.06	0.056	<i>qKLN.ak-3H</i>
	3HL	<i>uzu</i> - <i>e06m30.10.1</i>	67.53	15.24	-0.19	0.378	<i>uzu</i> - <i>e06m30.10.1</i>	65.53	12.15	-0.01	0.265	<i>uzu</i>
KA (mm ²)	6H	<i>ABG458.1</i> - <i>e11m18.12</i>	34.70	3.86	-0.08	0.073	ND					<i>qKWD.ak-6H</i>
	7HS	<i>cMWG704</i> - <i>e11m17.10.2</i>	86.02	12.08	-0.15	0.282	<i>cMWG704</i> - <i>e11m17.10.2</i>	82.02	10.83	-0.12	0.241	<i>dsp1</i>
	2HL	<i>vrs1</i> - MWG503	79.56	13.23	-2.13	0.350	<i>vrs1</i> - MWG503	81.56	14.61	-2.69	0.551	<i>vrs1</i>
STW (g)	3HL	<i>uzu</i> - <i>e06m30.10.1</i>	67.53	7.13	-1.53	0.173	<i>uzu</i> - <i>e06m30.10.1</i>	67.53	7.54	-2.11	0.329	<i>uzu</i>
	2HS	<i>ABG602</i> - <i>e13m23.6</i>	47.29	5.14	-9.20	0.137	<i>ABG602</i> - <i>e13m23.6</i>	47.29	4.49	-6.70	0.139	<i>PpdH1</i>
	3HL	<i>uzu</i> - <i>e06m30.10.1</i>	65.53	7.66	-11.78	0.207	<i>uzu</i> - <i>e06m30.10.1</i>	65.53	3.74	-6.00	0.107	<i>uzu</i>
HI (%)	3HL	MWG973 - <i>e06m30.8.3</i>	115.39	5.02	9.43	0.137	ND					<i>sdw1</i>
	5HL	<i>e12m26.9</i> - <i>e07m25.3</i>	103.09	4.66	-8.69	0.118	ND					<i>Sgh2</i>
	2HS	<i>e13m23.6</i> - <i>e13m31.7.1</i>	49.82	6.03	3.04	0.090	ND					<i>PpdH1</i>
HI (%)	2HL	MWG801 - <i>vrs1</i>	77.56	18.00	6.43	0.380	<i>vrs1</i> - MWG503	79.56	22.60	9.01	0.635	<i>vrs1</i>
	3HL	ND					<i>ABA001</i> - <i>uzu</i>	63.76	3.11	-2.95	0.064	<i>uzu</i>
	5HL	<i>e14m27.4.4</i> - MWG2230	86.37	3.19	2.14	0.050	ND					<i>qTPN.ak-5H</i>

1): Additive effect of "Azumamugi" allele.

2): Proportion of phenotypic variance explained by the detected QTLs.

3): New QTLs identified in this study are in bold face.

4): Not detected.

tivars by means of marker-assisted selection in breeding programs.

Acknowledgments

We thank Kayoko Muromachi for her help in the field evaluations. Research grants from the Ministry of Agriculture, Forestry and Fisheries of Japan (MAFF) and Japan Science and Technology Corporation (JST) are gratefully appreciated.

References

1. Ayoub, M. et al. (2002) QTLs affecting kernel size and shape in a two-rowed by six-rowed barley cross. *Theor. Appl. Genet.*, **105**, 237–247.
2. Basten, C. J., Weir, B. S. & Zeng, Z. B. (2000) QTL Cartographer, Version 1.14. Department of Statistics, North Carolina State University, Raleigh, North Carolina, USA.
3. Hayes, P. M. et al. (1993) Quantitative trait locus effects and environmental interaction in a sample of North American barley germplasm. *Theor. Appl. Genet.*, **87**, 392–401.
4. Kicherer, S. et al. (2000) Localising QTLs for leaf rust resistance and agronomic traits in barley (*Hordeum vulgare* L.). *Theor. Appl. Genet.*, **100**, 881–888.
5. Kjaer, B. & Jensen, J. (1996) Quantitative trait loci for grain yield and yield components in a cross between a six-rowed and a two-rowed barley. *Euphytica*, **90**, 39–48.
6. Kleinhofs, A. (2002) Integrating molecular and morphological/physiological marker maps. *Barley Genet. Newsl.*, **32**, 152–159.
7. Komatsuda, T., Annaka, T. & Oka, S. (1993) Genetic mapping of a quantitative trait locus (QTL) that enhances the shoot differentiation rate in *Hordeum vulgare* L. *Theor. Appl. Genet.*, **86**, 713–720.
8. Komatsuda, T. et al. (1998) Development of STS markers closely linked to the *vrs1* locus in barley, *Hordeum vulgare*. *Genome*, **41**, 680–685.
9. Kosambi, D. D. (1944) The estimation of map distance from recombination values. *Ann. Eugen.*, **12**, 172–175.
10. Laurie, D. A. et al. (1995) RFLP mapping of five major genes and eight quantitative trait loci controlling flowering time in a winter $\times$ spring barley (*Hordeum vulgare* L.) cross. *Genome*, **38**, 575–585.
11. Mano, Y. et al. (1999) Map construction of sequence-tagged sites (STSs) in barley (*Hordeum vulgare* L.). *Theor. Appl. Genet.*, **98**, 937–946.
12. Mano, Y. et al. (2001) Construction of a genetic map of barley (*Hordeum vulgare* L.) cross ‘Azumamugi’ $\times$ ‘Kanto Nakate Gold’ using a simple and efficient amplified fragment-length polymorphism system. *Genome*, **44**, 284–292.
13. Mano, Y. & Komatsuda, T. (2002) Identification of QTLs controlling tissue-culture traits in barley (*Hordeum vulgare* L.). *Theor. Appl. Genet.*, **105**, 708–715.
14. Marquez-Cedillo, L. A. et al. (2000) QTL analysis of malting quality in barley based on the doubled-haploid progeny of two elite North American varieties representing different germplasm groups. *Theor. Appl. Genet.*, **101**, 173–184.
15. Marquez-Cedillo, L. A. et al. (2001) QTL analysis of agronomic traits in barley based on the doubled haploid progeny of two elite North American varieties representing different germplasm groups. *Theor. Appl. Genet.*, **103**, 625–637.
16. Qi, X., Stam, P. & Lindhout, P. (1996) Comparison and integration of four barley genetic maps. *Genome*, **39**, 379–394.
17. Sameri, M. & Komatsuda, T. (2004) Identification of quantitative trait loci (QTLs) controlling heading time in the population generated from a cross between oriental and occidental barley cultivars (*Hordeum vulgare* L.). *Breed. Sci.*, **54**, 327–332.
18. Sameri, M., Takeda, K. & Komatsuda, T. (2006) Quantitative trait loci controlling agronomic traits in recombinant inbred lines from a cross between oriental- and occidental-type barley cultivars. *Breed. Sci.*, **56**, 243–252.
19. Sayed-Tabatabaei, B. E. et al. (1999) Sequence-tagged sites for barley genome mapping. *Bull. Natl. Inst. Agrobiol. Resour.*, **13**, 11–22.
20. Takahashi, R. (1951) Studies on the classification and geographical distribution of the Japanese barley varieties. II. Correlative inheritance of some quantitative characters with the ear type. *Ber. Ohara Inst. Landw. Forsch.*, **9**, 383–398.
21. Takahashi, R. (1955) The origin and evolution of cultivated barley. In *Advances in genetics* 7, ed. Demerec, M., Academic Press, New York, 227–266.
22. Takahashi, R. & Yasuda, S. (1956) Genetic studies of spring and winter habit of growth in barley. *Ber. Ohara Inst. Landw. Biol. Okayama Univ.*, **10**, 245–308.
23. Takahashi, R., Hayashi, J. & Moriya, I. (1963) The effect of the *uzu* gene on productive traits in barley. I. Evaluation of the genetic effects in different genetic backgrounds. *Ber. Ohara Inst. Landw. Biol. Okayama Univ.*, **12**, 24–42.
24. Takahashi, R., Hayashi, J. & Moriya, I. (1975) Studies on barley breeding with two-rowed and six-rowed varietal crosses. I. Effects of the genes *V* and *v* on agronomic characters. *Jpn. J. Breed.*, **25**, 334–342 [In Japanese with English summary].