REVIEW

Interaction between *Torymus sinensis* (Hymenoptera: Torymidae) and *T. beneficus*, Introduced and Indigenous Parasitoids of the Chestnut Gall Wasp *Dryocosmus kuriphilus* (Hymenoptera: Cynipidae)

Kaori YARA*

Biodiversity Division, National Institute for Agro-Environmental Sciences (Tsukuba, Ibaraki 305–8604, Japan)

Abstract

Torymus sinensis is a parasitoid wasp that was introduced from China to Japan to control the invasive chestnut gall wasp *Dryocosmus kuriphilus*. Interaction between *T. sinensis* and the indigenous parasitoid *T. beneficus* has been of interest since *T. sinensis* was first released in chestnut orchards, as hybridization between them might impede the success of biological control by the introduced parasitoid. Such apprehensions disappeared with the drastic decrease in damage caused by *D. kuriphilus*. However, the emergence of morphologically intermediate individuals between them after the introduction of the parasitoid has triggered renewed interest in hybridization, specifically on the non-target effects of biological control. In this review, the interaction between both species, specifically hybridization, species composition and species displacement, is surveyed. Molecular markers are used, not only because *T. sinensis* and *T. beneficus* are similar morphologically, but also because the latter has two emergence strains (early- and late-spring). Surveys show that 1) *T. sinensis* has displaced both emergence strains of *T. beneficus*; 2) hybridization between *T. sinensis* and early-spring *T. beneficus* occurs at a low frequency ($\leq 1\%$); but that 3), hybrid F_1 s between *T. sinensis* and late-spring *T. beneficus* occur at a much higher frequency ($\leq 20\%$). A study of phylogenetic relationships indicates that the late-spring *T. beneficus* is closer to the different species *T. sinensis* than the early-spring *T. beneficus*. This can partly explain why *T. sinensis* hybridizes more readily with the late-spring *T. beneficus* than with the early-spring *T. beneficus*.

Discipline: Plant protection

Additional key words: classical biological control, displacement, hybridization, molecular marker, non-target effect

Introduction

The chestnut gall wasp *Dryocosmus kuriphilus* Yasumatsu, thought to have invaded from China in the 1940s, is a serious pest affecting chestnut trees in Japan. Although the indigenous parasitoid *Torymus beneficus* Yasumatsu et Kamijo was experimentally released in chestnut orchards¹⁶, it was unable to suppress *D. kuriphilus* populations. The parasitoid *Torymus sinensis* Kamijo was then imported from China into Japan and released at 64 sites from 1982 to 1999¹⁴. This introduced parasitoid

very effectively reduced the damage caused by *D. kuriphilus*^{5,10,13}. This project is regarded as one of the most famous and successful cases of classical biological control in Japan. However, the ecological characteristics of *T. sinensis* and *T. beneficus* remain poorly understood, partly due to the difficulty of rearing them in the laboratory and, as mentioned below, that of discriminating morphologically between them. As for *T. beneficus*, for example, it is not yet clear what gall insects they exploited before the invasion of *D. kuriphilus* in Japan; *T. sinensis* is thought to be a specialist parasitoid of *D. kuriphilus*⁹.

Interaction between *T. sinensis* and *T. beneficus* has

Present address:

Tea Research Station, NARO Institute of Vegetable and Tea Science (Shimada, Shizuoka 428–8501, Japan)

*Corresponding author: e-mail yara@affrc.go.jp

Received 27 December 2012; accepted 7 June 2013.

attracted attention since the first release of *T. sinensis* in 1982: their morphological and ecological characteristics are so similar that researchers were concerned that hybridization between them might impair the effectiveness of biological control by the introduced parasitoid, e.g. through the increase in sterile individuals. Such apprehensions subsided in the face of evidence that damage by *D. kuriphilus* plummeted in chestnut orchards⁶. However, the appearance of individuals morphologically intermediate between *T. sinensis* and *T. beneficus* was reported in the 1990s^{1,7}, which sparked a new interest in hybridization, specifically the non-target effects of biological control.

Recent progress in the use of molecular markers has elicited a better understanding of insect interactions, especially hybridization, between closely-related species that are morphologically very similar. This also applies for the case of *T. sinensis* and *T. beneficus*¹⁷. Only adult females of both species can be discriminated empirically, based on a slight difference in either the length of the ovipositor or the ratio of the length of the ovipositor sheath to the thorax; but the males cannot be empirically discriminated^{7,11,21}. Furthermore, because *T. beneficus* has two emergence strains, namely early-spring and late-spring^{8,11}, molecular markers could provide a more detailed insight into interactions between these two species/strains.

In this review, I will discuss the interaction between *T. sinensis* and *T. beneficus*, especially hybridization, species composition and species displacement, as revealed by the use of molecular markers.

Molecular markers for *T. sinensis* and both emergence strains of *T. beneficus*

1. Isozyme markers

When morphologically intermediate individuals between both *Torymus* parasitoids appeared in the field, isozyme (or allozyme) markers were initially used to discriminate them. In *T. sinensis* and *T. beneficus* (including both emergence strains), a total of 25 isozymes were tested for available markers, only three of which (malic enzyme, adenylic acid, and aspartate aminotransferase) proved usable to discriminate between early-spring *T. beneficus* and the other parasitoid species/strain (*T. sinensis* and late-spring *T. beneficus*)^{2,3} (Noda et al., unpublished data). However, even these three isozymes cannot discriminate between *T. sinensis* and the late-spring *T. beneficus*. Hybridization between *T. sinensis* and the early-spring *T. beneficus* was therefore analyzed using an isozyme (malic enzyme), focusing on sites where the early-spring *T. beneficus* dominated.

2. DNA markers

Later, as replacing malic enzyme, the internal transcribed spacer 2 (ITS2) of the rRNA coding region of nuclear DNA and the cytochrome oxidase subunit I (COI) region of mtDNA were developed as discrimination markers between *T. sinensis* and *T. beneficus*^{18,20}. Fragment analysis, which makes it possible to detect slight differences in the length of PCR products, is used as the ITS2 marker¹⁸, while PCR-RFLP is used as the COI marker²⁰. Both ITS2 and COI are superior to malic enzyme in their ability to discriminate, especially when used together. Neither, however, can clearly analyze hybridization of the *Torymus* parasitoids, since (1) the ITS2 marker cannot discriminate between some *T. sinensis* individuals and the late-spring *T. beneficus*; and (2) the COI marker on mtDNA is maternally inherited.

Subsequently, the internal transcribed spacer 1 (ITS1) of the rRNA coding region of nuclear DNA was developed²³. Now, three kinds of allele-specific PCR, using four designed specific primers, make it possible to distinguish *Torymus* species, the emergence strains of *T. beneficus*, and F₁ (first filial) individuals.

Hybridization between *T. sinensis* and *T. beneficus*

1. The early-spring *T. beneficus*

The frequency of the malic enzyme genotype of the F₁ between *T. sinensis* and the early-spring *T. beneficus* was 13% (n = 15) in Tsukuba, Ibaraki Prefecture in 1992³, and 8% (n = 93) on the Oki Islands of Shimane Prefecture in 1996¹⁵. The F₁ genotype frequency was 1% (n = 821) in specimens collected from 20 populations at 11 sites in the Tokyo Metropolis and Nagano, Ibaraki, Saitama, and Gunma prefectures from 1991 to 1995¹⁹. These studies demonstrate that hybridization had, in fact, occurred between *T. sinensis* and the early-spring *T. beneficus*, although its frequency was much lower than we expected.

Adult female specimens (n = 418) collected from Obuse, Nagano Prefecture during 1993–2001, were analyzed using both ITS2 and COI²⁰. From 1993 to 1995, all specimens were the early-spring *T. beneficus* (Fig. 1). *Torymus sinensis* was first detected in 1996; the early-spring *T. beneficus* eventually appeared to be displaced by *T. sinensis*. Simultaneously with this displacement, we found only one individual descendant of the F₁ hybrid; it had both genotypes of *T. sinensis* in ITS2 and one of the early-spring *T. beneficus* in COI. Hybridization between them was thus rare, and thus did not appear to be the main factor driving the displacement of the early *T. beneficus*. *Torymus sinensis* excels early-spring *T. beneficus* in reproductive ability¹².

The larvae of *Torymus* parasitoids ($n = 448$), collected from the same sampling site described as above (Obuse) from 1996 to 1998 and in 2000, were also analyzed using ITS2 (Fig. 2). The frequency of the F_1 genotype between *T. sinensis* and the early-spring *T. beneficus* exceeded that at the adult stage, suggesting that (1) hybridization between *T. sinensis* and the early-spring *T. beneficus* did in fact occur, but that (2) most of these F_1 s larvae did not survive to the adult stage.

2. The late-spring *T. beneficus*

From the samples above (adult female specimens ($n = 418$) collected from Obuse, Nagano Prefecture during 1993-2001), there were also 10 individuals with both the genotype of *T. sinensis* in ITS2 and that of the late-spring *T. beneficus* in COI²⁰. This is the first report of hybridization occurring in the field between *T. sinensis* and the late-spring *T. beneficus*. It also suggests that the late-spring *T. beneficus* hybridized with *T. sinensis* elsewhere and that the hybrids or their descendants migrated to the research site, since originally only the early-spring *T. beneficus* had been present. These results led to research into possible interaction with late-spring *T. beneficus*. Reanalysis using ITS1 showed 49 individuals to be hybrids (F_1 and their descendants) with the late-spring *T. beneficus* (Fig. 1).

Adult female *Torymus* parasitoids ($n = 200$), collected from Tobu, Nagano Prefecture from 1993-1996, were

Fig. 1. Annual changes in the frequency of types of nDNA ITS2, ITS1 and mtDNA COI in adult female *Torymus* parasitoids in Obuse, 1993-2001

Ts: *T. sinensis*; TbE: early-spring *T. beneficus*; TbL: late-spring *T. beneficus*; F1(Ts&TbE): hybrid between *T. sinensis* and early-spring *T. beneficus*; F1(Ts&TbL): hybrid between *T. sinensis* and late-spring *T. beneficus*²⁰.

analyzed using ITS1²¹. Increasing numbers of F_1 hybrids were detected every year from 1993 to 1996, reaching 22% (Fig. 3). It appears that *T. sinensis* can hybridize more readily with the late-spring *T. beneficus* than with the early-spring *T. beneficus*. The morphological indices of F_1 hybrids identified using the ITS1 marker ranged be-

Fig. 2. Annual changes in the frequency of ITS2 type in larvae of *Torymus* parasitoids in Obuse, 1996-2000

Ts: *T. sinensis*; TbE: early-spring *T. beneficus*; F1(Ts&TbE): hybrid between *T. sinensis* and early-spring *T. beneficus*; unidentified 1: *T. sinensis* or late-spring *T. beneficus*; unidentified 2: *T. sinensis* or hybrid between *T. sinensis* and late-spring *T. beneficus*.

Fig. 3. Annual changes in frequency of ITS1 type in adult females of *Torymus* parasitoids in Tobu, 1993-1996

Ts: *T. sinensis*; TbL: late-spring *T. beneficus*; F1(Ts&TbL): hybrid between *T. sinensis* and the late-spring *T. beneficus*²¹.

tween indices of *T. sinensis* and the late-spring *T. beneficus*, with considerable overlap. Simultaneously with the occurrence of the F_1 hybrids, the indigenous late-spring *T. beneficus* was rapidly displaced by *T. sinensis*. These facts suggest that hybridization is a more significant factor in the displacement of the late-spring *T. beneficus* by *T. sinensis* than in the displacement of the early-spring *T. beneficus*.

Phylogenetic relationship

A phylogenetic tree was constructed to examine the genetic relationships between *T. sinensis* and *T. beneficus* (early- and late-spring strains) collected from various places in Japan, China, and South Korea, using mtDNA COI sequences (1129bp)¹⁷ (Fig. 4). There were two distinct groups in the phylogenetic tree obtained: A and B. Group A consisted of two subgroups: A1 and A2 (Fig. 5). *Torymus sinensis* belonged to Subgroup A1 within Group A, while Subgroup A2 consisted of late-spring *T. beneficus*. Early-spring *T. beneficus* belonged to Group B. These results show that the late-spring *T. beneficus* is closer to the different species *T. sinensis* than the early-spring *T.*

beneficus. This may partly explain why *T. sinensis* hybridizes more easily with the late-spring *T. beneficus* than with the early-spring *T. beneficus*.

Further remarks

Recent progress in the use of molecular markers has revealed complicated interactions among *Torymus* species/strains. For example, our study has shown their unexpected interactions in a Japanese chestnut orchard in Obuse of Nagano Prefecture where *T. koreanus*, endemic in Korea, was recently discovered⁴; at the site, early-spring and late-spring *T. beneficus* were almost completely displaced by *T. sinensis* for several years, whereas this was not the case for *T. koreanus* (Table 1). Further progress in molecular phylogenetic studies will shed light on the underlying mechanisms of the complicated interactions among *Torymus* species/strains and the patterns of non-target effects of classical biological control.

Acknowledgements

We wish to express our gratitude to Dr. T. Shimoda

Fig. 4. Collection sites of *Torymus* spp. to analyze the phylogenetic relationship

(National Agricultural Research Center) for helpful comments and suggestions on the manuscript. We appreciate both anonymous reviewers' comments that improved this article immensely. Our studies reviewed in this article were partly supported by a Grant-in-Aid (Bio Cosmos Program BCP-00-I-B-1) from the Ministry of Agriculture, Forestry and Fisheries, and a Grant-in-Aid for Scientific Research (No. 19380037) from the Japan Society for the Promotion of Science.

References

1. Aoto, I. & Murakami, Y. (1992) Dispersion of a *Torymus* (*Syntomaspis*) *sinensis* population in Fukuoka Prefecture (Hymenoptera: Torymidae). *Proc. Assoc. Pl. Prot. Kyushu*, **38**, 193–196 [In Japanese with English summary].
2. Izawa, H. et al. (1992) Isozyme discrimination between an imported parasitoid wasp, *Torymus sinensis* Kamijo and its sibling species, *T. beneficus* Yasumatsu et Kamijo (Hymenoptera: Torymidae), attacking *Dryocosmus kuriphilus* Ya-

Fig. 5. Phylogenetic tree inferred from 1,129-bp sequences of the mitochondrial COI gene of *Torymus* spp. using the neighbor-joining method

Ts: *Torymus sinensis*; TbE: early-spring *T. beneficus*; TbL: late-spring *T. beneficus*. Each OTU (operational taxonomic unit) is denoted by the collection site followed by the number of specimens. *Torymus geranii* and *T. celticolus* was used as outgroups. Bootstrap probabilities (500 replicates) are provided on each branch if they exceed 70%. The scale of distances is shown under the tree¹⁷.

Table 1. Composition of *Torymus* parasitoids collected from galls on *Dryocosmus kuriphilus* in Obuse²²

Year	n	<i>Torymus</i> species ^a (%)					
		Tk	TbE	TbL	Ts	F ₁ (Ts&TbE)	F ₁ (Ts&TbL)
1992	24	20.8	58.3	20.8	0.0	0.0	0.0
1994	24	12.5	54.2	16.7	8.3	4.1	4.1
1996	24	16.7	8.3	0.0	58.3	0.0	16.7

a Tk: *Torymus koreanus*; TbE: early-spring *T. beneficus*; TbL: late-spring *T. beneficus*; Ts: *T. sinensis*; F₁(Ts&TbE): F₁ hybrid between *T. sinensis* and early-spring *T. beneficus*; F₁(Ts&TbL): F₁ hybrid between *T. sinensis* and late-spring *T. beneficus*.

- sumatsu (Hymenoptera: Cynipidae). *Jpn. J. Appl. Entomol. Zool.*, **36**, 58–60 [In Japanese with English summary].
3. Izawa, H. et al. (1996) Use of malic enzyme to detect hybrids between *Torymus sinensis* and *T. beneficus* (Hymenoptera: Cynipidae) attacking *Dryocosmus kuriphilus* (Hymenoptera: Cynipidae) and possibility of natural hybridization. *Jpn. J. Appl. Entomol. Zool.*, **40**, 205–208 [In Japanese with English summary].
 4. Matsuo, K. et al. (2011) Finding of *Torymus koreanus* (Hymenoptera: Torymidae) attacking *Dryocosmus kuriphilus* (Hymenoptera: Cynipidae) in Japan. *Entomol. Sci.*, **14**, 100–102.
 5. Moriya, S. et al. (1989a) The use of *Torymus sinensis* to control chestnut gall wasp, *Dryocosmus kuriphilus*, in Japan. *FFTC Technical Bull.*, **118**, 1–12.
 6. Moriya, S. et al. (1989b) Decline of the chestnut gall wasp population, *Dryocosmus kuriphilus* Yasumatsu (Hymenoptera: Cynipidae) after the establishment of *Torymus sinensis* Kamijo (Hymenoptera: Torymidae). *Appl. Entomol. Zool.*, **24**, 231–233.
 7. Moriya, S. et al. (1992) Interspecific relationship between an introduced parasitoid, *Torymus sinensis* Kamijo, as a biological control agent of the chestnut gall wasp, *Dryocosmus kuriphilus* Yasumatsu, and an endemic parasitoid, *T. beneficus* Yasumatsu et Kamijo. *Acta Phytopathologica Entomol. Hungarica*, **27**, 479–483.
 8. Murakami, Y. (1988) Ecotypes of *Torymus* (*Syntomaspis*) *beneficus* Yasumatsu et Kamijo (Hymenoptera: Torymidae) with different seasonal prevalences of adult emergence. *Appl. Entomol. Zool.*, **23**, 81–87.
 9. Murakami, Y. (1997) *Natural Enemies of the Chestnut Gall Wasp: Approaches to Biological Control*. Kyushu University Press, Fukuoka, Japan, pp.308 [In Japanese].
 10. Murakami, Y. et al. (2001) Colonization by imported *Torymus* (*Syntomaspis*) *sinensis* Kamijo (Hymenoptera: Torymidae) of the chestnut gall wasp (Hymenoptera: Cynipidae). (7) Success in the eighteenth year after release in Kumamoto. *Kyushu Pl. Prot. Res.*, **47**, 132–134 [In Japanese with English summary].
 11. Ôtake, A. (1987) Comparison of some morphological characters among two strains of *Torymus beneficus* Yasumatsu et Kamijo and *T. sinensis* Kamijo (Hymenoptera: Torymidae). *Appl. Entomol. Zool.*, **22**, 600–609.
 12. Piao, C. & Moriya, S. (1992) Longevity and oviposition of *Torymus sinensis* Kamijo and two strains of *T. beneficus* Yasumatsu et Kamijo (Hymenoptera: Torymidae). *Jpn. J. Appl. Entomol. Zool.*, **36**, 113–118 [In Japanese with English summary].
 13. Shiga, M. (1999) Classical biological control of the chestnut gall wasp, *Dryocosmus kuriphilus*: Present status and interactions between an introduced parasitoid, *Torymus sinensis*, and native parasitoids. In *Biological Invasions of Ecosystem by Pests and Beneficial Organisms*, eds. Yano, E. et al., NIAES, Tsukuba, 175–188.
 14. Shirai, Y. et al. (1999) Records of release of *Torymus sinensis* Kamijo (Hymenoptera: Torymidae), an introduced biological control agent of chestnut gall wasp in Japan. *Bull. Natl. Inst. Fruit Tree Sci.*, **33**, 163–178 [In Japanese with English summary].
 15. Toda, S. et al. (2000) Occurrence and hybridization of two parasitoid wasps, *Torymus sinensis* Kamijo and *T. beneficus* Yasumatsu et Kamijo (Hymenoptera: Torymidae) in the Oki islands. *Appl. Entomol. Zool.*, **35**, 151–154.
 16. Torii, T. (1959) Studies on the biological control of the chestnut gall wasp, *Dryocosmus kuriphilus* Yasumatsu (Hymenoptera, Cynipidae), with particular reference to the utilization of its indigenous natural enemies. *J. Fac. Agr. Shinshu Univ.*, **2**, 71–149.
 17. Yara, K. (2004) Relationship between the introduced and indigenous parasitoids *Torymus sinensis* and *T. beneficus* (Hymenoptera: Torymidae) as inferred from mt-DNA (COI) sequences. *Appl. Entomol. Zool.*, **39**, 427–433.
 18. Yara, K. (2006) Identification of *Torymus sinensis* and *T. beneficus* (Hymenoptera: Torymidae), introduced and indigenous parasitoids of the chestnut gall wasp *Dryocosmus kuriphilus* (Hymenoptera: Cynipidae), using the ribosomal ITS2 region. *Biol. Control*, **36**, 15–21.
 19. Yara, K. et al. (2000) Detection of hybrids between introduced *Torymus sinensis* and native *T. beneficus* (Hymenoptera: Torymidae) in central Japan, using malic enzyme. *Appl. Entomol. Zool.*, **35**, 201–206.
 20. Yara, K. et al. (2007) Displacement of *Torymus beneficus* (Hymenoptera, Torymidae) by *T. sinensis*, an indigenous and introduced parasitoid of the chestnut gall wasp, *Dryocosmus kuriphilus* (Hymenoptera: Cynipidae), in Japanese chestnut fields: possible involvement in hybridization. *Biol. Control*, **42**, 148–154.
 21. Yara, K. et al. (2010) Hybridization between introduced *Torymus sinensis* (Hymenoptera: Torymidae) and indigenous *T. beneficus* (late-spring strain), parasitoids of the Asian chestnut gall wasp *Dryocosmus kuriphilus* (Hymenoptera: Cynipidae). *Biol. Control*, **54**, 14–18.
 22. Yara, K. et al. (2012) Influence of the introduced parasitoid *Torymus sinensis* (Hymenoptera: Torymidae) on *T. koreanus* and *T. beneficus* as indigenous parasitoids of the chestnut gall wasp *Dryocosmus kuriphilus* (Hymenoptera: Cynipidae) on chestnut trees in Nagano Prefecture, Japan. *Appl. Entomol. Zool.*, **47**, 55–60.
 23. Yara, K. & Kunimi, Y. (2009) Discrimination of *Torymus sinensis* Kamijo (Hymenoptera: Torymidae) and *T. beneficus* Yasumatsu et Kamijo and their hybrids by allele-specific PCR. *Appl. Entomol. Zool.*, **44**, 275–280.