Erratum

In the article “Fasting Plasma Glucose and Incidence of Diabetes --- Implication for the Threshold for Impaired Fasting Glucose: Results from the Population-Based Omiya MA Cohort Study” by Masayuki Kato., which appeared in JAT 2009, 16: 857-861, are incorrect. The editorial staff apologizes to the authors and all readers. The correct are presented with corresponding below.

Original Article

Fasting Plasma Glucose and Incidence of Diabetes --- Implication for the Threshold for Impaired Fasting Glucose: Results from the Population-Based Omiya MA Cohort Study

Masayuki Kato¹, Mitsuhiro Noda², Hiroshi Suga³, Masahiko Matsumoto³, and Yasunori Kanazawa⁴ for the Omiya MA Cohort Study Group

False

Correct

Results: The incidence of diabetes increased as the baseline FPG level increased and a similar pattern was observed irrespective of sex or age. The hazard ratios compared with subjects with FPG <85 mg/dL adjusted for possible confounding factors were 3.83 (95% confidence interval (95% CI); 2.41–6.08) for subjects with 100 to 104 mg/dL FPG and 7.97 (95% CI; 4.98–12.4) for subjects with 105 to 109 mg/dL FPG.

Results: The incidence of diabetes increased as the baseline FPG level increased and a similar pattern was observed irrespective of sex or age. The hazard ratios compared with subjects with FPG <85 mg/dL adjusted for possible confounding factors were 3.83 (95% confidence interval (95% CI); 2.41–6.08) for subjects with 100 to 104 mg/dL FPG and 7.87 (95% CI; 4.98–12.4) for subjects with 105 to 109 mg/dL FPG.