Erratum

In this article “The Ingestion of a Fructose-Containing Beverage Combined with Fat Cream Exacerbates Postprandial Lipidemia in Young Healthy Women” by Hiromi Saito, et al., which appeared in JAT 2015, 22: 85-94 are incorrect. Table 1. is corrected as below:

Original Article

The Ingestion of a Fructose-Containing Beverage Combined with Fat Cream Exacerbates Postprandial Lipidemia in Young Healthy Women

Hiromi Saito, Maiko Kato, Akihiro Yoshida and Michitaka Naito

Incorrect

Table 1. Anthropometric and clinical characteristics

Trial	F100	F90G45	F55G45	G100
Age (years)	20.7 ± 0.2			
Height (cm)	158.7 ± 1.3			
Weight (kg)	52.0 ± 1.7	51.9 ± 1.8	51.5 ± 1.7	51.9 ± 1.7
BMI (kg/m ²)	20.7 ± 0.8	20.6 ± 0.8	20.5 ± 0.8	20.7 ± 0.8
Waist (cm)	70.9 ± 1.8	70.9 ± 1.6	70.7 ± 1.6	70.8 ± 1.7
W/H	0.80 ± 0.01	0.80 ± 0.01	0.80 ± 0.01	0.80 ± 0.01
VFA (cm ²)	26.5 ± 4.5	28.2 ± 4.6	26.9 ± 4.2	25.3 ± 3.7
HbA1c (NGSP) (%)	5.3 ± 0.1	5.4 ± 0.1	5.3 ± 0.1	5.3 ± 0.1
HOMA-IR	1.3 ± 0.2	1.2 ± 0.1	1.4 ± 0.1	1.2 ± 0.2
TC (mg/dL)	162.7 ± 6.9	158.6 ± 6.4	166.3 ± 5.9	163.0 ± 6.4
LDL-C (mg/dL)	88.4 ± 4.4	85.7 ± 3.7	92.3 ± 3.9	88.0 ± 3.4
HDL-C (mg/dL)	63.8 ± 3.4	61.1 ± 3.5	63.8 ± 2.8	62.6 ± 4.0
Lp(a) (mg/dL)	18.5 ± 8.8			
apoAI (mg/dL)	149.2 ± 5.5	145.9 ± 6.0	148.4 ± 4.8	149.8 ± 7.0
apoAII (mg/dL)	25.4 ± 1.0	25.3 ± 0.9	25.3 ± 0.8	25.8 ± 1.0
apoCII (mg/dL)	2.4 ± 0.2	2.6 ± 0.3	2.5 ± 0.2	2.6 ± 0.3
apoCIII (mg/dL)	7.0 ± 0.5	6.9 ± 0.5	7.1 ± 0.5	7.3 ± 0.6
apoE (mg/dL)	3.9 ± 0.2	4.1 ± 0.3	4.0 ± 0.2	4.0 ± 0.2
SBP (mmHg)	98.2 ± 1.9	99.9 ± 1.6	98.7 ± 1.9	96.9 ± 1.2
DBP (mmHg)	61.8 ± 1.5	62.9 ± 1.1	60.6 ± 1.5	61.9 ± 1.5
PR (beats/min)	66.8 ± 2.0	68.6 ± 2.0	67.3 ± 1.6	67.4 ± 1.4

All values are presented as the mean ± SEM.

SBP: systolic blood pressure, DBP: diastolic blood pressure, PR: pulse rate

Correct

Table 1. Anthropometric and clinical characteristics

Trial	F100	F90G10	F55G45	G100
Age (years)	20.7 ± 0.2			
Height (cm)	158.7 ± 1.3			
Weight (kg)	52.0 ± 1.7	51.9 ± 1.8	51.5 ± 1.7	51.9 ± 1.7
BMI (kg/m ²)	20.7 ± 0.8	20.6 ± 0.8	20.5 ± 0.8	20.7 ± 0.8
Waist (cm)	70.9 ± 1.8	70.9 ± 1.6	70.7 ± 1.6	70.8 ± 1.7
W/H	0.80 ± 0.01	0.80 ± 0.01	0.80 ± 0.01	0.80 ± 0.01
VFA (cm ²)	26.5 ± 4.5	28.2 ± 4.6	26.9 ± 4.2	25.3 ± 3.7
HbA1c (NGSP) (%)	5.3 ± 0.1	5.4 ± 0.1	5.3 ± 0.1	5.3 ± 0.1
HOMA-IR	1.3 ± 0.2	1.2 ± 0.1	1.4 ± 0.1	1.2 ± 0.2
TC (mg/dL)	162.7 ± 6.9	158.6 ± 6.4	166.3 ± 5.9	163.0 ± 6.4
LDL-C (mg/dL)	88.4 ± 4.4	85.7 ± 3.7	92.3 ± 3.9	88.0 ± 3.4
HDL-C (mg/dL)	63.8 ± 3.4	61.1 ± 3.5	63.8 ± 2.8	62.6 ± 4.0
Lp(a) (mg/dL)	18.5 ± 8.8			
apoAI (mg/dL)	149.2 ± 5.5	145.9 ± 6.0	148.4 ± 4.8	149.8 ± 7.0
apoAII (mg/dL)	25.4 ± 1.0	25.3 ± 0.9	25.3 ± 0.8	25.8 ± 1.0
apoCII (mg/dL)	2.4 ± 0.2	2.6 ± 0.3	2.5 ± 0.2	2.6 ± 0.3
apoCIII (mg/dL)	7.0 ± 0.5	6.9 ± 0.5	7.1 ± 0.5	7.3 ± 0.6
apoE (mg/dL)	3.9 ± 0.2	4.1 ± 0.3	4.0 ± 0.2	4.0 ± 0.2
SBP (mmHg)	98.2 ± 1.9	99.9 ± 1.6	98.7 ± 1.9	96.9 ± 1.2
DBP (mmHg)	61.8 ± 1.5	62.9 ± 1.1	60.6 ± 1.5	61.9 ± 1.5
PR (beats/min)	66.8 ± 2.0	68.6 ± 2.0	67.3 ± 1.6	67.4 ± 1.4

All values are presented as the mean ± SEM.

SBP: systolic blood pressure, DBP: diastolic blood pressure, PR: pulse rate