



# Associations between Socioeconomic Status and the Prevalence and Treatment of Hypercholesterolemia in a General Japanese Population: NIPPON DATA2010

Naoko Fujiyoshi<sup>1</sup>, Hisatomi Arima<sup>1,2</sup>, Atsushi Satoh<sup>1,2</sup>, Toshiyuki Ojima<sup>3</sup>, Nobuo Nishi<sup>4</sup>, Nagako Okuda<sup>5</sup>, Aya Kadota<sup>1,6</sup>, Takayoshi Ohkubo<sup>7</sup>, Atsushi Hozawa<sup>8</sup>, Naoki Nakaya<sup>8</sup>, Akira Fujiyoshi<sup>1</sup>, Tomonori Okamura<sup>9</sup>, Hirotsugu Ueshima<sup>1,6</sup>, Akira Okayama<sup>10</sup> and Katsuyuki Miura<sup>1,6</sup>, NIPPON DATA2010 Research Group

<sup>1</sup>Department of Public Health, Shiga University of Medical Science, Shiga, Japan

<sup>2</sup>Department of Preventive Medicine and Public Health, Faculty of Medicine, Fukuoka University, Fukuoka, Japan

<sup>3</sup>Department of Community Health and Preventive Medicine, Hamamatsu University School of Medicine, Shizuoka, Japan

<sup>4</sup>International Center for Nutrition and Information, National Institute of Health and Nutrition, National Institutes of Biomedical Innovation, Health and Nutrition, Tokyo, Japan

<sup>5</sup>Department of Health and Nutrition, University of Human Arts and Sciences, Saitama, Japan

<sup>6</sup>Center for Epidemiologic Research in Asia, Shiga University of Medical Sciences, Shiga, Japan

<sup>7</sup>Department of Hygiene and Public Health Teikyo University School of Medicine, Tokyo, Japan

<sup>8</sup>Department of Preventive Medicine and Epidemiology, Tohoku Medical Megabank Organization, Tohoku University, Sendai, Japan

<sup>9</sup>Department of Preventive Medicine and Public Health, Keio University, Tokyo, Japan

<sup>10</sup>Research Institute of Strategy for Prevention, Tokyo, Japan

**Aim:** To investigate associations between socioeconomic status (SES) and the prevalence and treatment status of hypercholesterolemia in a general Japanese population.

**Methods:** In 2010, we established a cohort study of 2417 adults (age 20–91 yr) from 300 randomly selected areas across Japan who participated in the National Health and Nutrition Survey of Japan. We cross-sectionally examined an association between SES and (1) prevalence of hypercholesterolemia in 2417 participants (999 men and 1418 women) and (2) not receiving medication for hypercholesterolemia in 654 participants (215 men and 439 women). SES included employment status, marital status, length of education, and household expenditures. Hypercholesterolemia was defined as a total serum cholesterol level of  $\geq 6.21$  mmol/L (240 mg/dL) or the use of lipid-lowering medications.

**Results:** The overall prevalence of hypercholesterolemia was 21.5% in men and 31.0% in women. In men, the lowest quintile of household expenditures was associated with a higher prevalence of hypercholesterolemia (28.3%) compared with the upper 4 quintiles (19.9%) (multivariable-adjusted odds ratio 1.66; 95% confidence interval [CI] 1.16–2.38). Among participants with hypercholesterolemia, 55.4% of men and 55.1% of women were not receiving medication. Unmarried men were more likely to be untreated (75.0%) than married men (50.9%) (multivariable-adjusted odds ratio 2.53; 95%CI 1.05–6.08). SES had no significant effects in women.

**Conclusion:** In a general population of Japanese men, low household expenditures were associated with a higher prevalence of hypercholesterolemia, and unmarried men with hypercholesterolemia were less likely to receive medication.

**Key words:** Hypercholesterolemia, Socioeconomic status, Treatment, Risk factor, Japan

Copyright©2018 Japan Atherosclerosis Society

This article is distributed under the terms of the latest version of CC BY-NC-SA defined by the Creative Commons Attribution License.

## Introduction

Cardiovascular disease (CVD) is one of the leading causes of premature death in Japan as well as other regions of the world<sup>1,2</sup>. Hypercholesterolemia is one

of potential key risk factors for fatal and nonfatal CVD<sup>2-7</sup>. In Japan, however, cholesterol levels have increased significantly since the late 1960s and have reached a similar level as that in Western countries<sup>2,8-10</sup>. Effective CVD prevention requires a strategy based on the

reasons why cholesterol levels remain high in Japan.

A number of epidemiological studies have suggested that indicators of socioeconomic status (SES) such as education, household income or expenditures, marital status, and employment status were associated with cardiovascular risk factors including hypercholesterolemia<sup>11-15</sup>. Some studies also reported the heterogeneous effects of SES on serum lipid levels between men and women<sup>16, 17</sup>. Furthermore, it has been shown that low SES was associated with limited access to health-care services and preventive care<sup>18-20</sup>. However, there is limited evidence regarding separate associations between SES and the prevalence or treatment of hypercholesterolemia in men and women, particularly in Japan.

## Aim

The objective of the present analysis was to examine the associations between household expenditures, employment status, length of education, and marital status and the prevalence and treatment of hypercholesterolemia in a cross-sectional analysis of a representative, general population of Japanese men and women.

## Methods

### Study Population

In 2010, a prospective cohort study on CVDs—National Integrated Project for Prospective Observation of Non-communicable Disease And its Trends in the Aged (NIPPON DATA2010)—was established<sup>21-23</sup>. This study was performed with the National Health and Nutrition Survey of Japan (NHNS) in November 2010 (NHNS2010) and the Comprehensive Survey of Living Conditions (CSLC) in June 2010 (CSLC2010), both of which are conducted by the Ministry of Health, Labour and Welfare of Japan. Details regarding NHNS2010 and CSLC2010 have been described elsewhere<sup>24-26</sup>.

In November 2010, 8815 residents (age  $\geq 1$  yr) from 300 randomly selected districts across Japan participated in the NHNS2010 dietary survey. Among 7229 participants  $\geq 20$  yr, 3873 participants (1598 men and 2275 women) had a blood test, and 2898 (1239 men and 1659 women) agreed to participate in the baseline NIPPON DATA2010 survey, which also

included electrocardiographic analysis, urinalysis, and a questionnaire regarding CVD. Trained interviewers obtained informed consent before enrollment. Of these 2898 participants, 7 participants were excluded because data from NHNS2010 or CSLC2010 could not be merged with baseline NIPPON DATA2010 data, and 276 participants were excluded because of missing data on SES (employment status, marital status, length of education, and/or equivalent household expenditure). One hundred ninety-eight participants with a history of CVD—which was defined as stroke, myocardial infarction, or angina pectoris—were also excluded. The remaining 2417 participants were included in the present study. This study consisted of two kinds of cross-sectional analyses that used the baseline NIPPON DATA2010 survey. Analysis 1 included all 2417 participants (999 men and 1418 women) to assess the prevalence of hypercholesterolemia. Analysis 2 included 654 participants with hypercholesterolemia (215 men and 439 women) to assess the lack of hypercholesterolemia treatment. We define treatment for hypercholesterolemia as use of lipid-lowering medication(s). Written informed consent was obtained from all participants, and the Institutional Review Board of Shiga University of Medical Science (No. 22-29, 2010) approved this study.

### Lipid Measurement and Definition of Hypercholesterolemia

Casual blood samples were obtained for NHNS 2010. Serum was separated and centrifuged soon after blood coagulation, and plasma samples were collected in siliconized tubes containing sodium fluoride and shipped to a central laboratory (SRL, Tokyo, Japan) for analysis. Serum total cholesterol was measured by the cholesterol dehydrogenase—ultraviolet (UV) method and standardized by the Centers for Disease Control and Prevention/US Collaborating Center for Reference Method Laboratory Network Research in Blood Lipids<sup>27</sup>. Triglycerides were measured using enzyme methods, and high-density lipoprotein (HDL) cholesterol and low-density lipoprotein (LDL) cholesterol were measured using direct methods (Cholestest<sup>®</sup>LDL Sekisui Medical). Information on the use of medications was collected using a self-administered questionnaire. Hypercholesterolemia was defined as total cholesterol level  $\geq 6.21$  mmol/L (240 mg/dL)<sup>28</sup> and/or the use of antihypercholesterolemic agents. Information on the use of lipid-lowering medications was obtained using a questionnaire and confirmed by trained staff. Untreated hypercholesterolemia was defined as not receiving lipid-lowering medications.

Address for correspondence: Hisatomi Arima, Department of Preventive Medicine and Public Health, Faculty of Medicine, Fukuoka University, 7-45-1 Nanakuma, Zuyounnku, Fukuoka city, Fukuoka, Japan  
E-mail: harima@fukuoka-u.ac.jp

Received: September 11, 2017

Accepted for publication: November 9, 2017

**Table 1.** Characteristics of all participants and participants with hypercholesterolemia by sex, Japan, 2010

	Men		Women	
	Total ( <i>n</i> = 999)	Hypercholesterolemia ( <i>n</i> = 215)	Total ( <i>n</i> = 1418)	Hypercholesterolemia ( <i>n</i> = 439)
Age, years	59.1 (15.4)	60.7 (13.4)	57.2 (15.8)	64.7 (10.9)
Unemployed	330 (33.0)	76 (35.3)	807 (56.9)	285 (64.9)
Unmarried	186 (18.6)	40 (18.6)	368 (26.0)	127 (28.9)
Length of education < 13 years	655 (65.6)	143 (66.5)	973 (68.6)	344 (78.4)
Lowest quintile of equivalent household expenditure	198 (19.9)	56 (26.0)	279 (19.7)	86 (19.6)
Body mass index, kg/m <sup>2</sup>	23.8 (3.2)	24.3 (3.4)	22.6 (3.5)	23.4 (3.4)
Systolic BP, mmHg	135.8 (17.8)	140 (18.6)	129.1 (20.0)	136.2 (19.4)
Diastolic BP, mmHg	82.5 (10.7)	83.9 (10.8)	77.1 (10.8)	79.6 (10.4)
Total cholesterol, mmol/L	5.27 (0.87)	6.06 (1.06)	5.40 (0.92)	6.13 (1.00)
HDL cholesterol, mmol/L	1.47 (0.39)	1.53 (0.45)	1.72 (0.40)	1.76 (0.45)
LDL cholesterol, mmol/L	3.09 (0.76)	3.64 (0.95)	3.09 (0.81)	3.64 (0.92)
Triglycerides, mmol/L <sup>†</sup>	1.45 (1.00–2.15)	1.66 (1.17–2.62)	1.10 (0.75–1.59)	1.37 (0.96–1.91)
Hypertensive <sup>§</sup>	554 (55.5)	138 (64.2)	567 (40.0)	247 (56.3)
Diabetes mellitus <sup>‡</sup>	125 (12.5)	43 (20.0)	115 (8.1)	58 (13.2)
Smoking				
Ex-smoker	408 (41.1)	95 (44.4)	123 (8.7)	34 (7.8)
Current smoker	292 (29.4)	62 (29.0)	102 (7.2)	22 (5.0)
Regular exercise <sup>*</sup>	370 (37.2)	91 (42.5)	443 (31.3)	169 (38.6)
Alcohol consumption, g/week <sup>¶</sup>	63.4 (0–241.9)	51.8 (0–241.9)	0 (0–17.3)	0 (0–5.8)

Abbreviations: BP, blood pressure; HDL, high-density lipoprotein; LDL, low-density lipoprotein. Data are shown as mean (standard deviation), median (interquartile range), or *n* (%).

<sup>†</sup> Triglycerides are shown as the median and interquartile range.

<sup>§</sup> Defined as participants who met 1 or more of the following criteria: 1) systolic BP ≥ 140 mm Hg and/or diastolic BP ≥ 90 mm Hg; or 2) receiving antihypertensive medication.

<sup>‡</sup> Defined as participants who met 1 or more of the following criteria: 1) fasting plasma glucose level ≥ 126 mg/dL (≥ 7.0 mmol/L); or 2) nonfasting plasma glucose level ≥ 200 mg/dL (≥ 11.1 mmol/L); 3) hemoglobin A1c ≥ 6.5%; or 4) receiving antidiabetic medication.

<sup>\*</sup> Defined as participants who engaged in ≥ 30 minutes of exercise for ≥ 2 days/week for > 1 year.

<sup>¶</sup> Alcohol consumption is shown as the median and interquartile range.

## Socioeconomic Status

Information on SES was collected using self-administered questionnaires for NHNS2010 (employment status), CSLC2010 (monthly household expenditures in May 2010; i.e., the month before CSLC 2010), and NIPPON DATA2010 (length of education, marital status). Equivalent household expenditures were calculated as the household expenditure divided by the square root of the number of family members. SES was defined as follows: (1) employment status (unemployed or employed [including self-employed]); (2) marital status (married, unmarried [including never married, divorced, and widowed]); (3) length of education (< 13 yr [primary school and/or high school] or ≥ 13 yr); and (4) equivalent household expenditure (upper 4 quintiles or the lowest quintile [ $< ¥82,000$  JPY (approximately \$750/month)]).

## Other Risk Factors

After a minimum of 5 min of rest in a seated posi-

tion, blood pressure (BP) was measured twice with an interval of ≥ 1 min by trained survey staff using a standard mercury sphygmomanometer with an appropriately sized cuff on the right arm. The mean of the two measurements was used in further analyses. Hypertension was defined as systolic BP ≥ 140 mm Hg and/or diastolic BP ≥ 90 mm Hg or the use of antihypertensive medications. Plasma glucose was measured using the hexokinase UV method, and hemoglobin A1c (HbA1c) was measured using a latex agglutination inhibition assay according to the standardized method of the Japan Diabetes Society (JDS). HbA1c was converted to the National Glycohemoglobin Standardization Program (NGSP) value using the following formula: HbA1c (NGSP) (%) = 1.02 × HbA1c (JDS) (%) + 0.25. A blood sample was defined as a fasting blood sample if it was obtained after ≥ 8 h of fasting. Participants were classified as diabetic if they met one of the following criteria<sup>29</sup>): (1) fasting plasma glucose level ≥ 126 mg/dL (≥ 7.0 mmol/L); (2) nonfasting plasma glucose level

**Table 2.** Association between socioeconomic status and hypercholesterolemia prevalence in 999 men

Socioeconomic status	N of participants	Hypercholesterolemia, n (%)	Odds ratio (95% confidence interval)	
			Model 1	Model 2
Employment status				
Employed	669	139 (20.8)	Reference	Reference
Unemployed	330	76 (23.0)	0.96 (0.66–1.40)	0.97 (0.67–1.41)
Marital status				
Married	813	175 (21.5)	Reference	Reference
Unmarried	186	40 (21.5)	1.09 (0.73–1.62)	1.05 (0.70–1.56)
Length of education				
≥ 13 years	344	72 (20.9)	Reference	Reference
< 13 years	655	143 (21.8)	0.97 (0.69–1.35)	0.94 (0.67–1.32)
Equivalent household expenditure				
Upper 4 quintiles	801	159 (19.9)	Reference	Reference
Lowest quintile	198	56 (28.3)	1.60 (1.12–2.29)	1.66 (1.16–2.38)

Model 1 was adjusted for age. Model 2 was adjusted like Model 1, in addition to adjustments for history of diabetes mellitus, hypertension, and type of house (own or rent; this was only included in the analysis of household expenditures).

≥ 200 mg/dL (≥ 11.1 mmol/L); (3) HbA1c ≥ 6.5%; or (4) currently receiving antidiabetic medications.

### Statistical Analysis

The characteristics of the study participants are presented as the mean and standard deviation for continuous variables and as the number and percentage for categorical variables. The association between SES and each outcome was assessed using age-adjusted logistic regression (Model 1) and multivariable-adjusted logistic regression (Model 2). The multivariable-adjusted model included age, diabetes mellitus, hypertension, and type of house (owned or rented; this was only included in the analyses of equivalent household expenditures). The type of house was determined using the CSLC2010 questionnaire and used as a covariate because expenditures included household rent but not mortgages. The results of the logistic regression analyses are reported as odds ratios (ORs) with corresponding 95% confidence intervals (95% CIs). The associations between SES and each outcome in the subgroups (defined by age or menopausal status) were compared by adding an interaction term to the statistical model.  $P < 0.05$  was considered statistically significant. All statistical analyses were performed using SAS (version 9.4 for Windows; SAS Institute Inc., Cary, NC, USA).

## Results

### Participant Characteristics

The characteristics of the participants that were included in each cross-sectional analysis are shown by sex in **Table 1**. Among 999 men and 1418 women,

330 men (33.0%) and 807 women (56.9%) were unemployed, 186 men (18.6%) and 368 women (26.0%) were unmarried, 655 men (65.6%) and 973 women (68.6%) had < 13 yr of education, and 198 men (19.8%) and 279 women (19.7%) were in the lowest quintile of equivalent household expenditures. Among 215 men (21.5%) and 439 women (31.0%) with hypercholesterolemia (Analysis 2), 76 men (35.3%) and 285 women (64.9%) were unemployed, 40 men (18.6%) and 127 women (28.9%) were unmarried, 143 men (66.5%) and 344 women (78.4%) had < 13 yr of education, and 56 men (26.0%) and 86 women (19.6%) were in the lowest quintile of equivalent household expenditures. The characteristics of the participants with hypercholesterolemia are also shown by sex and SES in **Supplementary Tables 1 and 2**.

### Associations between SES and Hypercholesterolemia Prevalence in Men

**Table 2** shows the associations between SES and hypercholesterolemia prevalence in men. The lowest quintile of household expenditures was associated with more prevalent hypercholesterolemia (28.3%) compared with higher quintiles (19.9%). The OR of the lowest quintile versus the higher quintiles was 1.66 (95%CI 1.16–2.38) after adjusting for confounding factors. This association remained significant even after further adjusting for other items related to SES: the OR of the lowest quintile versus the higher quintiles was 1.70 (95%CI 1.18–2.44). There was a trend toward a larger effect of household expenditures on hypercholesterolemia prevalence in participants < 65 yr than in those ≥ 65 yr, but the difference was not statistically sig-



**Table 3.** Association between socioeconomic status and hypercholesterolemia prevalence in 1418 women

Socioeconomic status	N of participants	Hypercholesterolemia, n (%)	Odds ratios (95% confidence intervals)	
			Model 1	Model 2
Employment status				
Employed	611	154 (25.2)	Reference	Reference
Unemployed	807	285 (35.3)	0.85 (0.65–1.12)	0.85 (0.65–1.12)
Marital status				
Married	1050	312 (29.7)	Reference	Reference
Unmarried	368	127 (34.5)	1.02 (0.78–1.34)	1.00 (0.76–1.31)
Length of education				
≥ 13 years	445	95 (21.4)	Reference	Reference
< 13 years	973	344 (35.4)	1.03 (0.76–1.39)	1.01 (0.74–1.36)
Equivalent household expenditure				
Upper 4 quintiles	1139	353 (31.0)	Reference	Reference
Lowest quintile	279	86 (30.8)	0.84 (0.62–1.13)	0.82 (0.61–1.11)

Model 1 was adjusted for age. Model 2 was adjusted like Model 1, in addition to adjustments for history of diabetes mellitus, hypertension, and type of house (own or rent; this was only included in the analysis of household expenditures).

nificant ( $P$  heterogeneity=0.06) (**Supplementary Table 3**). Other SES variables were not clearly associated with hypercholesterolemia prevalence. Similar findings were obtained when different definitions for hypercholesterolemia were used (LDL-C >4.14 mmol/L and non-HDL cholesterol [NHDLC] >4.91 mmol/L) (**Supplementary Table 4**).

#### Associations between SES and Hypercholesterolemia Prevalence in Women

**Table 3** shows the associations between SES and hypercholesterolemia prevalence in women. Employment status, marital status, length of education, and household expenditures were not clearly associated with hypercholesterolemia. There were no significant differences in the associations between SES and hypercholesterolemia in the subgroups defined by age, except for length of education ( $P$  heterogeneity=0.01) (**Supplementary Table 3**). There was also no heterogeneity in the effects of SES on hypercholesterolemia between premenopausal and postmenopausal women ( $P$  heterogeneity >0.1) (**Supplementary Table 5**).

#### Associations between SES and not Receiving Hypercholesterolemia Treatment in Men

**Table 4** shows the associations between SES and untreated hypercholesterolemia in men. Unmarried men were more likely to be untreated (75.0%) than married men (50.9%). The OR of unmarried versus married men was 2.53 (95%CI 1.05–6.08) after adjusting for confounding factors. There were no significant differences in the effects of marital status on untreated hypercholesterolemia between participants <65 and

≥65 yr ( $P$  heterogeneity=0.22) (**Supplementary Table 7**). Other SES variables were not clearly associated with untreated hypercholesterolemia. Similar findings were obtained when different definitions of hypercholesterolemia were used (LDL-C >4.14 mmol/L and NHDLC >4.91 mmol/L) (**Supplementary Table 6**).

#### Associations between SES and not Receiving Hypercholesterolemia Treatment in Women

**Table 5** shows the associations between SES and untreated hypercholesterolemia in women. Employment status, marital status, length of education, and household expenditures were not clearly associated with untreated hypercholesterolemia. There were heterogeneous associations between marital status with untreated hypercholesterolemia in women <65 yr (OR 4.63; 95%CI 1.30–16.42) and women ≥65 yr (OR 0.90; 95%CI 0.50–1.61) ( $P$  heterogeneity=0.01). There were no other significant differences in the associations between SES and untreated hypercholesterolemia in the subgroups defined by age ( $P$  heterogeneity >0.52) (**Supplementary Table 7**). There was no heterogeneity in the effects of SES on untreated hypercholesterolemia between premenopausal and postmenopausal women ( $P$  heterogeneity >0.9) (**Supplementary Table 5**).

## Discussion

In the present analysis of a nationwide survey of a general Japanese population, male participants in the lowest quintile of household expenditures had a higher prevalence of hypercholesterolemia compared with men in the upper quintiles. Unmarried men were less likely

**Table 4.** Association between socioeconomic status and untreated hypercholesterolemia in 215 men

Socioeconomic status	N of participants	Untreated, n (%)	Odds ratios (95% confidence intervals)	
			Model 1	Model 2
Employment status				
Employed	139	86 (61.9)	Reference	Reference
Unemployed	76	33 (43.4)	1.31 (0.64–2.66)	1.34 (0.65–2.77)
Marital status				
Married	175	89 (50.9)	Reference	Reference
Unmarried	40	30 (75.0)	2.13 (0.90–5.07)	2.53 (1.05–6.08)
Length of education				
≥ 13 years	72	47 (65.3)	Reference	Reference
< 13 years	143	72 (50.4)	0.83 (0.43–1.59)	0.94 (0.48–1.87)
Equivalent household expenditure				
Upper 4 quintiles	159	89 (56.0)	Reference	Reference
Lowest quintile	56	30 (53.6)	0.66 (0.33–1.32)	0.67 (0.33–1.36)

Model 1 was adjusted for age. Model 2 was adjusted like Model 1, in addition to adjustments for history of diabetes mellitus, hypertension, and type of house (own or rent; this was only included in the analysis of household expenditures).

to receive “drug” treatment for hypercholesterolemia compared with married men. These associations remained significant even after controlling for confounding factors such as age, diabetes mellitus, and hypertension. In contrast, SES was not clearly associated with hypercholesterolemia prevalence or not receiving treatment for hypercholesterolemia in women. These findings would provide important scientific insights to health policy-makers who underscore attenuation of inequality in health status as well as that in SES.

### SES and Hypercholesterolemia Prevalence

To the best of our knowledge, no previous epidemiologic studies have reported inverse associations between household expenditures and hypercholesterolemia, but an observational study in Greece reported higher hypercholesterolemia prevalence in men with less education<sup>30</sup>. Possible mechanisms underlying the link between low SES and hypercholesterolemia may involve lifestyle choices. In fact, it has been reported that people from socioeconomically disadvantaged backgrounds are less likely to choose food products that are low in saturated fat<sup>31, 32</sup>. Another possible mechanism may involve the limited utilization of health services among people with lower SES. For example, employees of small companies have been shown to be less likely to participate in health checkups than were the employees of large companies<sup>33</sup>. On the basis of the findings of the present analysis and previous epidemiologic studies, men with low SES are more likely to have hypercholesterolemia; in contrast, there were no significant associations between SES and hypercholesterolemia in women. The observed sex difference was

consistent with previous reports from Japanese populations, which indicated that disadvantage in health status associated with low SES may be more evident in men than in women<sup>34, 35</sup>. The mechanisms underlying heterogeneous effects of SES between men and women may involve differences in lifestyle, occupational status (including type of occupation), access to health care, and attitude toward health information<sup>35</sup>. Interestingly, the sex difference observed in our study differs from what was found in studies conducted in Europe or the United States, where women had a stronger association than in men. However, SES and health outcome could have a complex relationship, and we agree with Kagamimori and the colleagues in their notion that “SES must be interpreted within the economic, social, demographic and cultural contexts of a specific country”<sup>34</sup>.

### SES and Hypercholesterolemia Treatment

To the best of our knowledge, this is the first study to examine the association between marital status and hypercholesterolemia treatment status in a general population. In the present analysis, married men with hypercholesterolemia were more likely to be treated than unmarried men, but such associations were only observed in women < 65 yr. These findings are in line with a previous observational study that reported that, with regard to choosing between healthy behaviors and/or compliance with medical treatment, married men are more likely to be affected by their partners than were married women<sup>36, 37</sup>. It has also been reported that never-married Japanese men have a less healthy lifestyle and subsequently higher levels of cardiovascu-

**Table 5.** Association between socioeconomic status and untreated hypercholesterolemia in 439 women

Socioeconomic status	N of participants	Untreated, n (%)	Odds ratios (95% confidence intervals)	
			Model 1	Model 2
Employment status				
Employed	154	98 (63.6)	Reference	Reference
Unemployed	285	144 (50.5)	1.24 (0.77–2.00)	1.24 (0.76–2.00)
Marital status				
Married	312	176 (56.4)	Reference	Reference
Unmarried	127	66 (52.0)	1.24 (0.77–1.99)	1.35 (0.83–2.20)
Length of education				
≥ 13 years	95	70 (73.6)	Reference	Reference
< 13 years	344	172 (50.0)	0.61 (0.35–1.07)	0.66 (0.37–1.17)
Equivalent household expenditure				
Upper 4 quintiles	353	205 (58.1)	Reference	Reference
Lowest quintile	86	37 (43.0)	0.66 (0.39–1.12)	0.66 (0.39–1.13)

Model 1 was adjusted for age. Model 2 was adjusted like Model 1, in addition to adjustments for history of diabetes mellitus, hypertension, and type of house (own or rent; this was only included in the analysis of household expenditures).

lar risk factors than do married men<sup>38</sup>). Regarding other SES items than marital status, an observational study from India reported that patients with hypercholesterolemia who had had a longer length of education were more likely to receive drug treatment<sup>39, 40</sup>). In the present study, however, length of education, employment status, and household expenditures were not clearly associated with untreated hypercholesterolemia. The reason for this discrepancy is not clear, but the lack of associations in this study of a general Japanese population may be because of the universal health insurance coverage system that allows Japanese citizens to access medical care regardless of SES<sup>41</sup>).

### Strengths and Limitations

This is the first study to demonstrate the associations between various SES items with the prevalence of untreated hypercholesterolemia based on a nationwide survey of a Japanese general population. Additionally, we conducted multivariable-adjusted analyses, whereas some preceding reports reported results after adjusting for a limited number of covariates. In the present analysis, some of the results remained significant after adjusting for age, diabetes mellitus, and hypertension, which suggests that other factors may mediate the link between SES and hypercholesterolemia. Further studies are needed to elucidate the mechanisms underlying this link.

The present analysis has several limitations. First, because of the cross-sectional nature of this study, we were unable to determine whether there was a causal relationship between SES and the prevalence or treatment of hypercholesterolemia. The second limitation

is reporting bias for SES items (i.e., underreporting of very high status or overreporting of very low status), which could have weakened the association between SES and the prevalence and treatment of hypercholesterolemia. Third, we did not assess whether treatment (i.e., use of medication) was translated into an appropriate clinical goal, which can be an important indicator according to a recent report from a multicountry survey<sup>42</sup>). Further study is needed to address this issue. Finally, the findings of the present analysis may not be generalizable to other countries where lifestyle, access to medical care, and medical insurance coverage differ from those found in Japan.

### Conclusion

In a general population of Japanese men, low household expenditures were associated with a higher prevalence of hypercholesterolemia, and unmarried men with hypercholesterolemia were less likely to receive drug treatment. The findings indicate that preventing cardiovascular events require a combination of high-risk and population strategies that focus on socioeconomic as well as traditional risk factors.

### Acknowledgements

This study was supported by Health and Labour Sciences Research Grants from the Ministry of Health, Labour and Welfare, Japan (Comprehensive Research on Life-Style Related Diseases including Cardiovascular Diseases and Diabetes Mellitus [H22-Junkankitou-Seishuu-Sitei-017, H25-Junkankitou-Seishuu-Sitei-022]).

## Conflict of Interest

All authors have disclosed their potential conflict of interest as indicated.

## The NIPPON DATA2010 Research Group

Chairpersons: Katsuyuki Miura (Center for Epidemiologic Research in Asia, Department of Public Health, Shiga University of Medical Science, Otsu, Shiga), and Akira Okayama (Research Institute of Strategy for Prevention, Tokyo)

Research members: Hirotugu Ueshima (Center for Epidemiologic Research in Asia, Shiga University of Medical Science, Otsu, Shiga), Shigeyuki Saitoh (School of Health Sciences, Sapporo Medical University, Sapporo, Hokkaido), Kiyomi Sakata (Department of Hygiene and Preventive Medicine, Iwate Medical University, Morioka, Iwate), Atsushi Hozawa (Department of Preventive Medicine and Epidemiology, Tohoku Medical Megabank Organization, Tohoku University, Sendai, Miyagi), Takehito Hayakawa (Kinugasa Research Organization, Ritsumeikan University, Kyoto), Hiroshi Yanagawa (Jichi Medical University, Shimotsuke, Tochigi), Yosikazu Nakamura (Department of Public Health, Jichi Medical University, Shimotsuke, Tochigi), Tomonori Okamura (Department of Preventive Medicine and Public Health, Keio University, Tokyo), Nobuo Nishi (International Center for Nutrition and Information, National Institute of Health and Nutrition, National Institutes of Biomedical Innovation, Health and Nutrition, Tokyo), Nagako Okuda (Department of Health and Nutrition, University of Human Arts and Sciences, Saitama), Takayoshi Ohkubo (Department of Hygiene and Public Health Teikyo University School of Medicine, Tokyo), Fumiyoshi Kasagi (Institute of Radiation Epidemiology, Radiation Effects Association, Tokyo), Yoshitaka Murakami (Department of Medical Statistics, Toho University, Tokyo), Toru Izumi (Kitasato University, Sagami, Kanagawa), Yasuhiro Matsumura (Faculty of Health and Nutrition, Bunkyo University, Chigasaki, Kanagawa), Toshiyuki Ojima (Department of Community Health and Preventive Medicine, Hamamatsu University School of Medicine, Hamamatsu, Shizuoka), Shinkan Tokudome (Department of Public Health, Nagoya City University Graduate School of Medical Sciences, Aichi), Hideaki Toyoshima (Nagoya University, Nagoya, Aichi), Hideaki Nakagawa (Medical Research Institute, Kanazawa Medical University, Kanazawa, Ishikawa), Yoshikuni Kita (Faculty of Nursing Science, Tsuruga Nursing University, Tsuruga, Fukui), Aya Kadota (Center for Epidemiologic Research in Asia, Shiga University of Medical Science, Otsu, Shiga), Akira Fujiyoshi (Department

of Public Health, Shiga University of Medical Science, Otsu, Shiga), Naomi Miyamatsu (Department of Clinical Nursing, Shiga University of Medical Science, Otsu, Shiga), Yasuyuki Nakamura (Department of Food Science and Human Nutrition, Ryukoku University, Otsu, Shiga), Katsushi Yoshita (Osaka City University Graduate School of human life science, Osaka), Yoshihiro Miyamoto (Department of Preventive Cardiology, National Cerebral and Cardiovascular Center, Suita, Osaka), Kazunori Kodama (Radiation Effects Research Foundation, Hiroshima) and Yutaka Kiyohara (Hisayama Research Institute for Lifestyle Disorders, Hisayama-cho, Fukuoka), Kazuo Ueda (Murakami Memorial Hospital, Nakatsu, Oita).

## References

- 1) GBD2015 Mortality and Causes of Death Collaborators: Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*, 2016; 388: 1459-1544
- 2) Ueshima H, Sekikawa A, Miura K, Turin TC, Takashima N, Kita Y, Watanabe M, Kadota A, Okuda N, Kadowaki T, Nakamura Y and Okamura T: Cardiovascular disease and risk factors in Asia: a selected review. *Circulation*, 2008; 118: 2702-2709
- 3) Di Angelantonio E, Sarwar N, Perry P, Kaptoge S, Ray KK, Thompson A, Wood AM, Lewington S, Sattar N, Packard CJ, Collins R, Thompson SG and Danesh J: Major lipids, apolipoproteins, and risk of vascular disease. *JAMA*, 2009; 302: 1993-2000
- 4) Zhang X, Patel A, Horibe H, Wu Z, Barzi F, Rodgers A, MacMahon S and Woodward M: Cholesterol, coronary heart disease, and stroke in the Asia Pacific region. *Int J Epidemiol*, 2003; 32: 563-572
- 5) Lewington S, Whitlock G, Clarke R, Sherliker P, Emberson J, Halsey J, Qizilbash N, Peto R and Collins R: Blood cholesterol and vascular mortality by age, sex, and blood pressure: a meta-analysis of individual data from 61 prospective studies with 55,000 vascular deaths. *Lancet*, 2007; 370: 1829-1839
- 6) Okamura T, Tanaka H, Miyamatsu N, Hayakawa T, Kadowaki T, Kita Y, Nakamura Y, Okayama A and Ueshima H: The relationship between serum total cholesterol and all-cause or cause-specific mortality in a 17.3-year study of a Japanese cohort. *Atherosclerosis*, 2007; 190: 216-223
- 7) Tanabe N, Iso H, Okada K, Nakamura Y, Harada A, Ohashi Y, Ando T and Ueshima H: Serum total and non-high-density lipoprotein cholesterol and the risk prediction of cardiovascular events - the JALS-ECC. *Circ J*, 2010; 74: 1346-1356
- 8) Ueshima H: Explanation for the Japanese paradox: prevention of increase in coronary heart disease and reduction in stroke. *J Atheroscler Thromb*, 2007; 14: 278-286
- 9) Nguyen HN, Fujiyoshi A, Abbott RD and Miura K: Epidemiology of cardiovascular risk factors in Asian countries. *Circ J*, 2013; 77: 2851-2859
- 10) NIPPON DATA80 Research Group: Risk assessment chart



- for death from cardiovascular disease based on a 19-year follow-up study of a Japanese representative population. *Circ J*, 2006; 70: 1249-1255
- 11) Luepker RV, Rosamond WD, Murphy R, Sprafka JM, Folsom AR, McGovern PG and Blackburn H: Socioeconomic status and coronary heart disease risk factor trends. The Minnesota Heart Survey. *Circulation*, 1993; 88: 2172-2179
  - 12) Fukuda Y and Hiyoshi A: Associations of household expenditure and marital status with cardiovascular risk factors in Japanese adults: analysis of nationally representative surveys. *J Epidemiol*, 2013; 23: 21-27
  - 13) Liu K, Cedres LB, Stamler J, Dyer A, Stamler R, Nanas S, Berkson DM, Paul O, Lepper M, Lindberg HA, Marquardt J, Stevens E, Schoenberger JA, Shekelle RB, Collette P, Shekelle S and Garside D: Relationship of education to major risk factors and death from coronary heart disease, cardiovascular diseases and all causes, Findings of three Chicago epidemiologic studies. *Circulation*, 1982; 66: 1308-1314
  - 14) Stamler R, Shipley M, Elliott P, Dyer A, Sans S and Stamler J: Higher blood pressure in adults with less education. Some explanations from INTERSALT. *Hypertension*, 1992; 19: 237-241
  - 15) Stamler J, Elliott P, Appel L, Chan Q, Buzzard M, Dennis B, Dyer AR, Elmer P, Greenland P, Jones D, Kesteloot H, Kuller L, Labarthe D, Liu K, Moag-Stahlberg A, Nichaman M, Okayama A, Okuda N, Robertson C, Rodriguez B, Stevens M, Ueshima H, Horn LV and Zhou B: Higher blood pressure in middle-aged American adults with less education-role of multiple dietary factors: the INTERMAP study. *J Hum Hypertens*, 2003; 17: 655-775
  - 16) Backholer K, Peters SAE, Bots SH, Peeters A, Huxley RR and Woodward M: Sex differences in the relationship between socioeconomic status and cardiovascular disease: a systematic review and meta-analysis. *J Epidemiol Community Health*, 2017; 71: 550-557
  - 17) Park SJ, Kang HT, Nam CM, Park BJ, Linton JA and Lee YJ: Sex differences in the relationship between socioeconomic status and metabolic syndrome: the Korean National Health and Nutrition Examination Survey. *Diabetes Res Clin Pract*, 2012; 96: 400-406
  - 18) Schoenborn CA: Health promotion and disease prevention: United States, 1985, Vital and Health Statistics. Series 10, No. 163. National Center for Health Statistics (Government Printing), Washington, USA, 1988
  - 19) Hart JT: The inverse care law. *Lancet*, 1971; 1: 405-412
  - 20) Murata C, Yamada T, Chen CC, Ojima T, Hirai H and Kondo K: Barriers to health care among the elderly in Japan. *Int J Environ Res Public Health*, 2010; 7: 1330-1341
  - 21) Satoh A, Arima H, Ohkubo T, Nishi N, Okuda N, Ae R, Inoue M, Kurita S, Murakami K, Kadota A, Fujiyoshi A, Sakata K, Okamura T, Ueshima H, Okayama A and Miura K: Associations of socioeconomic status with prevalence, awareness, treatment, and control of hypertension in a general Japanese population: NIPPON DATA2010. *J Hypertens*, 2017; 35: 401-408
  - 22) Kogure M, Tsuchiya N, Hozawa A, Nakaya N, Nakamura T, Miyamatsu N, Tanaka H, Wakabayashi I, Higashiyama A, Okuda N, Takashima N, Fujiyoshi A, Kadota A, Ohkubo T, Okamura T, Ueshima H, Okayama A and Miura K: Does the flushing response modify the relationship between alcohol intake and hypertension in the Japanese population? NIPPON DATA2010. *Hypertens Res*, 2016; 39: 670-679
  - 23) Michikawa T, Okamura T, Nitta H, Nishiwaki Y, Takebayashi T, Ueda K, Kadota A, Fujiyoshi A, Ohkubo T, Ueshima H, Okayama A and Miura K: Cross-sectional association between exposure to particulate matter and inflammatory markers in the Japanese general population: NIPPON DATA2010. *Environ Pollut*, 2016; 213: 460-467
  - 24) Ikeda N, Shibuya K and Hashimoto H: Improving population health measurement in national household surveys: a simulation study of the sample design of the comprehensive survey of living conditions of the people on health and welfare in Japan. *J Epidemiol*, 2011; 21: 385-390
  - 25) Ikeda N, Takimoto H, Imai S, Miyachi M and Nishi N: Data Resource Profile: The Japan National Health and Nutrition Survey (NHNS). *Int J Epidemiol*, 2015; 44: 1842-1849
  - 26) Katanoda K and Matsumura Y: National Nutrition Survey in Japan--its methodological transition and current findings. *J Nutr Sci Vitaminol (Tokyo)*, 2002; 48: 423-432
  - 27) Nakamura M, Sato S, Shimamoto T, Konishi M and Yoshiike N: Establishment of long-term monitoring system for blood chemistry data by the national health and nutrition survey in Japan. *J Atheroscler Thromb*, 2008; 15: 244-249
  - 28) National Cholesterol Education Program: Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation*, 2002; 106: 3143-3421
  - 29) Seino Y, Nanjo K, Tajima N, Kadowaki T, Kashiwagi A, Araki E, Ito C, Inagaki N, Iwamoto Y, Kasuga M, Hanafusa T, Haneda M and Ueki K: Report of the committee on the classification and diagnostic criteria of diabetes mellitus. *J Diabetes Investig*, 2010; 1: 212-228
  - 30) Benetou V, Chloutsios Y, Zavitsanos X, Karalis D, Naska A and Trichopoulos A: Total cholesterol and HDL-cholesterol in relation to socioeconomic status in a sample of 11,645 Greek adults: the EPIC study in Greece. *European Prospective Investigation into Nutrition and Cancer. Scand J Public Health*, 2000; 28: 260-265
  - 31) Turrell G, Hewitt B, Patterson C and Oldenburg B: Measuring socio-economic position in dietary research: is choice of socio-economic indicator important? *Public Health Nutr*, 2003; 6: 191-200
  - 32) Darmon N and Drewnowski A: Does social class predict diet quality? *Am J Clin Nutr*, 2008; 87: 1107-1117
  - 33) Okamura T, Sugiyama D, Tanaka T and Dohi S: Worksite wellness for the primary and secondary prevention of cardiovascular disease in Japan: the current delivery system and future directions. *Prog Cardiovasc Dis*, 2014; 56: 515-521
  - 34) Kagamimori S, Gaina A and Nasermoaddeli A: Socioeconomic status and health in the Japanese population. *Soc Sci Med*, 2009; 68: 2152-2160
  - 35) Kondo K, Ashida T, Hirai H, Misawa J and Suzuki K:

- The Relationship between Socio-economic Status and the Loss of Healthy Aging, and Relevant Gender Differences in the Japanese Older Population AGES Project Longitudinal Study.[Article in Japanese] Iryo To Shakai, 2009; 22: 19-30
- 36) Umberson D: Family status and health behaviors: social control as a dimension of social integration. *J Health Soc Behav*, 1987; 28: 306-319
  - 37) Umberson D: Gender, marital status and the social control of health behavior. *Soc Sci Med*, 1992; 34: 907-917
  - 38) Kamon Y, Okamura T, Tanaka T, Hozawa A, Yamagata Z, Takebayashi T, Kusaka Y, Urano S, Nakagawa H, Kadowaki T, Miyoshi Y, Yamato H, Okayama A and Ueshima H: Marital status and cardiovascular risk factors among middle-aged Japanese male workers: the High-risk and Population Strategy for Occupational Health Promotion (HIPOP-OHP) study. *J Occup Health*, 2008; 50: 348-356
  - 39) Gupta R, Sharma KK, Gupta BK, Gupta A, Gupta RR and Deedwania PC: Educational status-related disparities in awareness, treatment and control of cardiovascular risk factors in India. *Heart Asia*, 2015; 7: 1-6
  - 40) Rodriguez CJ, Cai J, Swett K, Gonzalez HM, Talavera GA, Wruck LM, Wassertheil-Smoller S, Lloyd-Jones D, Kaplan R and Daviglus ML: High Cholesterol Awareness, Treatment, and Control Among Hispanic/Latinos: Results From the Hispanic Community Health Study/Study of Latinos. *J Am Heart Assoc*, 2015; 4: e001867
  - 41) Ikeda N, Saito E, Kondo N, Inoue M, Ikeda S, Satoh T, Wada K, Stickley A, Katanoda K, Mizoue T, Noda M, Iso H, Fujino Y, Sobue T, Tsugane S, Naghavi M, Ezzati M and Shibuya K: What has made the population of Japan healthy? *Lancet*, 2011; 378: 1094-1105
  - 42) Chiang CE, Ferrieres J, Gotcheva NN, Raal FJ, Shehab A, Sung J, Henriksson KM and Hermans MP: Suboptimal Control of Lipid Levels: Results from 29 Countries Participating in the Centralized Pan-Regional Surveys on the Undertreatment of Hypercholesterolaemia (CEPHEUS). *J Atheroscler Thromb*, 2016; 23: 567-587

**Supplementary Table 1.** Characteristics of 999 men by socioeconomic status, Japan, 2010

	Employment status			Marital status		
	Employed ( <i>n</i> = 669)	Unemployed ( <i>n</i> = 330)	<i>P</i>	Married ( <i>n</i> = 813)	Unmarried ( <i>n</i> = 186)	<i>P</i>
Age, years	53.3 (14.5)	70.7 (9.6)	< 0.001	60.8 (53.3)	51.5 (18.6)	< 0.001
Body mass index, kg/m <sup>2</sup>	23.9 (3.3)	23.6 (3.0)	0.20	23.8 (23.9)	23.8 (3.7)	0.35
Systolic BP, mmHg	133.4 (16.9)	140.7 (18.6)	< 0.001	136.3 (133.4)	133.7 (19.3)	0.05
Diastolic BP, mmHg	82.9 (10.5)	81.7 (10.8)	0.22	82.5 (82.9)	82.6 (11.4)	0.96
Total cholesterol, mmol/L	5.30 (0.86)	5.20 (0.90)	0.08	5.25 (5.30)	5.34 (0.97)	0.35
HDL cholesterol, mmol/L	1.49 (0.37)	1.44 (0.41)	0.02	1.47 (1.49)	1.46 (0.41)	0.45
LDL cholesterol, mmol/L	3.13 (0.75)	3.00 (0.79)	0.01	3.07 (3.13)	3.18 (0.81)	0.15
Triglycerides, mmol/L <sup>†</sup>	1.48 (0.99–2.16)	1.43 (1.02–2.13)	0.95	1.47 (1.03–2.10)	1.38 (0.98–2.29)	0.92
Hypertension <sup>§</sup>	323 (48.3)	231 (70.0)	< 0.001	459 (56.5)	95 (51.1)	0.21
Diabetes mellitus <sup>‡</sup>	70 (10.5)	55 (16.7)	0.01	103 (12.7)	22 (11.8)	0.85
Smoking						
Ex-smoker	235 (35.3)	173 (52.7)	< 0.001	351 (43.5)	57 (30.7)	0.01
Current smoker	231 (34.7)	61 (18.6)		226 (28.0)	66 (35.5)	
Regular exercise <sup>*</sup>	188 (28.2)	182 (55.3)	< 0.001	303 (37.4)	67 (36.2)	0.84
Alcohol consumption, g/week <sup>¶</sup>	63.4 (0–241.9)	51.8 (0–190.1)	< 0.001	63.4 (5.8–241.9)	34.6 (0–190.1)	0.01
	Length of education			Equivalent household expenditure		
	≥ 13 years ( <i>n</i> = 344)	< 13 years ( <i>n</i> = 655)	<i>P</i>	Upper 4 quintiles ( <i>n</i> = 813)	Lowest quintile ( <i>n</i> = 186)	<i>P</i>
Age, years	52.9 (15.9)	62.3 (14.1)	< 0.001	59.2 (15.4)	58.7 (15.6)	0.59
Body mass index, kg/m <sup>2</sup>	24.0 (3.4)	23.7 (3.0)	0.35	23.8 (3.2)	23.8 (3.1)	0.97
Systolic BP, mmHg	131.3 (16.6)	138.2 (18.0)	< 0.001	135.9 (17.5)	135.7 (18.8)	0.82
Diastolic BP, mmHg	82.1 (10.4)	82.8 (10.8)	0.18	82.5 (10.6)	82.8 (10.9)	0.99
Total cholesterol, mmol/L	5.3 (0.87)	5.3 (0.87)	0.51	5.27 (0.86)	5.26 (0.93)	0.72
HDL cholesterol, mmol/L	1.5 (0.36)	1.5 (0.40)	0.27	1.47 (0.39)	1.45 (0.37)	0.79
LDL cholesterol, mmol/L	3.1 (0.75)	3.1 (0.77)	0.15	3.08 (0.74)	3.11 (0.86)	0.69
Triglycerides, mmol/L <sup>†</sup>	1.4 (0.97–2.07)	1.48 (1.04–2.18)	0.14	1.44 (1.02–2.15)	1.52 (0.96–2.15)	0.89
Hypertension <sup>§</sup>	154 (44.8)	400 (61.1)	< 0.001	443 (55.3)	111 (56.1)	0.91
Diabetes mellitus <sup>‡</sup>	33 (9.6)	92 (14.1)	0.05	107 (13.4)	18 (9.1)	0.13
Smoking						
Ex-smoker	135 (39.4)	273 (42.0)	0.003	324 (40.7)	84 (42.9)	0.21
Current smoker	85 (24.8)	207 (31.9)		228 (28.6)	64 (32.7)	
Regular exercise <sup>*</sup>	120 (34.9)	250 (38.3)	0.31	304 (38.1)	66 (33.5)	0.27
Alcohol consumption, g/week <sup>¶</sup>	51.8 (0–241.9)	80.6 (0–241.9)	0.05	241.9 (0–63.4)	241.9 (0–51.8)	0.21

Abbreviations: BP, blood pressure; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

Data are shown as mean (standard deviation), median (interquartile range), or *n* (%).<sup>†</sup> Triglycerides are shown as the median and interquartile range.<sup>§</sup> Defined as participants who met 1 or more of the following criteria: 1) systolic BP ≥ 140 mm Hg and/or diastolic BP ≥ 90 mm Hg; or 2) receiving antihypertensive medication.<sup>‡</sup> Defined as participants who met 1 or more of the following criteria: 1) fasting plasma glucose level ≥ 126 mg/dL (≥ 7.0 mmol/L); or 2) nonfasting plasma glucose level ≥ 200 mg/dL (≥ 11.1 mmol/L); 3) hemoglobin A1c ≥ 6.5%; or 4) receiving antidiabetic medication.<sup>\*</sup> Defined as participants who engaged in ≥ 30 minutes of exercise for ≥ 2 days/week for > 1 year.<sup>¶</sup> Alcohol consumption is shown as the median and interquartile range.

**Supplementary Table 2.** Characteristics of 1418 women by socioeconomic status, Japan, 2010

	Employment status			Marital status		
	Employed ( <i>n</i> = 611)	Unemployed ( <i>n</i> = 807)	<i>P</i>	Married ( <i>n</i> = 1050)	Unmarried ( <i>n</i> = 368)	<i>P</i>
Age, years	50.0 (13.9)	62.7 (15.0)	<0.001	56.5 (14.2)	59.3 (19.7)	<0.001
Body mass index, kg/m <sup>2</sup>	22.2 (3.5)	22.9 (3.5)	<0.001	22.5 (3.4)	22.8 (3.9)	0.52
Systolic BP, mmHg	123.6 (18.3)	133.1 (20.2)	<0.001	128.2 (19.2)	131.3 (22.0)	0.03
Diastolic BP, mmHg	75.7 (10.4)	78.2 (11.0)	<0.001	77.5 (10.7)	76.1 (11.2)	0.05
Total cholesterol, mmol/L	5.34 (0.91)	5.45 (0.93)	0.027	5.38 (0.90)	5.45 (1.00)	0.46
HDL cholesterol, mmol/L	1.78 (0.39)	1.68 (0.39)	<0.001	1.73 (0.38)	1.72 (0.44)	0.44
LDL cholesterol, mmol/L	3.03 (0.80)	3.13 (0.82)	0.02	3.08 (0.78)	3.11 (0.89)	0.83
Triglycerides, mmol/L <sup>†</sup>	0.99 (0.68–1.47)	1.19 (0.81–1.68)	<0.001	1.10 (0.75–1.56)	1.10 (1.10–1.36)	0.61
Hypertension <sup>§</sup>	173 (28.3)	394 (48.8)	<0.001	388 (37.0)	179 (48.6)	<0.001
Diabetes mellitus <sup>‡</sup>	33 (5.4)	82 (10.2)	0.002	78 (7.4)	37 (10.1)	0.14
Smoking						
Ex-smoker	61 (10.0)	62 (7.7)	<0.001	96 (9.2)	27 (7.4)	0.02
Current smoker	62 (10.2)	40 (5.0)		64 (6.1)	38 (10.4)	
Regular exercise <sup>*</sup>	133 (21.8)	310 (38.6)	<0.001	317 (30.3)	126 (34.2)	0.18
Alcohol consumption, g/week <sup>¶</sup>	0.0 (0–28.8)	0.0 (0–5.8)	<0.001	0.0 (0–17.3)	0.0 (0–17.3)	0.23
	Length of education			Equivalent household expenditures		
	≥ 13 years ( <i>n</i> = 445)	< 13 years ( <i>n</i> = 973)	<i>P</i>	Upper 4 quintiles ( <i>n</i> = 1139)	Lowest quintile ( <i>n</i> = 279)	<i>P</i>
Age, years	47.3 (13.8)	61.8 (14.5)	<0.001	56.6 (15.6)	59.7 (16.3)	<0.001
Body mass index, kg/m <sup>2</sup>	21.7 (3.1)	23.0 (3.6)	<0.001	22.5 (3.4)	22.8 (3.8)	0.54
Systolic BP, mmHg	121.4 (17.9)	132.5 (19.9)	<0.001	128.6 (19.8)	130.8 (20.6)	0.10
Diastolic BP, mmHg	75.4 (10.5)	77.9 (10.9)	<0.001	77.3 (10.8)	76.5 (11.0)	0.48
Total cholesterol, mmol/L	5.29 (0.94)	5.45 (0.91)	0.001	5.42 (0.92)	5.33 (0.93)	0.15
HDL cholesterol, mmol/L	1.80 (0.39)	1.69 (0.39)	<0.001	1.74 (0.40)	1.68 (0.40)	0.02
LDL cholesterol, mmol/L	3.01 (0.82)	3.13 (0.81)	0.002	3.10 (0.81)	3.02 (0.82)	0.10
Triglycerides, mmol/L <sup>†</sup>	0.93 (0.65–1.37)	1.17 (0.81–1.67)	<0.001	1.07 (0.73–1.56)	1.18 (0.78–1.70)	0.08
Hypertension <sup>§</sup>	102 (22.9)	465 (47.8)	<0.001	439 (38.5)	128 (45.9)	0.03
Diabetes mellitus <sup>‡</sup>	13 (2.9)	102 (10.5)	<0.001	87 (7.6)	28 (10.0)	0.23
Smoking						
Ex-smoker	48 (10.8)	75 (7.7)	0.10	99 (8.7)	24 (8.6)	0.96
Current smoker	27 (6.1)	75 (7.7)		83 (7.3)	19 (6.8)	
Regular exercise <sup>*</sup>	112 (25.2)	331 (34.1)	0.001	355 (31.2)	88 (31.7)	0.95
Alcohol consumption, g/week <sup>¶</sup>	0.0 (0–40.3)	0.0 (0–5.8)	<0.001	0.0 (0–17.3)	0.0 (0–17.3)	<0.001

Abbreviations: BP, blood pressure; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

Data are shown as mean (standard deviation), median (interquartile range), or *n* (%).<sup>†</sup> Triglycerides are shown as the median and interquartile range.<sup>§</sup> Defined as participants who met 1 or more of the following criteria: 1) systolic BP ≥ 140 mm Hg and/or diastolic BP ≥ 90 mm Hg; or 2) receiving antihypertensive medication.<sup>‡</sup> Defined as participants who met 1 or more of the following criteria: 1) fasting plasma glucose level ≥ 126 mg/dL (≥ 7.0 mmol/L); or 2) nonfasting plasma glucose level ≥ 200 mg/dL (≥ 11.1 mmol/L); 3) hemoglobin A1c ≥ 6.5%; or 4) receiving antidiabetic medication.<sup>\*</sup> Defined as participants who engaged in ≥ 30 minutes of exercise for ≥ 2 days/week for > 1 year.<sup>¶</sup> Alcohol consumption is shown as the median and interquartile range.



**Supplementary Table 3.** Association between socioeconomic status and hypercholesterolemia prevalence stratified by age in 999 men and 1418 women

	Men			Women		
	Odds ratio (95% confidence intervals)		<i>P</i> for interaction	Odds ratio (95% confidence intervals)		<i>P</i> for interaction
	< 65 years	≥ 65 years		< 65 years	≥ 65 years	
Employment status						
Employed	Reference	Reference		Reference	Reference	
Unemployed	1.61 (0.88–2.94)	0.91 (0.56–1.49)	0.06	0.83 (0.58–1.18)	1.05 (0.67–1.64)	0.50
Marital status						
Married	Reference	Reference		Reference	Reference	
Unmarried	1.40 (0.85–2.31)	0.71 (0.33–1.51)	0.25	1.58 (1.01–2.45)	1.20 (0.82–1.76)	0.65
Length of education						
≥ 13 years	Reference	Reference		Reference	Reference	
< 13 years	1.06 (0.69–1.62)	0.71 (0.41–1.22)	0.14	0.99 (0.68–1.45)	0.72 (0.42–1.26)	0.01
Equivalent household expenditure						
Upper 4 quintiles	Reference	Reference		Reference	Reference	
Lowest quintile	2.27 (1.43–3.61)	1.02 (0.56–1.87)	0.06	0.74 (0.46–1.21)	0.94 (0.63–1.39)	0.71

Model included age, history of diabetes mellitus, hypertension, and type of house (own or rent; this was only included in the analysis of household expenditures).

**Supplementary Table 4.** Multivariable-adjusted odds ratios and 95% confidence intervals for the effect of socioeconomic status on hypercholesterolemia according to sex

	Men ( <i>n</i> = 999)			Women ( <i>n</i> = 1418)		
	TC	LDL-C	NHDL-C	TC	LDL-C	NHDL-C
	≥ 6.21 mmol/l	≥ 4.14 mmol/l	≥ 4.91 mmol/l	≥ 6.21 mmol/l	≥ 4.14 mmol/l	≥ 4.91 mmol/l
Employment status						
Employed	Reference	Reference	Reference	Reference	Reference	Reference
Unemployed	0.97 (0.67–1.41)	1.04 (0.70–1.56)	1.12 (0.75–1.66)	0.85 (0.65–1.12)	1.02 (0.76–1.38)	0.97 (0.72–1.31)
Marital status						
Married	Reference	Reference	Reference	Reference	Reference	Reference
Unmarried	1.05 (0.70–1.56)	0.92 (0.59–1.43)	1.04 (0.68–1.59)	1.00 (0.76–1.31)	1.11 (0.83–1.49)	1.08 (0.80–1.45)
Length of education						
≥ 13 years	Reference	Reference	Reference	Reference	Reference	Reference
< 13 years	0.94 (0.67–1.32)	1.01 (0.70–1.46)	1.02 (0.71–1.45)	1.01 (0.74–1.36)	1.25 (0.89–1.76)	1.43 (1.01–2.03)
Equivalent household expenditure						
Upper 4 quintiles	Reference	Reference	Reference	Reference	Reference	Reference
Lowest quintile	1.66 (1.16–2.38)	1.51 (1.02–2.22)	1.61 (1.10–2.35)	0.82 (0.60–1.11)	0.97 (0.70–1.34)	0.90 (0.65–1.24)

Abbreviations: TC, total cholesterol; LDL-C, low-density lipoprotein cholesterol; NHDL-C, non-high-density lipoprotein cholesterol.

Model included age, history of diabetes mellitus, hypertension, and type of house (own or rent; this was only included in the analysis of household expenditures).

**Supplementary Table 5.** Association between socioeconomic status and hypercholesterolemia prevalence and untreated hypercholesterolemia in women stratified by menopausal status

	Odds ratio of hypercholesterolemia (95% confidence interval)		<i>P</i> for interaction	Odds ratio of untreated hypercholesterolemia (95% confidence interval)		<i>P</i> for interaction
	Premenopausal	Postmenopausal		Premenopausal	Postmenopausal	
Employment status						
Employed	Reference	Reference		Reference	Reference	
Unemployed	1.52 (0.75–3.06)	0.91 (0.64–1.29)	0.19	Not calculable	1.25 (0.71–2.21)	0.94
Marital status						
Married	Reference	Reference		Reference	Reference	
Unmarried	1.82 (0.83–3.96)	1.23 (0.87–1.73)	0.72	Not calculable	1.47 (0.85–2.55)	0.97
Length of education						
≥ 13 years	Reference	Reference		Reference	Reference	
< 13 years	1.27 (0.64–2.54)	0.95 (0.65–1.40)	0.38	Not calculable	0.62 (0.32–1.19)	0.98
Equivalent household expenditure						
Upper 4 quintiles	Reference	Reference		Reference	Reference	
Lowest quintile	1.63 (0.69–3.84)	0.70 (0.48–1.03)	0.14	Not calculable	0.77 (0.41–1.46)	0.99

Model included age, history of diabetes mellitus, hypertension, and type of house (own or rent; this was only included in the analysis of household expenditures).

**Supplementary Table 6.** Multivariable-adjusted odds ratio (OR) and 95% confidence interval (95%CI) for the effect of socioeconomic status on untreated hypercholesterolemia according to sex

	Men ( <i>n</i> = 215)			Women ( <i>n</i> = 439)		
	TC ≥ 6.21 mmol/l	LDL-C ≥ 4.14 mmol/l	NHDL-C ≥ 4.91 mmol/l	TC ≥ 6.21 mmol/l	LDL-C ≥ 4.14 mmol/l	NHDL-C ≥ 4.91 mmol/l
Employment status						
Employed	Reference	Reference	Reference	Reference	Reference	Reference
Unemployed	1.34 (0.65–2.77)	1.68 (0.66–4.24)	1.80 (0.79–4.10)	1.24 (0.76–2.00)	1.59 (0.87–2.93)	1.50 (0.86–2.61)
Marital status						
Married	Reference	Reference	Reference	Reference	Reference	Reference
Unmarried	2.53 (1.05–6.08)	3.03 (1.07–8.59)	2.79 (1.12–6.93)	1.35 (0.83–2.20)	2.00 (1.08–3.71)	1.82 (1.03–3.20)
Length of education						
≥ 13 years	Reference	Reference	Reference	Reference	Reference	Reference
< 13 years	0.94 (0.48–1.87)	1.06 (0.47–2.40)	0.90 (0.43–1.89)	0.66 (0.37–1.17)	0.81 (0.40–1.64)	0.97 (0.50–1.89)
Equivalent household expenditure						
Upper 4 quintiles	Reference	Reference	Reference	Reference	Reference	Reference
Lowest quintile	0.67 (0.33–1.36)	0.64 (0.27–1.53)	0.64 (0.29–1.37)	0.66 (0.39–1.13)	0.87 (0.44–1.70)	0.73 (0.39–1.37)

Abbreviations: TC, total cholesterol; LDL-C, low-density lipoprotein cholesterol; NHDL-C, non-high-density lipoprotein cholesterol.

Model included age, history of diabetes mellitus, hypertension, and type of house (own or rent; this was only included in the analysis of household expenditures).

**Supplementary Table 7.** Association between socioeconomic status and untreated hypercholesterolemia in 215 men and 439 women stratified by age

	Men			Women		
	Odds ratio (95% confidence interval)		<i>P</i> for interaction	Odds ratio (95% confidence interval)		<i>P</i> for interaction
	< 65 years	≥ 65 years		< 65 years	≥ 65 years	
Employment status						
Employed	Reference	Reference		Reference	Reference	
Unemployed	1.40 (0.45–4.34)	1.42 (0.53–3.82)	0.95	1.01 (0.50–2.04)	1.57 (0.79–3.13)	0.52
Marital status						
Married	Reference	Reference		Reference	Reference	
Unmarried	3.71 (1.12–12.33)	2.20 (0.46–10.44)	0.22	4.63 (1.30–16.42)	0.90 (0.50–1.61)	0.01
Length of education						
≥ 13 years	Reference	Reference		Reference	Reference	
< 13 years	1.05 (0.42–2.67)	0.84 (0.30–2.33)	0.82	0.91 (0.39–2.08)	0.52 (0.23–1.14)	0.56
Equivalent household expenditure						
Upper 4 quintiles	Reference	Reference		Reference	Reference	
Lowest quintile	0.72 (0.29–1.76)	0.62 (0.19–2.09)	0.71	0.51 (0.20–1.33)	0.73 (0.38–1.38)	0.72

Model included age, history of diabetes mellitus, hypertension, and type of house (own or rent; this was only included in the analysis of household expenditures).