Finding Importance for Universal Screening of Familial Hypercholesterolemia

Shinji Yokoyama

Chubu University, Aichi, Japan

See article vol. 28: 000-000

In this issue of Journal of Atherosclerosis and Thrombosis, Matsunaga et al report finding familial hypercholesterolemia (FH) in universal screening by measuring plasma low density lipoprotein cholesterol (LDLC) of school children in Kagawa Prefectural region¹⁾. Among the 15665 children subjected, 67 were screened as LDLC $\geq$ 140 mg/dL for further genetic analysis and clinical diagnosis of FH. Common causative gene mutations for FH were identified in 41 and those for sitosterolemia in 1, while Japan Atherosclerosis Society (JAS) guideline for pediatric FH²⁾ identified only 19. No homozygote was identified in this particular survey. The prevalence of the confirmed FH in this cohort is thus as least 1/382, and potentially as high as 1/220 since more than 40% of clinical FH may not be identified by common FH gene mutations³⁾. We find important messages in these results.

First, clinical diagnosis of FH was found difficult as sensitivity and specificity of the JAS guideline for pediatric FH was only 32 and 77%, respectively¹⁾. The guideline must be reviewed and revised to improve its clinical significance by collecting more data from practical application attempt like this study. It seems more sensitive and specific to screen the candidates simply by LDLC directly for genetic analysis. This approach apparently identified most of the FH patients with common mutations having seen their prevalence in the study. Second, the study demonstrated the practical feasibility to introduce nationwide universal screening for FH children, which would not only save their lives and/or dramatically improve their quality of life in the future, but also would substantially reduce the social cost for public health.

Development of coronary heart disease (CHD) is dependent on age-integrated LDLC so that more reduction of the risk can be expected by the earlier lowering LDLC in FH patients^{4, 5)}. Onset of CHD has been in fact delayed after introduction of statins into clinical use⁶⁾. However, contribution of FH is still high among Japanese CHD patients according to the recent studies, 2.7%–5.7% for all ages and 7.8%–15.4% in those under 60 year-old⁷⁻⁹⁾, indicating overall CHD risk of FH is 10 times higher than general population and would be 30 times under 60. It is, therefore, important to find FH patients as early as possible to initiate LDLC-lowering treatment.

CHD is still the top cause of death in Western world, while it remains relatively low in Japan. According to Statistics Japan, annual death by myocardial infarction (MI) and by other CHD both increased during the post-war time until it reached around 50 and 30 thousand cases (total nearly 80 thousand) toward the end of the 20th century, and then both started declining¹⁰⁾. The numbers came down to around 33 and 35 thousand (total 68 thousand), respectively, throughout the past 25 years (Fig. 1). The cost of the treatment of CHD in Japanese Healthcare System is around 750 billion yen (BY) but showing slight gradual decrease by a few percent reflecting the decreasing trend of the cases during the ten-year period from 2008 to 2018¹¹⁾. This trend is somewhat more apparent as decreasing from 190 to 170 BY for the patients under the age of 65 (Fig. 1). The loss of life by FH-CHD is therefore estimated 2000–2500 a year and the medical cost for all FH-CHD is some 25 BY a year. Accordingly, 90% of them, 1800–2250 FH lives and 22.5 BY a year, can be saved when their risk is reduced to the same level as general population by treating LDLC from their childhood⁴⁾. Instead, the cost for LDLC-lowering treatment can be around 6 BY a year assuming that all

Address for correspondence: Shinji Yokoyama, Chubu University, Aichi, Japan E-mail: syokoyam@isc.chubu.ac.jp

Received: June 15, 2021 Accepted for publication: June 19, 2021

Copyright©2021 Japan Atherosclerosis Society

This article is distributed under the terms of the latest version of CC BY-NC-SA defined by the Creative Commons Attribution License.

Advance Publication Journal of Atherosclerosis and Thrombosis

Accepted for publication: June 19, 2021 Published online: July 10, 2021

Fig. 1. Chronological profile of death by coronary heart disease (CHD) and myocardial infarction (MI) (on left axis) and the cost of medical care for CHD in all ages and under 65 (on right axis). The data are taken from Statistics Japan^{10, 11)}.

500 thousand FH heterozygotes are prescribed generic super-strong statins at the level 3 (rosuvastatin 10 mg or atorvastatin 20 mg a day), estimated 30 yen a day.

Universal screening of FH is thus an important, urgent, practical, and feasible issue for Japanese public health with respect to significant reduction of CHD, for saving lives, improving quality of life, reducing the direct medical cost and easing burden on medical resources, and eventually resulting in saving a large amount of social cost. Ministry of Health, Welfare and Labour of Japan currently enforces universal screening tests for 23 inborn metabolic diseases such as phenylketonuria, in which development of mental retardation can be prevented by introducing specific appropriate diet and life-styles¹²⁾. FH is a more common and easy-to-find metabolic disorder and its serious future outcome is well preventable. More substantial amount of social cost can be saved by enforcement of universal mass screening system for FH.

Acknowledgement

The author is grateful to Health, Labor and Welfare Sciences Research Grant for Research on Rare and Intractable Diseases and to a Grant-in-Aid from MEXT (21K11586).

Conflicts of Interest

The author has no conflict of interest to declare related to this article.

References

- 1) Matsunaga K, Mizobuchi A, Fu HY, Ishikawa S, Tada H, Kawashiri M, Yokota I, Sasaki T, Ito S, Kunikata J, Iwase T, Hirao T, Yokoyama K, Hoshikawa Y, Fujisawa T, Dobashi K, Kusaka T, Minamino T: Universal screening for familial hypercholesterolemia in children in Kagawa, Japan. *J Atheroscler Thromb*, 2021; In Press: doi: <http://doi.org/10.5551/jat.62780>
- 2) Harada-Shiba M, Ohta T, Ohtake A, Ogura M, Dobashi K, Nohara A, Yamashita S, Yokote K: Guidance for pediatric familial hypercholesterolemia 2017. *J Atheroscler Thromb*, 2018; 25: 539-553
- 3) Hori M, Ohta N, Takahashi A, Masuda H, Isoda R, Yamamoto S, Son C, Ogura M, Hosoda K, Miyamoto Y, Harada-Shiba M: Impact of LDLR and PCSK9 pathogenic variants in Japanese heterozygous familial hypercholesterolemia patients. *Atherosclerosis*, 2019; 289: 101-108
- 4) Mabuchi H, Koizumi J, Shimizu M, Takeda R: Development of coronary heart disease in familial hypercholesterolemia. *Circulation*, 1989; 79: 225-232
- 5) Nordestgaard BG, Chapman MJ, Humphries SE, Ginsberg HN, Masana L, Descamps OS, Wiklund O, Hegele RA, Raal FJ, Defesche JC, Wiegman A, Santos RD, Watts GF, Parhofer KG, Hovingh GK, Kovanen PT, Boileau C, Averna M, Borén J, Bruckert E, Catapano AL, Kuivenhoven JA, Pajukanta P, Ray K, Stalenhoef AF, Stroes E, Taskinen MR, Tybjærg-Hansen A: Familial hypercholesterolaemia is underdiagnosed and undertreated in the general population: Guidance for clinicians to prevent coronary heart disease: Consensus statement of the European Atherosclerosis Society. *Eur Heart J*, 2013; 34: 3478-3490a
- 6) Harada-Shiba M, Sugisawa T, Makino H, Abe M, Tsushima M, Yoshimasa Y, Yamashita T, Miyamoto Y, Yamamoto A, Tomoike H, Yokoyama S: Impact of statin treatment on the clinical fate of heterozygous familial hypercholesterolemia. *J Atheroscler Thromb*, 2010; 17: 667-674
- 7) Ohmura H, Fukushima Y, Mizuno A, Niwa K, Kobayashi Y, Ebina T, Kimura K, Ishibashi S, Daida H: Estimated prevalence of heterozygous familial hypercholesterolemia in patients with acute coronary syndrome. *Int Heart J*, 2017; 58: 88-94
- 8) Harada-Shiba M, Ako J, Arai H, Hirayama A, Murakami Y, Nohara A, Ozaki A, Uno K, Nakamura M: Prevalence of familial hypercholesterolemia in patients with acute coronary syndrome in Japan: Results of the EXLORE-J study. *Atherosclerosis*, 2018; 277: 362-368
- 9) Tanaka N, Teramoto T, Yokoyama S: Application of the Japanese guidelines for the diagnosis of familial hypercholesterolemia in general practice: It is to be validated in international harmonization. *J Atheroscler Thromb*, 2019; 26: 93-98
- 10) <https://www.e-stat.go.jp/dbview?sid=0003411672> (in Japanese)
- 11) <https://www.e-stat.go.jp/stat-search/files?page=1&toukei=00450032&tstat=000001020931> (in Japanese)
- 12) https://www.jsms.gr.jp/download/MHLW_MCH_20180330.pdf (in Japanese)