Oxidative Stress in Digestive Disease

Guest Editor: Yuji Naito

Oxidative Stress and Ischemia-Reperfusion Injury in Gastrointestinal Tract and Antioxidant, Protective Agents

Makoto Sasaki* and Takashi Joh

Internal Medicine and Bioregulation, Nagoya City University Graduate School of Medical Sciences, 1 Kawasumi, Mizuho, Nagoya City 467-8601, Japan

Received: 21 August, 2006; Accepted 7 September, 2006

Summary Exacerbation of hypoxic injury after reoxygenation is a crucial mechanism mediating organ injury in transplantation, and in myocardial, hepatic, gastrointestinal, cerebral, renal, and other ischemic syndromes. The occlusion and reperfusion of the splanchnic artery is a useful animal model to elucidate the mechanism of gastrointestinal injury induced by ischemia-reperfusion (I/R). Although xanthine oxidase is a major source of reactive oxygen species (ROS), which plays an important role in the I/R-induced intestinal injury, there are many other sources of intracellular ROS. Various treatment modalities have been successfully applied to attenuate the I/R injury in animal models. This review focuses on the role of oxidant stress in the mechanism of I/R injury and the use of antioxidant agents for its treatment.

Key Words: oxidant stress, ischemia-reperfusion, antioxidant

Introduction

Oxidant stress, such as that due to free radicals and/or reactive oxygen species is known to cause organ injury. A growing body of evidence indicates that oxidative stress plays an important role in the pathogenesis of many clinical conditions [1–3] involving cardiovascular diseases [4, 5], liver diseases [6, 7], lung disease [8, 9], gastrointestinal disorders [10–12], neurological disorders [13, 14], muscle damage [15], diabetes [16], and aging [17]. The involvement of free radicals in gastrointestinal injury observed after ischemia-reperfusion (I/R) has also been reported [18–20]. Superoxide and hydrogen peroxide are considered to be the major free radicals contributing to gastrointestinal injury

after I/R. Acute gastrointestinal mucosal lesions can be by abolishing these reactive oxygen species [21].

The present review focuses on the oxidant stress during ischemia and reperfusion injury in the gastrointestinal tract and the use of antioxidant agents in its treatment.

Oxidant Stress during Ischemia-Reperfusion Injury in the Gastrointestinal Tract

An important assumption that oxygen free radicals play a crucial role in the pathogenesis of I/R has been supported by many lines of evidence that superoxide dismutase (SOD)—a highly specific scavenger of superoxide—prevents I/R induced gastrointestinal injury [18, 22]. Additionally, reperfusion-induced oxidant stress in the gastrointestinal tract is supported by reduced glutathione (GSH) consumption and concomitant formation of oxidized glutathione (glutathione disulfide: GSSG) in the gastrointestinal mucosa that is subjected to I/R [23, 24]. These changes are signifi-

*To whom correspondence should be addressed.

Tel: +81-52-853-8211 Fax: +81-52-852-0952

E-mail: msasaki@med.nagoya-cu.ac.jp

cantly reversed by both SOD and allopurinol. Conversely, there are reports that the disappearance of most of the GSH occurs during the ischemic period and GSH is not oxidized during the I/R episode [25].

By using electron spin resonance spectrometry and low-level chemiluminescence, Nilsson and co-workers demonstrated that there is a burst of oxidant formation immediately after 2–5 min of reperfusion of the ischemic intestine [26, 27]. This oxidant formation after reperfusion was supposedly derived from the electron transport chains of the mitochondria [28, 29], xanthine oxidase (XO) metabolism [30, 31], endothelial cells [32, 33], prostaglandins [34], and activated neutrophils [35, 36].

Since superoxide is more efficient as a reducing agent than as an oxidizing agent, its formation is not considered to induce high levels of cytotoxicity. Spontaneous or SOD-catalyzed dismutation of superoxide ($O_2^{\cdot-}$) produces hydrogen peroxide (H_2O_2). Additionally, superoxide—a relatively low-energy radical—is responsible for the production of highly reactive and harmful hydroxyl radicals ($HO^{\cdot}$) by the Haber-Weiss reaction.

Typically, $HO^{\cdot}$ causes biological damage by stimulating the free chain reaction of the lipid side chains of the membrane phospholipids and DNA strand breakage and causing organelle and cell disruption [37].

Relative Contributions of Ischemia and Reperfusion to Mucosal Injury

The interruption of blood supply results in an ischemic injury. Paradoxically, restoration of the blood supply causes additional cell injury that is referred to as reperfusion injury. There are several lines of evidence in the oxygen radical hypothesis of gastrointestinal I/R injury that tend to support an important assumption that the tissue injury observed after reperfusion is due to the reintroduction of oxygen rather than a delayed manifestation of injury incurred during the ischemic period. Parks and Granger have demonstrated that the mucosal injury produced after 3 h ischemia followed by 1 h of reperfusion was significantly greater than that produced after 4 h of ischemia without reperfusion [38]. They also observed that 3 h of intestinal ischemia followed by 1 h of reperfusion with deoxygenated perfusate produced significantly less mucosal injury than that produced by reperfusion with oxygenated whole blood. Furthermore, the assumption that reoxygenation results in greater mucosal injury after reperfusion is supported by the consistent observation that antioxidants and inhibitors of oxy-radical formation (e.g., allopurinol) attenuate only that component of the mucosal injury that manifests after reperfusion [39]. Ates and co-workers demonstrated that the administration of antioxidants immediately before reperfusion is as effective in attenuating

the mucosal injury as it would be if the antioxidant were to be administered before ischemia [40]. Oxy-radical production during the reperfusion period is largely responsible for the injury observed in intestinal models of I/R.

Xanthine Oxidase (XO)

The concept that XO-derived oxidants mediate the intestinal injury associated with I/R was first proposed in 1981 [41]. Subsequently, several studies have supported this concept and it is now considered that XO is a major source of oxidant generation during I/R injury in the gastrointestinal tract.

In comparison with the other organs, the intestine can generate high amounts of oxidants during I/R. The intestinal mucosa has tremendous capacity to oxidize hypoxanthine via the XO, which exists in normal healthy tissues predominantly as a NAD^+ -reducing xanthine dehydrogenase (XD) [42]. Under ischemic conditions, XD rapidly converts to XO by ischemia-mediated protease, and XO is capable of reducing molecular oxygen to both $O_2^{\cdot-}$ and H_2O_2 (Fig. 1) [39]. However, there are also adverse reports that the activity of XO decreases during ischemia but increases subsequently following reperfusion [43].

Intestinal ischemia reduces cellular ATP levels rapidly and completely within 20 min while increasing the concentration of AMP and hypoxanthine [23, 44, 45]. Menguy *et al.* also reported that hemorrhagic shock resulted in a 75% reduction in ATP, a 27% reduction in ADP, and a 350% increase in AMP concentrations in the gastric mucosa within 15 min [46]. Therefore, 30-min of ischemia is sufficient to produce prolonged functional and structural changes in a rat intestine [47]. ATP depletion results in loss of ATP-dependent ion channel regulation, producing passive ion flux across the cell membrane. Increased intracellular Ca^{2+} is harmful, one of its important consequences being the activation of a calcium-dependent protease, which cleaves of XD to form

Fig. 1. Mechanism of xanthine oxidase-mediated free radical injury. Modified from Granger *et al* [39].

XO. Fructose-1,6-biphosphate (F16BP), which is a high-energy glycolytic intermediate [48], prevents the depletion of intracellular ATP during ischemia. Thus, it prevents the conversion of XD to XO [49] and inhibits free radical production, [50] thereby protecting the intestine during the I/R injury [51].

It can be estimated (assuming a tissue PaO_2 of 20 mmHg) that the XO activity measured in a cat intestine subjected to I/R injury produced superoxide and hydrogen peroxide fluxes of 1 and 40 $\text{nmol} \cdot \text{min}^{-1} \cdot \text{g}^{-1}$, respectively [39]. These rates of oxidant production are cytotoxic to cells in culture [52] and exceed the fluxes required to increase microvascular permeability. Parks *et al.* have demonstrated that intra-arterial infusion of XO (to achieve a plasma activity of ~ 2 mU/ml) in nonischemic intestines produced an increase in microvascular permeability that is comparable to that observed after I/R [53]. Furthermore, the cytotoxicity is largely prevented by SOD and dimethyl sulfoxide (DMSO), a hydroxyl radical scavenger, which supports the assumption that these oxidants mediate the increased microvascular permeability induced by arterial XO infusion.

Previous research supports the hypothesis that the intestine appears to be the most sensitive to I/R injury among the other internal organs [39, 54]. The intestine is composed of labile enterocytes that are easily injured by episodes of ischemia, and subsequent reperfusion results in further damage to the intestinal mucosa. Enterocytes that are located at the microvilli tips are more sensitive to the effect of ischemia than those present in the crypts. This is because of their location at the end of the distribution of a central arteriole and a relative lack of collateral blood flow, which can result in a lower PaO_2 in the distal enterocytes as when compared with that in the crypts [55]. Intestinal XO activity, which is one of the major sources of free radicals, is found primarily in the mucosal layer with an increasing gradient of activity from the villus base to the tip. This is an additional reason for the increased sensitivity of the villus tip to I/R injury as compared to that of the base [54]. Additionally, the differentiated enterocytes—located at the tip of the microvilli—are inherently more sensitive to ischemia-induced injury than their undifferentiated counterparts that are located at the base [56].

Nitric Oxide (NO)

The role of nitric oxide (NO) in I/R injury is still controversial. Inhibition of NO in certain models of I/R injury causes tissue dysfunction, whereas it proves beneficial in others.

NO is synthesized from L-arginine by a family of enzymes known as NO synthase (NOS). The constitutive forms of NOS (neural NOS: nNOS and endothelial NOS: eNOS) are critical to normal physiology, and the inhibition of these

enzymes causes tissue injury [57]. However, the inducible form of NOS (iNOS), which is only expressed in the response to cytokines and growth factors, produces a large amount of NO that contributes to the pathophysiology of I/R injury. Therefore, while the use of selective inhibitors of iNOS, namely, N(6)-(iminoethyl)-L-lysine (L-NIL) and ONO-1714 has been shown to be beneficial [58–60], the use of non-selective inhibitors, namely, N(G)-nitro-L-arginine methyl ester (L-NAME) has been shown to be deleterious in gastrointestinal I/R injury [61]. Additionally, iNOS knockout mice are more resistant to intestinal I/R-induced bacterial translocation and mucosal injury, further supporting the role of iNOS as an important mediator of I/R injury in the intestine [62]. The nonselective NO inhibitors such as NG-monomethyl-L-arginine (L-NMMA) and L-NAME aggravate the gastric mucosal injury due to I/R, and this effect is blocked by L-arginine [63].

Using an NO-sensitive electrode, Wada and his co-workers measured the NO concentrations in the gastric mucosa and concluded that NO has an important pathological role in acute gastric mucosal injury induced by I/R [64]. They state that the NO concentration in the gastric mucosa increases during the ischemic period; subsequently, reperfusion causes a rapid decrease to normal levels. NOS inhibitors prevent both the increase in NO concentration during the ischemic episode and the mucosal injury induced by I/R.

NO has many beneficial effects in the I/R injury scavenging of oxygen free radicals, reduction of leukocyte adhesion to the mesenteric endothelium, and maintenance of normal vascular permeability [65–67]. During the early phase of reperfusion, after ischemia, an increase in $\text{O}_2^{\cdot -}$ limits the accumulation of NO and prevents any of the beneficial effects of NO; as a result, the leukocytes tend to adhere to the endothelium [68]. Supplementation of NO using NO donors such as FK-409, molsidomine, and nitroprusside have been shown to attenuate gastrointestinal I/R injury [58, 69–71].

Endothelial Cells and Polymorphonuclear Neutrophils (PMNs)

Both endothelial cells and polymorphonuclear neutrophils (PMNs) are another potential source of oxidants in the gastrointestinal tract. NADPH oxidase is found in both these cells, and reoxygenation promotes oxidant generation by NADPH [32, 33]. Moreover, activated PMNs secrete a variety of enzymes such as myeloperoxidase (MPO) and elastase that can injure parenchymal cells and the microvasculature [72].

Intravital microscopic studies of tissue exposed to I/R injury reveal an acute inflammatory response that is characterized by increased adhesion and emigration of PMNs in postcapillary venules, microvascular permeability, and

mucosal injury [73]. Reduced PMNs and monoclonal antibodies that interfere with the adhesion and emigration of the PMNs across venules provide significant protection against I/R injury [74, 75]. Determination of tissue-associated MPO activity showed that the infiltration of PMNs in the mucosal layer during the reperfusion period is significantly more than that during ischemia [35]. The increased mucosal MPO activity after reperfusion can be significantly attenuated by the use of antioxidants. However, this attenuation of MPO activity is reflective of the blockage of the recruitment of PMNs into the mucosal layer rather than the inhibition of MPO catalytic activity [39]. These observations suggest that the adhesive interaction between PMNs and endothelial cells plays a crucial role in I/R injury and this process mainly depends on the oxidant stress.

Antioxidants: Protective Agents against Ischemia-Reperfusion Injury

Under normal conditions, the oxygen molecule undergoes a tetravalent reduction by the cytochrome system in the mitochondria to form water. However, one to two percent of the oxygen molecules that escape this pathway undergoes univalent reduction and generates oxygen-derived free radicals. Under normal conditions, these radicals are neutralized by endogenous antioxidant enzymes such as SOD and GSH, thereby having no deleterious effect on the cells [76]. However, oxidative stress occurs when the production of oxidants exceeds the capacity of the antioxidant defense systems of the cell, or when the effectiveness of the antioxidant defense system decreases. The human body has many natural antioxidants; however, not all of them are capable of protection against the attack of oxidants induced by the I/R condition.

Many animal studies have shown that antioxidant agents are useful in protection against I/R-induced gastrointestinal injury. XO inhibitors such as allopurinol, significantly reduce the severity of intestinal mucosal lesions and gastric lesions induced by I/R, [18, 19, 22, 77–80]. Enteral administration of XO inhibitors such as pterin aldehyde or folic acid also attenuates the I/R-induced increase in microvascular permeability [81]. Although XO plays a crucial role in mucosal damage during reperfusion, it is also provoke the injury caused during the ischemic episodes [22].

Table lists antioxidant agents that have been reported to be helpful in reducing I/R-induced tissue damage in the intestine and stomach separately. The agents that have been used are acetaminophen [82], allopurinol [18, 19, 22, 23, 77–80, 83–87], astragalus membranaceus [88], anti-thrombin III [89], bilirubin [90, 91], captopril [92], cimetidine [93], 5-(2-amino-ethylamino)-1-phenyl-2-pentanone (compound IA) [94], CV-6209—a platelet-activating factor antagonist [95], cystathionine [96], cyclosporine [97, 98], desferrioxamine [99–101], dimethyl sulfoxide (DMSO) [19, 83, 102–104],

edaravone [105], epidermal growth factor (EGF) [106, 107, 108], ellagic acid—one of the polyphenols [109], fullerene [110], glutamine [103, 111–113], honey [102, 114], IT-066—a novel histamine H₂-receptor antagonist (H₂RA) [115], mannitol [116, 117], melatonin and its precursor L-tryptophan [40, 118–127], methylprednisolone [116], N-acetylcysteine (NAC) [117, 128–131], nitroglycerine [132, 133], pyrrolidine dithiocarbamate (PDTC) [134, 135], pyruvate [136], rapamycin [97], rebamipide [100, 103, 137, 138], rotenone [139], selenium [140, 141], SOD [18, 19, 22, 77–79, 85, 142], sofalcone [143, 144], sucralfate [79, 145–148], T-593—an H₂RA [149, 150], 2,2,6,6-tetramethylpiperidine-1-oxyl (TPL) [151], trimetazidine [152, 153], verapamil [87, 154, 155, 156], vitamin C [116, 117, 157–159], vitamin E [116], and zinc N-(3-aminopropionyl)-L-histidine (Z-103) [21].

Intravenous administration of N-acetylcysteine (NAC), which is a free radical scavenger, has been proved to be beneficial in protection against I/R injury of intestine [117, 128]; however, there are some controversial reports in this regard [147, 160]. Some reports have demonstrated that NAC treatment resulted in significant aggravation of the gastric mucosal injury after I/R injury [147, 161]. Although intraluminal perfusion of NAC increased the hexosamine concentration, gastric mucus was significantly decreased. This resulted in the exacerbation of the mucosal injury.

Some types of H₂RA, which inhibit acid secretion by parietal cells, are potent antioxidants and it has been suggested that these exert a protective effect on I/R-induced gastric mucosal injury. Although the luminal acid was completely neutralized by NaOH instead of T-593 (an H₂RA), no reduction in the mucosal clearance was observed. This indicated that the endogenous luminal acid does not play an important role in gastric injury induced by localized I/R stress [149]. Kitano demonstrated that compared to luminal perfusion with an HCl solution alone, the luminal perfusion with a solution containing HCl and H₂RA, cimetidine (3 mmol/l) significantly reduced the total area of erosions. Additionally, cimetidine (3 mmol/l) inhibited hydroxyl radical-induced lipid peroxidation of human erythrocyte membranes by 60% *in vitro*; this indicated that cimetidine possesses a protective effect against acute gastric mucosal injury induced by I/R not only due to the suppression of gastric acid secretion but also due to its antioxidant properties when it is present at high concentrations in the intragastric environment [93]. Furthermore, Naito *et al.* [115] showed that IT-066 (an H₂RA) scavenged the superoxide and hydroxyl radicals generated by the hypoxanthine-XO system and the hydrogen peroxide-ferrous iron system, respectively, by a spin trapping method using 5,5-dimethyl-1-pyrroline-N-oxide.

Mucoprotective agents such as sofalcon, sucralfate, and rebamipide have also been suggested to have antioxidant

Table. A list of antioxidant agents used against ischemia-reperfusion injury in the intestine and stomach

Agent	Intestine			Stomach		
	Author	Outcome	Reference	Author	Outcome	Reference
Acetaminophen				Nakamoto	↓ mucosal lesion	82
Allopurinol	Parks	↓ histological damage, ↑ blood flow	38	Itoh	↓ mucosal lesion, ↓ XO	18
	Schoenberg	↓ mucosal damage	23	Perry	↓ mucosal permeability	19
	Haglund	↓ mucosal damage	77	Laudanno	↓ mucosal lesion	79
	Bilbao	↓ histological damage, ↓ mortality	78	Alarcon de la Lastra	↓ mucosal lesion, ↓ mucosal neutrophils	84
	Boros	↓ histological damage, ↑ blood flow	80	Zollei	↓ mucosal permeability, ↓ mucosal lesion, ↓ histological damage	85
	Horne	↓ mucosal damage	83			
	Hakguder	↑ motility	86			
	Kulah	↓ histological damage, ↓ MDA, ↑ GSH, ↓ LDH	87			
	Hei	↓ histological damage, ↑ NO, ↓ Endothelin-1, ↑ SOD, ↑ GSH	88			
Anti-thrombin III	Ozden	↓ histological damage, ↓ MPO, ↓ MDA	89			
Bilirubin	Ceran	↓ histological damage, ↓ MDA	90			
	Hammerman	↓ histological damage, ↓ thiobarbituric acid-reducing substances	91			
Captopril	Buyukgebiz	↓ histological damage, ↓ MDA	92			
Cimetidine				Kitano	↓ mucosal lesion	93
Compound IA	Poussios	↓ histological damage, ↓ MDA	94			
CV-6209				Yoshikawa	↓ mucosal lesion, ↓ TBARS	95
Cystathionine				Wada	↓ mucosal lesion, ↓ TBARS	96
Cyclosporine	Puglisi	↓ histological damage	97, 98			
Desferrioxamine	Balogh	↓ MDA, ↓ TBARS, ↓ 4-hydroxy-alkenals	101	Andrews	↓ histological damage	99
				Kurokawa	↓ mucosal permeability	100
				Perry	↓ mucosal permeability	19
DMSO	Horne	↓ mucosal damage	83	Ali	↓ mucosal lesion, ↓ vascular permeability, ↑ NP-SH	102
	Kojima	↓ apoptosis	103			
	Dabareiner	↓ microvascular permeability, ↓ edema	104			
Edaravone	Tomatsuri	↓ mucosal damage, ↓ histological damage, ↓ TBARS, ↓ MPO, ↓ CINC-1	105			
EGF	Berlanga	↓ intraluminal bleeding, ↓ MPO, ↓ MDA	106			
	Villa	↓ intestinal permeability, ↓ histological damage	107			
	Martin	↓ mucosal damage	108			
Ellagic acid				Iino	↓ mucosal lesion, ↓ lipid peroxidation	109
Fullerenol	Lai	↓ histological damage, ↓ MDA, ↑ GSH	110			
Glutamin	Harward	↑ GSH, conjugated diene (a by-product of lipid peroxidation)	112	Stein	↓ mucosal lesion	111
	Basoglu	↓ NO, ↑ GSH	113			
	Kojima	↓ apoptosis	103			
Honey	Koltuksuz	↓ mucosal damage, ↓ MDA	114	Ali	↓ mucosal lesion, ↓ vascular permeability, ↑ NP-SH	102
IT-066				Naito	↓ mucosal lesion, ↓ O ₂ , ↓ OH, ↓ lipid peroxidation	115
Mannitol	Gunel	↓ histological damage, ↓ MDA, ↑ GSH	116			
	Byrka-Owczarek	↓ histological damage, ↑ blood flow	117			
Melatonin (L-tryptophan)	Kazez	↓ histological damage, ↓ MDA	121	Konturek	↓ histological damage, ↑ blood flow	118, 119
	Ustundag	↑ GSH-Px, ↑ SOD, ↑ selenium	122	Brzozowski	↓ mucosal lesion	120
	Sileri	↓ bacterial translocation	126	Cabeza	↓ histological damage, ↓ XO, ↑ SOD, ↑ GSH, ↑ PGE ₂	123, 124
	Ates	↑ catalase, ↑ SOD, ↑ GSH-Px, ↓ MDA	40	Bulbuler	↓ mucosal lesion, ↓ histological damage	125
	Ozacmak	↓ MDA, ↑ contraction, ↓ histological damage, ↑ GSH	127			
	Gunel	↑ GSH	116			
Methylprednisolone						
NAC	Cuzzocrea	↓ nitrosine, ↓ PARS, ↓ MPO, ↓ MDA, ↓ P-selectin, ↓ ICAM-1	128			

Table. A list of antioxidant agents used agints ischemia-reperfusion injury in the intestine and stomach (continued)

Agent	Intestine			Stomach		
	Author	Outcome	Reference	Author	Outcome	Reference
Nitroglycerine	Sun	↓ endothelial and epithelial permeability, ↓ MPO, ↓ IL-1beta	129			
	Montero	↓ histological damage	130			
	Olanders	↓ endothelial permeability,	131			
	Byrka-Owczarek	↑ blood flow	117			
	Dun	↓ histological damage, ↓ MDA, ↓ LDH	132			
PDTC	Khanna	↓ intestinal permeability, ↓ histological damage,	133			
	Mallick	↑ blood flow, ↑ HO, ↓ LDH, ↓ histological damage	134	El Eter	↓ vascular permeability, ↓ LDH, ↓ TNF-alpha, ↓ TBARS, ↓ NF-kappaB, ↑ GSH	135
Pyruvate	Cicalese	↓ histological damage, ↓ free radical	136			
Rapamycin	Puglisi	↓ histological damage	97			
Rebamipide	Kojima	↓ apoptosis	103	Kim	↓ MPO, ↓ MDA, ↑ SOD, ↑ NOS	137
Rotenone	Ichikawa	↓ mucosal damage, ↓ TBARS, ↓ CINC-1, ↓ TNF-alpha	139	Kurokawa	↓ mucosal permeability	100
				Hiratsuka	↓ histological damage, ↓ COX2	138
Selenium	Yoshida	↓ histological damage, ↑ blood pressure	140			
	Ozturk	↓ histological damage, ↓ bacterial translocation, ↓ MDA	141			
SOD	Parks	↓ histological damage, ↑ blood flow	22	Itoh	↓ mucosal lesion	18
	Haglund	↓ mucosal damage	77	Perry	↓ mucosal permeability	19
	Bilbao	↓ histological damage	78	Laudanno	↓ mucosal lesion	79
	Riaz	↓ MDA, ↓ neutrophil rolling & adhesion	142	Zollei	↓ mucosal permeability, ↓ mucosal lesion, ↓ histological damage	85
Sofalcone				Yoshikawa	↓ mucosal lesion, ↓ lipid peroxidation	143
				Momo	↓ mucosal lesion, ↓ MDA	144
Sucralfate	Sencan	↓ histological damage, ↓ enterocyte apoptosis	148	Laudanno	↓ mucosal lesion	79
				al-Swayeh	↓ mucosal lesion, ↓ vascular permeability, ↓ superoxide, ↑ NP-SH	145
				Wada	↓ mucosal lesion, ↓ TBARS	146
				Mojzis	↓ mucosal lesion, ↑ mucus	147
T-593 (H2RA)				Seno	↓ mucosal permeability	149
				Naito	↓ mucosal lesion	150
TPL	Udassin	↓ intestinal permeability	151			
Trimetazidine	Tsimoyiannis	↓ peritoneal adhesion	152			
	Tetik	↓ histological damage, ↓ MDA, ↓ MPO	153			
Verapamil	Mocan	↓ histological damage	155	Sato	↓ mucosal lesion, ↓ lipid peroxidation	154
	Kulah	↓ histological damage, ↓ MDA, ↓ GSH	87	al-Dohayan	↓ mucosal lesion	156
Vitamin C	Nakamura	↓ histological damage, ↓ lipid peroxides, ↓ GSSH/GSH	159	Ekman	↓ mucosal bleeding, ↑ vascular patency	157, 158
	Gunel	↓ histological damage, ↓ MDA, ↓ GSH	116			
	Byrka-Owczarek	↓ histological damage, ↑ blood flow	117			
Vitamin E	Gunel	↑ GSH	116			
Z-103				Yoshikawa	↓ mucosal lesion	21

Compound 1A: 5-(2-amino-ethylamino)-1-phenyl-2-pentanone

DMSO: dimethyl sulfoxide

EGF: epidermal growth factor

NAC: N-acetylcysteine

PDTC: pyrrolidinedithiocarbamate

SOD: superoxide dismutase

TPL: 2,2,6,6-tetramethylpiperidine-1-oxyl

Z-103: zinc N-(3-aminopropionyl)-L-histidine

properties; thus, they may prevent I/R-induced mucosal injury. High concentrations of sucralfate (3–10 mg/ml) reduced the superoxide radicals generated by leukocytes or xanthine-XO, and protected erythrocyte membrane ghosts against lipid peroxidation induced by hydrogen peroxide and Fe^{2+} *in vitro* [146]. The administration of sucralfate, either before ischemia or before reperfusion, prevented the gastrointestinal mucosal injury due to I/R stress [79, 145–148]. Rebamipide also reduced gastrointestinal damage by the pretreatment of ischemic changes [100, 103, 137, 138]. Only intraluminal administration of sofalcon before the onset of ischemia has been proven to be effective in providing gastric mucosal protection against I/R injury [143, 144].

Conclusion

Oxygen-derived free radicals play a major role in the pathophysiological mechanism of ischemia-reperfusion injury of the intestine. The sources of reactive oxygen species are abundant, and the attenuation of oxidant stress minimizes the extent of the ischemia-reperfusion injury. Therefore, antioxidant agents play an important role in the treatment of intestinal injury induced by ischemia-reperfusion.

References

- [1] Cross, C.E., Halliwell, B., Borish, E.T., Pryor, W.A., Ames, B.N., Saul, R.L., McCord, J.M., and Harman, D.: Oxygen radicals and human disease. *Ann. Intern. Med.*, **107**, 526–545, 1987.
- [2] Halliwell, B. and Gutteridge, J.M.: Free radicals and antioxidant protection: mechanisms and significance in toxicology and disease. *Hum. Toxicol.*, **7**, 7–13, 1988.
- [3] Halliwell, B., Gutteridge, J.M., and Cross, C.E.: Free radicals, antioxidants, and human disease: where are we now? *J. Lab. Clin. Med.*, **119**, 598–620, 1992.
- [4] McCord, J.M.: Oxygen-derived free radicals in postischemic tissue injury. *N. Engl. J. Med.*, **312**, 159–163, 1985.
- [5] Jeroudi, M.O., Hartley, C.J., and Bolli, R.: Myocardial reperfusion injury: role of oxygen radicals and potential therapy with antioxidants. *Am. J. Cardiol.*, **73**, 2B–7B, 1994.
- [6] Comporti, M.: Lipid peroxidation and cellular damage in toxic liver injury. *Lab. Invest.*, **53**, 599–623, 1985.
- [7] Poli, G., Albano, E., and Dianzani, M.U.: The role of lipid peroxidation in liver damage. *Chem. Phys. Lipids*, **45**, 117–142, 1987.
- [8] Kalra, J., Chaudhary, A.K., and Prasad, K.: Increased production of oxygen free radicals in cigarette smokers. *Int. J. Exp. Pathol.*, **72**, 1–7, 1991.
- [9] Ryrfeldt, A., Bannenberg, G., and Moldeus, P.: Free radicals and lung disease. *Br. Med. Bull.*, **49**, 588–603, 1993.
- [10] Otamiri, T. and Sjobahl, R.: Oxygen radicals: their role in selected gastrointestinal disorders. *Dig. Dis. Sci.*, **9**, 133–141, 1991.
- [11] Simmonds, N.J. and Rampton, D.S.: Inflammatory bowel disease—a radical view. *Gut*, **34**, 865–868, 1993.
- [12] Das, D. and Banerjee, R.K.: Effect of stress on the antioxidant enzymes and gastric ulceration. *Mol. Cell. Biochem.*, **125**, 115–125, 1993.
- [13] Adams, J.D., Jr. and Odunze, I.N.: Oxygen free radicals and Parkinson's disease. *Free Radic. Biol. Med.*, **10**, 161–169, 1991.
- [14] Dickinson, M.J. and Singh, I.: Down's syndrome, dementia, and superoxide dismutase. *Br. J. Psychiatry*, **162**, 811–817, 1993.
- [15] Jackson, M.J. and O'Farrell, S.: Free radicals and muscle damage. *Br. Med. Bull.*, **49**, 630–641, 1993.
- [16] Oberley, L.W.: Free radicals and diabetes. *Free Radic. Biol. Med.*, **5**, 113–124, 1988.
- [17] Harman, D.: The aging process. *Proc. Natl. Acad. Sci. USA*, **78**, 7124–7128, 1981.
- [18] Itoh, M. and Guth, P.H.: Role of oxygen-derived free radicals in hemorrhagic shock-induced gastric lesions in the rat. *Gastroenterology*, **88**, 1162–1167, 1985.
- [19] Perry, M.A., Wadhwa, S., Parks, D.A., Pickard, W., and Granger, D.N.: Role of oxygen radicals in ischemia-induced lesions in the cat stomach. *Gastroenterology*, **90**, 362–367, 1986.
- [20] Parks, D.A., Bulkley, G.B., and Granger, D.N.: Role of oxygen-derived free radicals in digestive tract diseases. *Surgery*, **94**, 415–422, 1983.
- [21] Yoshikawa, T., Naito, Y., Tanigawa, T., Yoneta, T., Yasuda, M., Ueda, S., Oyamada, H., and Kondo, M.: Effect of zinc-carnosine chelate compound (Z-103), a novel antioxidant, on acute gastric mucosal injury induced by ischemia-reperfusion in rats. *Free Radic. Res. Commun.*, **14**, 289–296, 1991.
- [22] Parks, D.A., Bulkley, G.B., Granger, D.N., Hamilton, S.R., and McCord, J.M.: Ischemic injury in the cat small intestine: role of superoxide radicals. *Gastroenterology*, **82**, 9–15, 1982.
- [23] Schoenberg, M.H., Fredholm, B.B., Haglund, U., Jung, H., Sellin, D., Younes, M., and Schildberg, F.W.: Studies on the oxygen radical mechanism involved in the small intestinal reperfusion damage. *Acta. Physiol. Scand.*, **124**, 581–589, 1985.
- [24] Younes, M., Mohr, A., Schoenberg, M.H., and Schildberg, F.W.: Inhibition of lipid peroxidation by superoxide dismutase following regional intestinal ischemia and reperfusion. *Res. Exp. Med. (Berl.)*, **187**, 9–17, 1987.
- [25] Gibson, D.D., Brackett, D.J., Squires, R.A., Balla, A.K., Lerner, M.R., McCay, P.B., and Pennington, L.R.: Evidence that the large loss of glutathione observed in ischemia/reperfusion of the small intestine is not due to oxidation to glutathione disulfide. *Free Radic. Biol. Med.*, **14**, 427–433, 1993.
- [26] Turrens, J.F., Giulivi, C., Pinus, C., Roldan, E., Lavagno, C., and Boveris, A.: Low level chemiluminescence from isolated rat hepatocytes, intact lung and intestine *in situ*. *Basic Life Sci.*, **49**, 239–242, 1988.
- [27] Nilsson, U.A., Lundgren, O., Haglund, E., and Bylund-Fellenius, A.C.: Radical production during *in vivo* intestinal ischemia and reperfusion in the cat. *Am. J. Physiol.*, **257**,

- G409–G414, 1989.
- [28] Jaeschke, H. and Mitchell, J.R.: Mitochondria and xanthine oxidase both generate reactive oxygen species in isolated perfused rat liver after hypoxic injury. *Biochem. Biophys. Res. Commun.*, **160**, 140–147, 1989.
 - [29] Zhang, J. and Piantadosi, C.A.: Mitochondrial oxidative stress after carbon monoxide hypoxia in the rat brain. *J. Clin. Invest.*, **90**, 1193–1199, 1992.
 - [30] Parks, D.A., Williams, T.K., and Beckman, J.S.: Conversion of xanthine dehydrogenase to oxidase in ischemic rat intestine: a reevaluation. *Am. J. Physiol.*, **254**, G768–G774, 1988.
 - [31] Bindoli, A., Cavallini, L., Rigobello, M.P., Coassin, M., and Di Lisa, F.: Modification of the xanthine-converting enzyme of perfused rat heart during ischemia and oxidative stress. *Free Radic. Biol. Med.*, **4**, 163–167, 1988.
 - [32] Zulueta, J.J., Yu, F.S., Hertig, I.A., Thannickal, V.J., and Hassoun, P.M.: Release of hydrogen peroxide in response to hypoxia-reoxygenation: role of an NAD(P)H oxidase-like enzyme in endothelial cell plasma membrane. *Am. J. Respir. Cell Mol. Biol.*, **12**, 41–49, 1995.
 - [33] Liu, J.Q., Zelko, I.N., and Folz, R.J.: Reoxygenation-induced constriction in murine coronary arteries: the role of endothelial NADPH oxidase (gp91phox) and intracellular superoxide. *J. Biol. Chem.*, **279**, 24493–24497, 2004.
 - [34] Grisham, M.B. and Granger, D.N.: Metabolic sources of reactive oxygen metabolites during oxidant stress and ischemia with reperfusion. *Clin. Chest. Med.*, **10**, 71–81, 1989.
 - [35] Grisham, M.B., Hernandez, L.A., and Granger, D.N.: Xanthine oxidase and neutrophil infiltration in intestinal ischemia. *Am. J. Physiol.*, **251**, G567–G574, 1986.
 - [36] Grace, P.A.: Ischaemia-reperfusion injury. *Br. J. Surg.*, **81**, 637–647, 1994.
 - [37] Aust, S.D., Morehouse, L.A., and Thomas, C.E.: Role of metals in oxygen radical reactions. *J. Free Radic. Biol. Med.*, **1**, 3–25, 1985.
 - [38] Parks, D.A. and Granger, D.N.: Contributions of ischemia and reperfusion to mucosal lesion formation. *Am. J. Physiol.*, **250**, G749–G753, 1986.
 - [39] Granger, D.N.: Role of xanthine oxidase and granulocytes in ischemia-reperfusion injury. *Am. J. Physiol.*, **255**, H1269–H1275, 1988.
 - [40] Ates, B., Yilmaz, I., Geckil, H., Iraz, M., Birincioglu, M., and Fiskin, K.: Protective role of melatonin given either before ischemia or prior to reperfusion on intestinal ischemia-reperfusion damage. *J. Pineal. Res.*, **37**, 149–152, 2004.
 - [41] Granger, D.N., Rutili, G., and McCord, J.M.: Superoxide radicals in feline intestinal ischemia. *Gastroenterology*, **81**, 22–29, 1981.
 - [42] Parks, D.A. and Granger, D.N.: Xanthine oxidase: biochemistry, distribution and physiology. *Acta. Physiol. Scand. Suppl.*, **548**, 87–99, 1986.
 - [43] Harisch, G., Kretschmer, M., Richter, T., and Pickel, M.: Investigations on the influence of copper succinate on the production of superoxide anion radicals by bovine small intestinal mucosa cells. *Zentralbl. Veterinarmed A*, **36**, 576–584, 1989.
 - [44] Younes, M., Schoenberg, M.H., Jung, H., Fredholm, B.B., Haglund, U., and Schildberg, F.W.: Oxidative tissue damage following regional intestinal ischemia and reperfusion in the cat. *Res. Exp. Med. (Berl.)*, **184**, 259–264, 1984.
 - [45] Blum, H., Summers, J.J., Schnall, M.D., Barlow, C., Leigh, J.S., Jr., Chance, B., and Buzby, G.P.: Acute intestinal ischemia studies by phosphorus nuclear magnetic resonance spectroscopy. *Ann. Surg.*, **204**, 83–88, 1986.
 - [46] Menguy, R., Desbaillets, L., and Masters, Y.F.: Mechanism of stress ulcer: influence of hypovolemic shock on energy metabolism in the gastric mucosa. *Gastroenterology*, **66**, 46–55, 1974.
 - [47] Robinson, J.W., Mirkovitch, V., Winistorfer, B., and Saegesser, F.: Response of the intestinal mucosa to ischaemia. *Gut*, **22**, 512–527, 1981.
 - [48] Juergens, T.M. and Hardin, C.D.: Fructose-1,6-bisphosphate as a metabolic substrate in hog ileum smooth muscle during hypoxia. *Mol. Cell. Biochem.*, **154**, 83–93, 1996.
 - [49] Lazzarino, G., Tavazzi, B., Di Pierro, D., and Giardina, B.: Ischemia and reperfusion: effect of fructose-1,6-bisphosphate. *Free Radic. Res. Commun.*, **16**, 325–339, 1992.
 - [50] Akimitsu, T., White, J.A., Carden, D.L., Gute, D.C., and Korthuis, R.J.: Fructose-1,6-diphosphate or adenosine attenuate leukocyte adherence in postischemic skeletal muscle. *Am. J. Physiol.*, **269**, H1743–H1751, 1995.
 - [51] Sola, A., De Oca, J., Alfaro, V., Xaus, C., Jaurrieta, E., and Hotter, G.: Protective effects of exogenous fructose-1,6-bisphosphate during small bowel transplantation in rats. *Surgery*, **135**, 518–526, 2004.
 - [52] Simon, R.H., Scoggin, C.H., and Patterson, D.: Hydrogen peroxide causes the fatal injury to human fibroblasts exposed to oxygen radicals. *J. Biol. Chem.*, **256**, 7181–7186, 1981.
 - [53] Parks, D.A., Shah, A.K., and Granger, D.N.: Oxygen radicals: effects on intestinal vascular permeability. *Am. J. Physiol.*, **247**, G167–G170, 1984.
 - [54] Granger, D.N., Hollwarth, M.E., and Parks, D.A.: Ischemia-reperfusion injury: role of oxygen-derived free radicals. *Acta. Physiol. Scand. Suppl.*, **548**, 47–63, 1986.
 - [55] Takeyoshi, I., Zhang, S., Nakamura, K., Ikoma, A., Zhu, Y., Starzl, T.E., and Todo, S.: Effect of ischemia on the canine large bowel: a comparison with the small intestine. *J. Surg. Res.*, **62**, 41–48, 1996.
 - [56] Hinnebusch, B.F., Ma, Q., Henderson, J.W., Siddique, A., Archer, S.Y., and Hodin, R.A.: Enterocyte response to ischemia is dependent on differentiation state. *J. Gastrointest. Surg.*, **6**, 403–409, 2002.
 - [57] Kubes, P. and McCafferty, D.M.: Nitric oxide and intestinal inflammation. *Am. J. Med.*, **109**, 150–158, 2000.
 - [58] Kubes, P.: Ischemia-reperfusion in feline small intestine: a role for nitric oxide. *Am. J. Physiol.*, **264**, G143–G149, 1993.
 - [59] Naito, Y., Takagi, T., Ichikawa, H., Tomatsuri, N., Kuroda, M., Isozaki, Y., Katada, K., Uchiyama, K., Kokura, S., Yoshida, N., Okanoue, T., and Yoshikawa, T.: A novel potent inhibitor of inducible nitric oxide inhibitor, ONO-1714, reduces intestinal ischemia-reperfusion injury in rats. *Nitric Oxide*, **10**, 170–177, 2004.

- [60] Moore-Olufemi, S.D., Xue, H., Allen, S.J., Moore, F.A., Stewart, R.H., Laine, G.A., and Cox, C.S., Jr.: Inhibition of intestinal transit by resuscitation-induced gut edema is reversed by L-NIL. *J. Surg. Res.*, **129**, 1–5, 2005.
- [61] Kim, H. and Hwan Kim, K.: Role of nitric oxide and mucus in ischemia/reperfusion-induced gastric mucosal injury in rats. *Pharmacology*, **62**, 200–207, 2001.
- [62] Suzuki, Y., Deitch, E.A., Mishima, S., Lu, Q., and Xu, D.: Inducible nitric oxide synthase gene knockout mice have increased resistance to gut injury and bacterial translocation after an intestinal ischemia-reperfusion injury. *Crit. Care Med.*, **28**, 3692–3696, 2000.
- [63] Iwata, F., Joh, T., Yokoyama, Y., and Itoh, M.: Role of endogenous nitric oxide in ischaemia-reperfusion injury of rat gastric mucosa. *J. Gastroenterol. Hepatol.*, **13**, 997–1001, 1998.
- [64] Wada, K., Kamisaki, Y., Ohkura, T., Kanda, G., Nakamoto, K., Kishimoto, Y., Ashida, K., and Itoh, T.: Direct measurement of nitric oxide release in gastric mucosa during ischemia-reperfusion in rats. *Am. J. Physiol.*, **274**, G465–G471, 1998.
- [65] Wink, D.A., Hanbauer, I., Krishna, M.C., DeGraff, W., Gamson, J., and Mitchell, J.B.: Nitric oxide protects against cellular damage and cytotoxicity from reactive oxygen species. *Proc. Natl. Acad. Sci. USA*, **90**, 9813–9817, 1993.
- [66] Lefer, A.M. and Lefer, D.J.: The role of nitric oxide and cell adhesion molecules on the microcirculation in ischaemia-reperfusion. *Cardiovasc. Res.*, **32**, 743–751, 1996.
- [67] Kosonen, O., Kankaanranta, H., Malo-Ranta, U., and Moilanen, E.: Nitric oxide-releasing compounds inhibit neutrophil adhesion to endothelial cells. *Eur. J. Pharmacol.*, **382**, 111–117, 1999.
- [68] Gauthier, T.W., Davenpeck, K.L., and Lefer, A.M.: Nitric oxide attenuates leukocyte-endothelial interaction via P-selectin in splanchnic ischemia-reperfusion. *Am. J. Physiol.*, **267**, G562–G568, 1994.
- [69] Payne, D. and Kubes, P.: Nitric oxide donors reduce the rise in reperfusion-induced intestinal mucosal permeability. *Am. J. Physiol.*, **265**, G189–G195, 1993.
- [70] Andrews, F.J., Malcontenti-Wilson, C., and O'Brien, P.E.: Protection against gastric ischemia-reperfusion injury by nitric oxide generators. *Dig. Dis. Sci.*, **39**, 366–373, 1994.
- [71] Kawata, K., Takeyoshi, I., Iwanami, K., Sunose, Y., Aiba, M., Ohwada, S., Matsumoto, K., and Morishita, Y.: A spontaneous nitric oxide donor ameliorates small bowel ischemia-reperfusion injury in dogs. *Dig. Dis. Sci.*, **46**, 1748–1756, 2001.
- [72] Weiss, S.J.: Oxygen, ischemia and inflammation. *Acta. Physiol. Scand. Suppl.*, **548**, 9–37, 1986.
- [73] Granger, D.N. and Korthuis, R.J.: Physiologic mechanisms of postischemic tissue injury. *Annu. Rev. Physiol.*, **57**, 311–332, 1995.
- [74] Hernandez, L.A., Grisham, M.B., Twohig, B., Arfors, K.E., Harlan, J.M., and Granger, D.N.: Role of neutrophils in ischemia-reperfusion-induced microvascular injury. *Am. J. Physiol.*, **253**, H699–H703, 1987.
- [75] Granger, D.N., Benoit, J.N., Suzuki, M., and Grisham, M.B.: Leukocyte adherence to venular endothelium during ischemia-reperfusion. *Am. J. Physiol.*, **257**, G683–G688, 1989.
- [76] Cerqueira, N.F., Hussni, C.A., and Yoshida, W.B.: Pathophysiology of mesenteric ischemia/reperfusion: a review. *Acta Cir. Bras.*, **20**, 336–343, 2005.
- [77] Haglund, U.H., Morris, J.B., and Bulkley, G.B.: Haemodynamic characterization of the isolated (denervated) parabolically perfused rat jejunum. *Acta. Physiol. Scand.*, **132**, 151–158, 1988.
- [78] Bilbao, J., Garcia-Alonso, I., Portugal, V., Barcelo, P., Ortiz, J., and Mendez, J.: Therapeutic usefulness of antioxidant drugs in experimental intestinal reperfusion syndrome. *Rev. Esp. Enferm. Dig.*, **80**, 237–241, 1991.
- [79] Laudanno, O.M., Bedini, O.A., Cesolari, J.A., and San Miguel, P.: Sucralfate in the prevention of acute gastric lesions induced by ischemia-reperfusion. *Acta. Gastroenterol. Latinoam.*, **21**, 85–88, 1991.
- [80] Boros, M., Karacsony, G., Kaszaki, J., and Nagy, S.: Reperfusion mucosal damage after complete intestinal ischemia in the dog: the effects of antioxidant and phospholipase A2 inhibitor therapy. *Surgery*, **113**, 184–191, 1993.
- [81] Granger, D.N., McCord, J.M., Parks, D.A., and Hollwarth, M.E.: Xanthine oxidase inhibitors attenuate ischemia-induced vascular permeability changes in the cat intestine. *Gastroenterology*, **90**, 80–84, 1986.
- [82] Nakamoto, K., Kamisaki, Y., Wada, K., Kawasaki, H., and Itoh, T.: Protective effect of acetaminophen against acute gastric mucosal lesions induced by ischemia-reperfusion in the rat. *Pharmacology*, **54**, 203–210, 1997.
- [83] Horne, M.M., Pascoe, P.J., Ducharme, N.G., Barker, I.K., and Grovum, W.L.: Attempts to modify reperfusion injury of equine jejunal mucosa using dimethylsulfoxide, allopurinol, and intraluminal oxygen. *Vet. Surg.*, **23**, 241–249, 1994.
- [84] Alarcon de la Lastra, A.C., Martin, M.J., Motilva, V., Jimenez, M., La Casa, C., and Lopez, A.: Gastroprotection induced by silymarin, the hepatoprotective principle of Silybum marianum in ischemia-reperfusion mucosal injury: role of neutrophils. *Planta Med.*, **61**, 116–119, 1995.
- [85] Zollei, I.: Experimental study of hypovolaemic shock-induced gastric mucosal lesions in the rat. *Ann. Acad. Med. Singapore*, **28**, 85–89, 1999.
- [86] Hakguder, G., Akgur, F.M., Ates, O., Olguner, M., Aktug, T., and Ozer, E.: Short-term intestinal ischemia-reperfusion alters intestinal motility that can be preserved by xanthine oxidase inhibition. *Dig. Dis. Sci.*, **47**, 1279–1283, 2002.
- [87] Kulah, B., Besler, H.T., Akdag, M., Oruc, T., Altinok, G., Kulacoglu, H., Ozmen, M.M., and Coskun, F.: The effects of verapamil vs. allopurinol on intestinal ischemia/reperfusion injury in rats. "An experimental study". *Hepatogastroenterology*, **51**, 401–407, 2004.
- [88] Hei, Z.Q., Huang, H.Q., Zhang, J.J., Chen, B.X., and Li, X.Y.: Protective effect of Astragalus membranaceus on intestinal mucosa reperfusion injury after hemorrhagic shock in rats. *World J. Gastroenterol.*, **11**, 4986–4991, 2005.
- [89] Ozden, A., Tetik, C., Bilgihan, A., Calli, N., Bostanci, B., Yis, O., and Duzcan, E.: Antithrombin III prevents 60 min

- warm intestinal ischemia reperfusion injury in rats. *Res. Exp. Med. (Berl.)*, **198**, 237–246, 1999.
- [90] Ceran, C., Sonmez, K., Turkyilmaz, Z., Demirogullari, B., Dursun, A., Duzgun, E., Basaklar, A.C., and Kale, N.: Effect of bilirubin in ischemia/reperfusion injury on rat small intestine. *J. Pediatr. Surg.*, **36**, 1764–1767, 2001.
- [91] Hammerman, C., Goldschmidt, D., Caplan, M.S., Kaplan, M., Bromiker, R., Eidelman, A.I., Gartner, L.M., and Hochman, A.: Protective effect of bilirubin in ischemia-reperfusion injury in the rat intestine. *J. Pediatr. Gastroenterol. Nutr.*, **35**, 344–349, 2002.
- [92] Buyukgebiz, O., Aktan, A.O., Yegen, C., Yalcin, A.S., Haklar, G., Yalin, R., and Ercan, Z.S.: Captopril increases endothelin serum concentrations and preserves intestinal mucosa after mesenteric ischemia-reperfusion injury. *Res. Exp. Med. (Berl.)*, **194**, 339–348, 1994.
- [93] Kitano, M., Wada, K., Kamisaki, Y., Nakamoto, K., Kishimoto, Y., Kawasaki, H., and Itoh, T.: Effects of cimetidine on acute gastric mucosal injury induced by ischemia-reperfusion in rats. *Pharmacology*, **55**, 154–164, 1997.
- [94] Poussios, D., Andreadou, I., Papalois, A., Rekkas, E., Gavalakis, N., Aroni, K., Kourounakis, P.N., Fotiadis, C., and Sechas, M.N.: Protective effect of a novel antioxidant non-steroidal anti-inflammatory agent (compound IA) on intestinal viability after acute mesenteric ischemia and reperfusion. *Eur. J. Pharmacol.*, **465**, 275–280, 2003.
- [95] Yoshikawa, T., Takahashi, S., Naito, Y., Ueda, S., Tanigawa, T., Yoshida, N., and Kondo, M.: Effects of a platelet-activating factor antagonist, CV-6209, on gastric mucosal lesions induced by ischemia-reperfusion. *Lipids*, **27**, 1058–1060, 1992.
- [96] Wada, K., Kamisaki, Y., Kitano, M., Nakamoto, K., and Itoh, T.: Protective effect of cystathionine on acute gastric mucosal injury induced by ischemia-reperfusion in rats. *Eur. J. Pharmacol.*, **294**, 377–382, 1995.
- [97] Puglisi, R.N., Strande, L., Santos, M., Schulte, G., Hewitt, C.W., and Whalen, T.V.: Beneficial effects of cyclosporine and rapamycin in small bowel ischemic injury. *J. Surg. Res.*, **65**, 115–118, 1996.
- [98] Puglisi, R.N., Strande, L., Santos, M., Doolin, E.J., Hewitt, C.W., and Whalen, T.V.: The effect of cyclosporine in gut ischemic injury: a computerized morphometric and enzymatic analysis. *J. Pediatr. Surg.*, **31**, 319–322, 1996.
- [99] Andrews, F.J., Malcontenti, C., and O'Brien, P.E.: Sequence of gastric mucosal injury following ischemia and reperfusion. Role of reactive oxygen metabolites. *Dig. Dis. Sci.*, **37**, 1356–1361, 1992.
- [100] Kurokawa, T., Joh, T., Ikai, M., Seno, K., Yokoyama, Y., and Itoh, M.: Rebamipide protects against oxygen radical-mediated gastric mucosal injury in rats. *Dig. Dis. Sci.*, **43**, 113S–117S, 1998.
- [101] Balogh, N., Krausz, F., Levai, P., Ribiczeyne, P.S., Vajdovich, P., and Gaal, T.: Effect of deferoxamine and L-arginine treatment on lipid peroxidation in an intestinal ischaemia-reperfusion model in rats. *Acta. Vet. Hung.*, **50**, 343–356, 2002.
- [102] Ali, A.T., al-Swayeh, O.A., al-Humayyd, M.S., Mustafa, A.A., al-Rashed, R.S., and al-Tuwaijiri, A.S.: Natural honey prevents ischaemia-reperfusion-induced gastric mucosal lesions and increased vascular permeability in rats. *Eur. J. Gastroenterol. Hepatol.*, **9**, 1101–1107, 1997.
- [103] Kojima, M., Iwakiri, R., Wu, B., Fujise, T., Watanabe, K., Lin, T., Amemori, S., Sakata, H., Shimoda, R., Oguzu, T., Ootani, A., Tsunada, S., and Fujimoto, K.: Effects of antioxidative agents on apoptosis induced by ischaemia-reperfusion in rat intestinal mucosa. *Aliment. Pharmacol. Ther.*, **18** Suppl 1, 139–145, 2003.
- [104] Dabareiner, R.M., White, N.A., Snyder, J.R., Feldman, B.F., and Donaldson, L.L.: Effects of Carolina rinse solution, dimethyl sulfoxide, and the 21-aminosteroid, U-74389G, on microvascular permeability and morphology of the equine jejunum after low-flow ischemia and reperfusion. *Am. J. Vet. Res.*, **66**, 525–536, 2005.
- [105] Tomatsuri, N., Yoshida, N., Takagi, T., Katada, K., Isozaki, Y., Imamoto, E., Uchiyama, K., Kokura, S., Ichikawa, H., Naito, Y., Okanoue, T., and Yoshikawa, T.: Edaravone, a newly developed radical scavenger, protects against ischemia-reperfusion injury of the small intestine in rats. *Int. J. Mol. Med.*, **13**, 105–109, 2004.
- [106] Berlanga, J., Prats, P., Ramirez, D., Gonzalez, R., Lopez-Saura, P., Aguiar, J., Ojeda, M., Boyle, J.J., Fitzgerald, A.J., and Playford, R.J.: Prophylactic use of epidermal growth factor reduces ischemia/reperfusion intestinal damage. *Am. J. Pathol.*, **161**, 373–379, 2002.
- [107] Villa, X., Kuluz, J.W., Schleien, C.L., and Thompson, J.F.: Epidermal growth factor reduces ischemia-reperfusion injury in rat small intestine. *Crit. Care. Med.*, **30**, 1576–1580, 2002.
- [108] Martin, A.E., Luquette, M.H., and Besner, G.E.: Timing, route, and dose of administration of heparin-binding epidermal growth factor-like growth factor in protection against intestinal ischemia-reperfusion injury. *J. Pediatr. Surg.*, **40**, 1741–1747, 2005.
- [109] Iino, T., Tashima, K., Umeda, M., Ogawa, Y., Takada, M., Takata, K., and Takeuchi, K.: Effect of ellagic acid on gastric damage induced in ischemic rat stomachs following ammonia or reperfusion. *Life Sci.*, **70**, 1139–1150, 2002.
- [110] Lai, H.S., Chen, Y., Chen, W.J., Chang, J., and Chiang, L.Y.: Free radical scavenging activity of fullerene on grafts after small bowel transplantation in dogs. *Transplant. Proc.*, **32**, 1272–1274, 2000.
- [111] Stein, H.J., Hinder, R.A., and Oosthuizen, M.M.: Gastric mucosal injury caused by hemorrhagic shock and reperfusion: protective role of the antioxidant glutathione. *Surgery*, **108**, 467–473; discussion 473–464, 1990.
- [112] Harward, T.R., Coe, D., Souba, W.W., Klingman, N., and Seeger, J.M.: Glutamine preserves gut glutathione levels during intestinal ischemia/reperfusion. *J. Surg. Res.*, **56**, 351–355, 1994.
- [113] Basoglu, M., Yildiran, I., Akcay, F., Kiziltunc, A., Kavak, I., and Oren, D.: Glutathione and nitric oxide concentrations in glutamine-infused rabbits with intestinal ischaemia/reperfusion. *Eur. J. Clin. Chem. Clin. Biochem.*, **35**, 415–419, 1997.
- [114] Koltuksuz, U., Ozen, S., Uz, E., Aydin, M., Karaman, A.,

- Gultek, A., Akyol, O., Gursoy, M.H., and Aydin, E.: Caffeic acid phenethyl ester prevents intestinal reperfusion injury in rats. *J. Pediatr. Surg.*, **34**, 1458–1462, 1999.
- [115] Naito, Y., Yoshikawa, T., Matsuyama, K., Yagi, N., Arai, M., Nakamura, Y., Kaneko, T., Yoshida, N., and Kondo, M.: Effect of a novel histamine H2 receptor antagonist, IT-066, on acute gastric injury induced by ischemia-reperfusion in rats, and its antioxidative properties. *Eur. J. Pharmacol.*, **294**, 47–54, 1995.
- [116] Gunel, E., Caglayan, F., Caglayan, O., Dilsiz, A., Duman, S., and Aktan, M.: Treatment of intestinal reperfusion injury using antioxidative agents. *J. Pediatr. Surg.*, **33**, 1536–1539, 1998.
- [117] Byrka-Owczarek, K., Stepievska-Mazur, K., Krason, M., Bohosiewicz, J., Koszutski, T., and Wojtynek, G.: The evaluation of the protective action of antioxidants on small intestine of rabbits experimentally injured by ischemia and reperfusion. *J. Pediatr. Surg.*, **39**, 1226–1229, 2004.
- [118] Konturek, P.C., Konturek, S.J., Majka, J., Zembala, M., and Hahn, E.G.: Melatonin affords protection against gastric lesions induced by ischaemia-reperfusion possibly due to its antioxidant and mucosal microcirculatory effects. *Eur. J. Pharmacol.*, **322**, 73–77, 1997.
- [119] Konturek, P.C., Konturek, S.J., Brzozowski, T., Dembinski, A., Zembala, M., Mytar, B., and Hahn, E.G.: Gastroprotective activity of melatonin and its precursor, L-tryptophan, against stress-induced and ischaemia-induced lesions is mediated by scavenge of oxygen radicals. *Scand. J. Gastroenterol.*, **32**, 433–438, 1997.
- [120] Brzozowski, T., Konturek, P.C., Konturek, S.J., Pajdo, R., Bielanski, W., Brzozowska, I., Stachura, J., and Hahn, E.G.: The role of melatonin and L-tryptophan in prevention of acute gastric lesions induced by stress, ethanol, ischemia, and aspirin. *J. Pineal. Res.*, **23**, 79–89, 1997.
- [121] Kazez, A., Demirbag, M., Ustundag, B., Ozercan, I.H., and Saglam, M.: The role of melatonin in prevention of intestinal ischemia-reperfusion injury in rats. *J. Pediatr. Surg.*, **35**, 1444–1448, 2000.
- [122] Ustundag, B., Kazez, A., Demirbag, M., Canatan, H., Halifeoglu, I., and Ozercan, I.H.: Protective effect of melatonin on antioxidative system in experimental ischemia-reperfusion of rat small intestine. *Cell Physiol. Biochem.*, **10**, 229–236, 2000.
- [123] Cabeza, J., Motilva, V., Martin, M.J., and de la Lastra, C.A.: Mechanisms involved in gastric protection of melatonin against oxidant stress by ischemia-reperfusion in rats. *Life Sci.*, **68**, 1405–1415, 2001.
- [124] Cabeza, J., Alarcon-de-la-Lastra, C., Jimenez, D., Martin, M.J., and Motilva, V.: Melatonin modulates the effects of gastric injury in rats: role of prostaglandins and nitric oxide. *Neurosignals*, **12**, 71–77, 2003.
- [125] Bulbuler, N., Akkus, M. A., Ilhan, Y. S., Baysal, F., Ozercan, I., Aygen, E., and Kirkil, C.: The effects of L-tryptophan and pentoxifylline on stress ulcer. *Ulus. Travma. Acil. Cerrahi. Derg.*, **9**, 90–95, 2003.
- [126] Sileri, P., Sica, G.S., Gentileschi, P., Venza, M., Benavoli, D., Jarzembowski, T., Manzelli, A., and Gaspari, A.L.: Melatonin reduces bacterial translocation after intestinal ischemia-reperfusion injury. *Transplant. Proc.*, **36**, 2944–2946, 2004.
- [127] Ozacmak, V.H., Sayan, H., Arslan, S.O., Altaner, S., and Aktas, R.G.: Protective effect of melatonin on contractile activity and oxidative injury induced by ischemia and reperfusion of rat ileum. *Life Sci.*, **76**, 1575–1588, 2005.
- [128] Cuzzocrea, S., Mazzon, E., Costantino, G., Serrano, I., De Sarro, A., and Caputi, A.P.: Effects of n-acetylcysteine in a rat model of ischemia and reperfusion injury. *Cardiovasc. Res.*, **47**, 537–548, 2000.
- [129] Sun, Z., Olanders, K., Lasson, A., Dib, M., Annborn, M., Andersson, K., Wang, X., and Andersson, R.: Effective treatment of gut barrier dysfunction using an antioxidant, a PAF inhibitor, and monoclonal antibodies against the adhesion molecule PECAM-1. *J. Surg. Res.*, **105**, 220–233, 2002.
- [130] Montero, E.F., Abrahao, M.S., Koike, M.K., Manna, M.C., and Ramalho, C.E.: Intestinal ischemia and reperfusion injury in growing rats: hypothermia and N-acetylcysteine modulation. *Microsurgery*, **23**, 517–521, 2003.
- [131] Olanders, K., Borjesson, A., Sun, Z.W., and Andersson, R.: Protective effects of N-acetyl-L-cysteine and platelet activating factor inhibition are not linked to intercellular adhesion molecule-1 expression after intestinal ischemia and reperfusion injury in rats. *Scand. J. Gastroenterol.*, **38**, 618–625, 2003.
- [132] Dun, Y., Hao, Y.B., Wu, Y.X., Zhang, Y., and Zhao, R.R.: Protective effects of nitroglycerin-induced preconditioning mediated by calcitonin gene-related peptide in rat small intestine. *Eur. J. Pharmacol.*, **430**, 317–324, 2001.
- [133] Khanna, A., Rossman, J., Caty, M.G., and Fung, H.L.: Beneficial effects of intraluminal nitroglycerin in intestinal ischemia-reperfusion injury in rats. *J. Surg. Res.*, **114**, 15–24, 2003.
- [134] Mallick, I.H., Yang, W.X., Winslet, M.C., and Seifalian, A.M.: Pyrrolidine dithiocarbamate reduces ischemia-reperfusion injury of the small intestine. *World J. Gastroenterol.*, **11**, 7308–7313, 2005.
- [135] El Eter, E., Hagar, H.H., Al-Tuwaijiri, A., and Arafa, M.: Nuclear factor-kappaB inhibition by pyrrolidine dithiocarbamate attenuates gastric ischemia-reperfusion injury in rats. *Can. J. Physiol. Pharmacol.*, **83**, 483–492, 2005.
- [136] Cicalese, L., Lee, K., Schraut, W., Watkins, S., Borle, A., and Stanko, R.: Pyruvate prevents ischemia-reperfusion mucosal injury of rat small intestine. *Am. J. Surg.*, **171**, 97–100; discussion 100–101, 1996.
- [137] Kim, C.D. and Hong, K.W.: Preventive effect of rebamipide on gastric lesions induced by ischemia-reperfusion in the rat. *J. Pharmacol. Exp. Ther.*, **275**, 340–344, 1995.
- [138] Hiratsuka, T., Futagami, S., Tatsuguchi, A., Suzuki, K., Shinji, Y., Kusunoki, M., Shinoki, K., Nishigaki, H., Fujimori, S., Wada, K., Miyake, K., Gudis, K., Tsukui, T., and Sakamoto, C.: COX-1 and COX-2 conversely promote and suppress ischemia-reperfusion gastric injury in mice. *Scand. J. Gastroenterol.*, **40**, 903–913, 2005.
- [139] Ichikawa, H., Takagi, T., Uchiyama, K., Higashihara, H.,

- Katada, K., Isozaki, Y., Naito, Y., Yoshida, N., and Yoshikawa, T.: Rotenone, a mitochondrial electron transport inhibitor, ameliorates ischemia-reperfusion-induced intestinal mucosal damage in rats. *Redox Rep.*, **9**, 313–316, 2004.
- [140] Yoshida, W.B., Alasio, T., Mazziotta, R., Qin, F., Kashani, M., Lee, S., Dardik, H., and Becker, R.: Effect of alpha-tocopherol, taurine and selenium on the attenuation of ischemia/reperfusion injury of splanchnic organs, *Cardiovasc. Surg.*, **6**, 178–187, 1998.
- [141] Ozturk, C., Avlan, D., Cinel, I., Cinel, L., Unlu, A., Camdeviren, H., Atik, U., and Oral, U.: Selenium pretreatment prevents bacterial translocation in rat intestinal ischemia/reperfusion model. *Pharmacol. Res.*, **46**, 171–175, 2002.
- [142] Riaz, A.A., Wan, M.X., Schafer, T., Dawson, P., Menger, M.D., Jeppsson, B., and Thorlacius, H.: Allopurinol and superoxide dismutase protect against leucocyte-endothelium interactions in a novel model of colonic ischaemia-reperfusion. *Br. J. Surg.*, **89**, 1572–1580, 2002.
- [143] Yoshikawa, T., Nakamura, S., Takahashi, S., Naito, Y., and Kondo, M.: Effect of sofalcone on gastric mucosal injury induced by ischemia-reperfusion and its antioxidant properties. *J. Clin. Gastroenterol.*, **17**, S111–S115, 1993.
- [144] Momo, K., Hoshina, K., Ishibashi, Y., and Saito, T.: Preventive effects of troxipide on a newly developed model of acute gastric mucosal lesion (AGML) induced by ischemia/reperfusion plus ammonia in the rat. *Nippon Yakurigaku Zasshi.*, **104**, 313–323, 1994.
- [145] al-Swayeh, O.A., al-Humayyd, M.S., Mustafa, A.A., al-Tuwaijri, A.S., al-Rashed, R.S., and Ali, A.T.: Sucralfate attenuates gastric mucosal lesions and increased vascular permeability induced by ischaemia and reperfusion in rats. *J. Gastroenterol. Hepatol.*, **12**, 481–489, 1997.
- [146] Wada, K., Kamisaki, Y., Kitano, M., Kishimoto, Y., Nakamoto, K., and Itoh, T.: Effects of sucralfate on acute gastric mucosal injury and gastric ulcer induced by ischemia-reperfusion in rats. *Pharmacology.*, **54**, 57–63, 1997.
- [147] Mojzis, J., Hegedusova, R., and Mirossay, L.: Role of mucus in ischemia/reperfusion-induced gastric mucosal injury in rats. *Physiol. Res.*, **49**, 441–446, 2000.
- [148] Sencan, A., Yilmaz, O., Ozer, E., Gunsar, C., Genc, K., Ulukus, C., Taneli, C., and Mir, E.: Does sucralfate prevent apoptosis occurring in the ischemia/reperfusion-induced intestinal injury? *Eur. J. Pediatr. Surg.*, **13**, 231–235, 2003.
- [149] Seno, K., Joh, T., Yokoyama, Y., and Itoh, M.: Role of endogenous acid in gastric mucosal injury induced by local ischemia-reperfusion in the rat. *J. Clin. Gastroenterol.*, **21**, S108–S112, 1995.
- [150] Naito, Y., Yoshikawa, T., Matsuyama, K., Yagi, N., Nakamura, Y., Nishimura, S., Kaneko, T., Yoshida, N., and Kondo, M.: Effect of the histamine H2-receptor antagonist (+/-)-(E)-1-[2-hydroxy-2-(4-hydroxyphenyl)ethyl]-3'-[2-[[[5-methylamino)methyl-2-furyl] methyl]thio]ethyl]-2'-(methylsulfonyl)guanidine on acute gastric mucosal injury in rats and its free-radical scavenging activities. *Arzneimittelforschung*, **47**, 845–848, 1997.
- [151] Udassin, R., Haskel, Y., and Samuni, A.: Nitroxide radical attenuates ischaemia/reperfusion injury to the rat small intestine. *Gut*, **42**, 623–627, 1998.
- [152] Tsimoyiannis, E.C., Lekkas, E.T., Paizis, J.B., Boullis, S.A., Page, P., and Kotoulas, O.B.: Prevention of peritoneal adhesions in rats with trimetazidine. *Acta. Chir. Scand.*, **156**, 771–774, 1990.
- [153] Tetik, C., Ozden, A., Calli, N., Bilgihan, A., Bostanci, B., Yis, O., and Bayramoglu, H.: Cytoprotective effect of trimetazidine on 60 minutes of intestinal ischemia-reperfusion injury in rats. *Transpl. Int.*, **12**, 108–112, 1999.
- [154] Sato, N., Kawano, S., Tsuji, S., and Kamada, T.: Microvascular basis of gastric mucosal protection. *J. Clin. Gastroenterol.*, **10**, S13–S18, 1988.
- [155] Mocan, H., Gedik, Y., Erduran, E., Mocan, Z.M., Okten, A., Gacar, N., and Pul, N.: The role of calcium channel entry blocker in experimental ischemia-reperfusion-induced intestinal injury. *Pol. J. Pharmacol.*, **47**, 179–183, 1995.
- [156] al-Dohayan, A.D. and al-Tuwaijri, A.S.: The potential synergistic effect of calcium channel blockers and alpha-tocopherol on gastric mucosal injury induced by ischaemia-reperfusion. *Eur. J. Gastroenterol. Hepatol.*, **8**, 1107–1110, 1996.
- [157] Ekman, T., Risberg, B., and Bagge, U.: Ascorbate reduces gastric bleeding after hemorrhagic shock and retransfusion in rats. *Eur. Surg. Res.*, **26**, 187–193, 1994.
- [158] Ekman, T., Risberg, B., Bagge, U., and Braide, M.: Blocking of endothelial-leukocyte interaction (rolling) does not improve reflow in the rat gastric mucosa after hemorrhagic shock and retransfusion. *Shock*, **2**, 257–261, 1994.
- [159] Nakamura, M., Ozaki, M., Fuchinoue, S., Teraoka, S., and Ota, K.: Ascorbic acid prevents ischemia-reperfusion injury in the rat small intestine. *Transpl. Int.*, **10**, 89–95, 1997.
- [160] Hazinedaroglu, S.M., Dulger, F., Kayaoglu, H.A., Pehlivan, M., Serinsoz, E., Canbolat, O., and Erverdi, N.: N-acetylcysteine in intestinal reperfusion injury: an experimental study in rats. *ANZ J. Surg.*, **74**, 676–678, 2004.
- [161] Seno, K., Joh, T., Yokoyama, Y., and Itoh, M.: Role of mucus in gastric mucosal injury induced by local ischemia/reperfusion. *J. Lab. Clin. Med.*, **126**, 287–293, 1995.